

Global Initiative for Chronic Obstructive Lung Disease



**GLOBAL STRATEGY FOR THE DIAGNOSIS,
MANAGEMENT, AND PREVENTION OF
CHRONIC OBSTRUCTIVE PULMONARY DISEASE**

UPDATED 2016

GLOBAL INITIATIVE FOR
CHRONIC OBSTRUCTIVE LUNG DISEASE

**GLOBAL STRATEGY FOR THE DIAGNOSIS, MANAGEMENT, AND
PREVENTION OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE
(UPDATED 2016)**



GLOBAL STRATEGY FOR THE DIAGNOSIS, MANAGEMENT, AND PREVENTION OF COPD (UPDATED 2016)

GOLD BOARD OF DIRECTORS (2015)

Marc Decramer, MD, Chair
University of Leuven
Leuven, Belgium

Alvar G. Agustí, MD
Hospital Clinic, Universitat de Barcelona
Barcelona Spain

Jean Bourbeau, MD
McGill University Health Centre
Montreal, Quebec, Canada

Bartolome R. Celli, MD
Brigham and Women's Hospital
Boston, Massachusetts USA

Rongchang Chen, MD
Guangzhou Institute of Respiratory Disease
Guangzhou, PRC

Gerard Criner, MD
Temple University School of Medicine
Philadelphia, Pennsylvania USA

Peter Frith, MD
Repatriation General Hospital, Adelaide
South Australia, Australia

David Halpin, MD
Royal Devon and Exeter Hospital
Devon, UK

M. Victorina López Varela, MD
Universidad de la República
Montevideo, Uruguay

Masaharu Nishimura, MD
Hokkaido Univ School of Medicine
Sapporo, Japan

Claus Vogelmeier, MD
University of Gießen and Marburg
Marburg, Germany

GOLD SCIENCE DIRECTOR

Suzanne S. Hurd, PhD
Vancouver, Washington, USA

GOLD SCIENCE COMMITTEE* (2015)

Claus Vogelmeier, MD
University of Giessen and Marburg
Marburg, Germany

Alvar G. Agustí, MD
Thorax Institute, Hospital Clinic
Univ. Barcelona, Ciberes, *Barcelona, Spain*

Antonio Anzueto, MD
University of Texas Health Science Center
San Antonio, Texas, USA

Marc Decramer, MD
Katholieke Universiteit Leuven
Leuven, Belgium

Leonardo M. Fabbri, MD
University of Modena & Reggio Emilia
Modena, Italy

Fernando Martinez, MD
University of Michigan School of Medicine
Ann Arbor, Michigan, USA

Nicolas Roche, MD
Hôpital Cochin
Paris, France

Roberto Rodriguez-Roisin, MD
Thorax Institute, Hospital Clinic
Univ. Barcelona, *Barcelona, Spain*

Donald Sin, MD
St. Paul's Hospital
Vancouver, Canada

Dave Singh, MD
University of Manchester
Manchester, UK

Robert Stockley, MD
University Hospital
Birmingham, UK

Jørgen Vestbo, MD
Hvidovre University Hospital, Hvidovre, Denmark
and University of Manchester
Manchester, England, UK

Jadwiga A. Wedzicha, MD
Univ College London
London, UK

*Disclosure forms for GOLD Committees are posted on the GOLD Website, www.goldcopd.org

GLOBAL STRATEGY FOR THE DIAGNOSIS, MANAGEMENT, AND PREVENTION OF COPD (REVISED 2011)

INVITED REVIEWERS

Joan-Albert Barbera, MD
Hospital Clinic, Universitat de Barcelona
Barcelona Spain

A. Sonia Buist, MD
Oregon Health Sciences University
Portland, OR, USA

Peter Calverley, MD
University Hospital Aintree
Liverpool, England, UK

Bart Celli, MD
Brigham and Women's Hospital
Boston, MA, USA

M. W. Elliott, MD
St. James's University Hospital
Leeds, England, UK

Yoshinosuke Fukuchi, MD
Juntendo University
Tokyo, Japan

Masakazu Ichinose, MD
Wakayama Medical University
Kimiedera, Wakayama, Japan

Christine Jenkins, MD
Woolcock Institute of Medical Research
Camperdown, NSW, Australia

H. A. M. Kerstjens, MD
University of Groningen
Groningen, The Netherlands

Peter Lange, MD
Hvidovre University Hospital
Copenhagen, Denmark

M. Victoria López Varela, MD
Universidad de la República
Montevideo, Uruguay

Maria Montes de Oca, MD
Hospital Universitario de Caracas
Caracas, Venezuela

Atsushi Nagai, MD
Tokyo Women's Medical University
Tokyo, Japan

Dennis Niewoehner, MD
Veterans Affairs Medical Center
Minneapolis, MN, USA

David Price, MD
University of Aberdeen
Aberdeen, Scotland, UK

Nicolas Roche, MD, PhD
University Paris Descartes
Paris, France

Sanjay Sethi, MD
State University of New York
Buffalo, NY, USA

GOLD NATIONAL LEADERS (Submitting Comments)

Lorenzo Corbetta, MD
University of Florence
Florence, Italy

Alexandru Corlateanu, MD, PhD
State Medical and Pharmaceutical University
Republic of Moldova

Le Thi Tuyet Lan, MD, PhD
University of Pharmacy and Medicine
Ho Chi Minh City, Vietnam

Fernando Lundgren, MD
Pernambuco, Brazil

E. M. Irušen, MD
University of Stellenbosch
South Africa

Timothy J. MacDonald, MD
St. Vincent's University Hospital
Dublin, Ireland

Takahide Nagase, MD
University of Tokyo
Tokyo, Japan

Ewa Nizankowska-Mogilnicka, MD, PhD
Jagiellonian University Medical College
Krakow, Poland

Magvannorov Oyunchimeg, MD
Ulaanbatar, Mongolia

Mostafizur Rahman, MD
NIDCH
Mohakhali, Dhaka, Bangladesh

PREFACE

In 2011, the Global Initiative for Chronic Obstructive Lung Disease (GOLD) released a consensus report, Global Strategy for the Diagnosis, Management, and Prevention of COPD. It recommended a major revision in the management strategy for COPD that was presented in the original 2001 document. Updated reports released in January 2013, January 2014, and January 2015 are based on scientific literature published since the completion of the 2011 document but maintain the same treatment paradigm. Assessment of COPD is based on the patient's level of symptoms, future risk of exacerbations, the severity of the spirometric abnormality, and the identification of comorbidities. The 2015 update adds an Appendix on Asthma COPD Overlap Syndrome, material prepared jointly by the GOLD and GINA Science Committees.


The GOLD report is presented as a "strategy document" for health care professionals to use as a tool to implement effective management programs based on available health care systems. The quadrant management strategy tool presented in this report is designed to be used in any clinical setting; it draws together a measure of the impact of the patient's symptoms and an assessment of the patient's risk of having a serious adverse health event in the future. Many studies have assessed the utility/relevance of this new tool; the main observations of these studies are shown in the table. Evidence will continue to be evaluated by the GOLD committees and management strategy recommendations modified as required.

Table: Summary Observations	Refs.
Choice of symptom measure (mMRC vs. CAT) influence category assignment	2-5
The prevalence of the four GOLD groups depends on the specific population studied, C being consistently the least prevalent	2;4-10
Groups differed in several clinical, functional, imaging and biological characteristics in addition to those used for their definition, including comorbidities	4;11;12
Prevalence of comorbidities and persistent systemic inflammation were highest in group B.	11
The new classification systems correlates with exercise capacity	5
A and D groups were relatively stable over time, where as groups B and C showed more temporal variability	11
Good prediction of exacerbations during follow-up	13
Conflicting results in relation to its capacity to predict mortality	5-7;14
B patients consistently have a mortality and hospitalization rate similar to C patients	11;13
Prescription appropriateness by GPs (in Italy) is better using new GOLD classification.	9
A real world observational study in five European countries and US identifies the frequent and potentially inappropriate use of inhaled steroids and bronchodilators in patients at low risk of exacerbations (A and B)	10

GOLD has been fortunate to have a network of international distinguished health professionals from multiple disciplines. Many of these experts have initiated investigations of the causes and prevalence of COPD in their countries, and have developed innovative approaches for the dissemination and implementation of the GOLD management strategy. The GOLD initiative will continue to work with National Leaders and other interested health care professionals to bring COPD to the attention of governments, public health officials, health care workers, and the general public to raise awareness of the burden of COPD and to develop programs for early detection, prevention and approaches to management.



Marc Decramer, MD
 Chair, GOLD Board of Directors
 Professor of Medicine
 CEO University Hospital Leuven
 University of Leuven, Leuven, Belgium



Claus Vogelmeier, MD
 Chair GOLD Science Committee
 Director, Internal Medicine Clinic
 University of Gießen and Marburg,
 School of Medicine
 Standort Marburg Baldingerstraße
 D-35043 Marburg Germany

REFERENCES

- 1: Agusti A, Hurd S, Jones P, Fabbri LM, Martinez F, Vogelmeier C et al. FAQs about the GOLD 2011 assessment proposal of COPD: a comparative analysis of four different cohorts. *Eur Respir J* 2013 November;42(5):1391-401.
- 2: Han MK, Mullerova H, Curran-Everett D, Dransfield DT, Washko GR, Regan EA et al. GOLD 2011 disease severity classification in COPD Gene: a prospective cohort study. *The Lancet Respir Med* 2013;1:43-50.
- 3: Jones PW, Adamek L, Nadeau G, Banik N. Comparisons of health status scores with MRC grades in COPD: implications for the GOLD 2011 classification. *Eur Respir J* 2013 September;42(3):647-54.
- 4: Jones PW, Nadeau G, Small M, Adamek L. Characteristics of a COPD population categorised using the GOLD framework by health status and exacerbations. *Respir Med* 2014 January;108(1):129-35.
- 5: Nishimura K, Oga T, Tsukino M, Hajiro T, Ikeda A, Jones PW. Reanalysis of the Japanese experience using the combined COPD assessment of the 2011 GOLD classification. *Respir Investig* 2014 March;52(2):129-35.
- 6: Soriano JB, Alfajeme I, Almagro P, Casanova C, Esteban C, Soler-Cataluna JJ et al. Distribution and prognostic validity of the new GOLD grading classification. *Chest* 2012;143(3):694-702.
- 7: Leivseth L, Brumpton BM, Nilsen TI, Mai XM, Johnsen R, Langhammer A. GOLD classifications and mortality in chronic obstructive pulmonary disease: the HUNT Study, Norway. *Thorax* 2013 October;68(10):914-21.
- 8: Haughney J, Gruffydd-Jones K, Roberts J, Lee AJ, Hardwell A, McGarvey L. The distribution of COPD in UK general practice using the new GOLD classification. *Eur Respir J* 2014 April;43(4):993-1002.
- 9: Maio S, Baldacci S, Martini F, Cerrai S, Sarno G, Borbotti M et al. COPD management according to old and new GOLD guidelines: an observational study with Italian general practitioners. *Curr Med Res Opin* 2014 June;30(6):1033-42.
- 10: Vestbo J, Vogelmeier C, Small M, Higgins V. Understanding the GOLD 2011 Strategy as applied to a real-world COPD population. *Respir Med* 2014 May;108(5):729-36.
- 11: Agusti A, Edwards LD, Celli B, Macnee W, Calverley PM, Mullerova H et al. Characteristics, stability and outcomes of the 2011 GOLD COPD groups in the ECLIPSE cohort. *Eur Respir J* 2013 September;42(3):636-46.
- 12: Sillen MJ, Franssen FM, Delbressine JM, Uszko-Lencer NH, Vanfleteren LE, Rutten EP et al. Heterogeneity in clinical characteristics and co-morbidities in dyspneic individuals with COPD GOLD D: findings of the DICES trial. *Respir Med* 2013 August;107(8):1186-94.
- 13: Lange P, Marott JL, Vestbo J, Olsen KR, Ingebrigtsen TS, Dahl M et al. Prediction of the clinical course of chronic obstructive pulmonary disease, using the new GOLD classification: a study of the general population. *Am J Respir Crit Care Med* 2012 November 15;186(10):975-81.
- 14: de Torres JP, Casanova C, Marin JM, Pinto-Plata V, Divo M, Zulueta JJ et al. Prognostic evaluation of COPD patients: GOLD 2011 versus BODE and the COPD comorbidity index COTE. *Thorax* 2014 September;69(9):799-804.

TABLE OF CONTENTS

Preface	iv	3. Therapeutic Options	19
Methodology and Summary of New Recommendations	ix	Key Points.....	20
Introduction	xiv	Smoking Cessation.....	20
1. Definition and Overview	1	Pharmacotherapies for Smoking Cessation.....	20
Key Points.....	2	Pharmacologic Therapy for Stable COPD.....	21
Definition.....	2	Overview of the Medications.....	21
Burden Of COPD.....	2	<i>Bronchodilators</i>	21
Prevalence.....	3	<i>Corticosteroids</i>	24
Morbidity.....	3	<i>Phosphodiesterase-4 Inhibitors</i>	25
Mortality.....	3	<i>Other Pharmacologic Treatments</i>	25
Economic Burden.....	3	Non-Pharmacologic Therapies.....	26
Social Burden.....	4	Rehabilitation.....	26
Factors That Influence Disease.....		Components of Pulmonary Rehabilitation Programs.....	27
Development And Progression.....	4	Other Treatments.....	28
Genes.....	4	Oxygen Therapy.....	28
Age and Gender.....	4	Ventilatory Support.....	29
Lung Growth and Development.....	4	Surgical Treatments.....	29
Exposure to Particles.....	5	Palliative Care, End-of-life Care, Hospice Care.....	30
Socioeconomic Status.....	5	4. Management of Stable COPD	31
Asthma/Bronchial Hyperreactivity.....	5	Key Points.....	32
Chronic Bronchitis.....	5	Introduction.....	32
Infections.....	5	Identify And Reduce Exposure to Risk Factors.....	33
Pathology, Pathogenesis And Pathophysiology.....	6	Tobacco Smoke.....	33
Pathology.....	6	Occupational Exposures.....	33
Pathogenesis.....	6	Indoor And Outdoor Pollution.....	33
Pathophysiology.....	6	Treatment of Stable COPD.....	33
2. Diagnosis and Assessment	9	Moving from Clinical Trials to Recommendations for Routine Practice Considerations.....	33
Key Points.....	10	Non-Pharmacologic Treatment.....	34
Diagnosis.....	10	<i>Smoking Cessation</i>	34
Symptoms.....	11	<i>Physical Activity</i>	34
Medical History.....	12	<i>Rehabilitation</i>	34
Physical Examination.....	12	<i>Vaccination</i>	34
Spirometry.....	12	Pharmacologic Treatment.....	35
Assessment Of Disease.....	12	<i>Bronchodilators - Recommendations</i>	35
Assessment of Symptoms.....	13	<i>Corticosteroids and Phosphodiesterase-4 Inhibitors - Recommendations</i>	37
Choice of Cut Points.....	13	Monitoring And Follow-Up.....	37
Spirometric Assessment.....	14	<i>Monitor Disease Progression and Development of Complications</i>	37
Assessment of Exacerbation Risk.....	14	<i>Monitor Pharmacotherapy and Other Medical Treatment</i>	37
Assessment of Comorbidities.....	15		
Combined COPD Assessment.....	15		
Additional Investigations.....	16		
Differential Diagnosis.....	17		

<i>Monitor Exacerbation History</i>	37
<i>Monitor Comorbidities</i>	37
<i>Surgery in the COPD Patient</i>	38
5. Management of Exacerbations	39
Key Points.....	40
Definition.....	40
Diagnosis.....	40
Assessment.....	41
Treatment Options.....	41
Treatment Setting.....	41
Pharmacologic Treatment.....	41
Respiratory Support.....	43
Hospital Discharge and Follow-up.....	45
Home Management of Exacerbations.....	45
Prevention of COPD Exacerbations.....	45
6. COPD and Comorbidities	47
Key Points.....	48
Introduction.....	48
Cardiovascular Disease.....	48
Osteoporosis.....	49
Anxiety and Depression.....	50
Lung Cancer.....	50
Infections.....	50
Metabolic Syndrome and Diabetes.....	50
Bronchiectasis.....	50
Impaired Cognitive Function.....	51
References	52
APPENDIX	
Diagnosis of Diseases of Chronic Airflow Limitation: Asthma, COPD and Asthma-COPD Overlap Syndrome (ACOS).....	A1
Figures	
Figure 1.1. Mechanisms Underlying Airflow Limitation in COPD.....	2
Figure 2.1A. Spirometry - Normal Trace.....	13
Figure 2.1B. Spirometry - Obstructive Disease.....	13
Figure 2.2. Relationship Between Health-Related Quality of Life, Post-Bronchodilator FEV ₁ and GOLD Spirometric Classification.....	14
Figure 2.3. Assessment Using Symptoms, Breathlessness, Spirometric Classification, and Risk of Exacerbations.....	15

Tables	
Table. Summary Observations.....	iv
Table. Description of Levels of Evidence.....	xvi
Table 2.1. Key Indicators for Considering a Diagnosis of COPD.....	10
Table 2.2. Causes of Chronic Cough.....	11
Table 2.3. Considerations in Performing Spirometry.....	12
Table 2.4. Modified Medical Research Council Questionnaire for Assessing the Severity of Breathlessness.....	13
Table 2.5. Classification of Severity of Airflow Limitation in COPD (Based on Post-Bronchodilator FEV ₁).....	14
Table 2.6. RISK IN COPD: Placebo-limb data from TORCH, Uplift, and Eclipse.....	15
Table 2.7. COPD and its Differential Diagnoses.....	18
Table 3.1. Treating Tobacco Use and Dependence: A Clinical Practice Guideline—Major Findings and Recommendations.....	20
Table 3.2. Brief Strategies to Help the Patient Willing to Quit.....	21
Table 3.3. Formulations and Typical Doses of COPD Medications.....	22
Table 3.4. Bronchodilators in Stable COPD.....	23
Table 3.5. Benefits of Pulmonary Rehabilitation in COPD.....	27
Table 4.1. Goals for Treatment of Stable COPD.....	32
Table 4.2. Model of Symptom/Risk of Evaluation of COPD.....	33
Table 4.3. Non-pharmacologic Management of COPD.....	34
Table 4.4. Initial Pharmacologic Management of COPD.....	36
Table 5.1. Assessment of COPD Exacerbations: Medical History.....	41
Table 5.2. Assessment of COPD Exacerbations: Signs of Severity.....	41
Table 5.3. Potential Indications for Hospital Assessment or Admission.....	41
Table 5.4. Management of Severe but Not Life-Threatening Exacerbations.....	42
Table 5.5. Therapeutic Components of Hospital Management.....	42
Table 5.6. Indications for ICU Admission.....	43
Table 5.7. Indications for Noninvasive Mechanical Ventilation.....	44

Table 5.8. Indications for Invasive Mechanical Ventilation.....	44
Table 5.9. Discharge Criteria.....	44
Table 5.10. Checklist of items to assess at time of Discharge from Hospital.....	44
Table 5.11. Items to Assess at Follow-Up Visit 4-6 Weeks After Discharge from Hospital.....	44

ACOS Tables

Table 1. Current definitions of asthma and COPD, and clinical description of ACOS.....	A2
Table 2a. Usual features of asthma, COPD and ACOS.....	A4
Table 2b. Features that favor asthma or COPD.....	A4
Table 3. Spirometric measures in asthma, COPD and ACOS.....	A6
Table 4. Summary of syndromic approach to diseases of chronic airflow limitation.....	A7
Table 5. Specialized investigations sometimes used in distinguishing asthma and COPD.....	A8

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

METHODOLOGY AND SUMMARY OF NEW RECOMMENDATIONS GLOBAL STRATEGY FOR DIAGNOSIS, MANAGEMENT AND PREVENTION OF COPD 2016 UPDATE¹

When the Global Initiative for Chronic Obstructive Lung Disease (GOLD) program was initiated in 1998, a goal was to produce recommendations for management of COPD based on the best scientific information available. The first report, *Global Strategy for Diagnosis, Management and Prevention of COPD* was issued in 2001. In 2006 and again in 2011 a complete revision was prepared based on published research. These reports, and their companion documents, have been widely distributed and translated into many languages and can be found on the GOLD website (www.goldcopd.org).

The GOLD Science Committee was established in 2002 to review published research on COPD management and prevention, to evaluate the impact of this research on recommendations in the GOLD documents related to management and prevention, and to post yearly updates on the GOLD website. Its members are recognized leaders in COPD research and clinical practice with the scientific credentials to contribute to the task of the Committee and are invited to serve in a voluntary capacity.

Updates of the 2011-revised report were released in January 2013, 2014, and 2015. This fourth update, released January 2016, is based on the impact of publications from July 1, 2014 through June 30, 2015. Posted on the website along with the updated documents is a list of all the publications reviewed by the Committee.

Process: To produce the updated documents a Pub Med search is completed using search fields established by the Committee: 1) *COPD, All Fields, Adult: 19+ years, only items with abstracts, Clinical Trial, Meta-analyses, Human*. The first search included publications for July 1, 2014 – February 28, 2015

¹The Global Strategy for Diagnosis, Management and Prevention of COPD (updated 2016), the Pocket Guide (updated 2016) and the complete list of references examined by the Committee are available on the GOLD website www.goldcopd.org.

²Members (2014-2015): C. Volgelmeier, Chair; A. Agustí, A. Anzueto, L. Fabbri, F. Martinez, N. Roche, R. Rodriguez-Roisin, D. Sin, D. Singh, R. Stockley, J. Vestbo, W. Wedzicha.

(publications already reviewed for the 2015 update were excluded) for review by the Committee during the meeting in May 2015. The second search included publications for January 1 – May 31, 2015 for review by the Committee during the meeting in September 2015. Publications in peer review journals not captured by Pub Med can be submitted to the Chair, GOLD Science Committee, providing an abstract and the full paper are submitted in (or translated into) English.

Members of the Committee receive a summary of citations and all abstracts. Each abstract is assigned to two Committee members, although all members are offered the opportunity to provide an opinion on any abstract. Members evaluate the abstract or, up to her/his judgment, the full publication, by answering four specific written questions from a short questionnaire, and to indicate if the scientific data presented impacts on recommendations in the GOLD report. If so, the member is asked to specifically identify modifications that should be made.

The GOLD Science Committee meets twice yearly to discuss each publication that was considered by at least 1 member of the Committee to potentially have an impact on the COPD management. The full Committee then reaches a consensus on whether to include it in the report, either as a reference supporting current recommendations, or to change the report. In the absence of consensus, disagreements are decided by an open vote of the full Committee. At its annual meeting in December, the final review and approval of all recommendations is provided by the GOLD Board of Directors.

Recommendations by the GOLD Committees for use of any medication are based on the best evidence available from the published literature and not on labeling directives from government regulators. The Committee does not make recommendations for therapies that have not been approved by at least one regulatory agency.

As an example of the workload of the Committee, for the 2016 update, between July 1, 2014 and May 31, 2015, 173 articles met the search criteria. Of the 173 papers, 15 were identified to have an impact on the GOLD report posted on the website in January 2016 either by: A) modifying, that is, changing the text or introducing a concept requiring a new recommendation to the report; B) confirming, that is, adding or replacing an existing reference; or C) requiring modification for clarification of the text.

SUMMARY OF RECOMMENDATIONS IN THE 2016 UPDATE

A. Additions to the text

Page 5, right column, paragraph 4, insert statement line 5 and add reference:

...and cough and sputum production is associated with increased mortality in mild to moderate COPD⁶¹⁶.

Reference 616: Putcha N, Drummond MB, Connett JE, Scanlon PD, Tashkin DP, Hansel NN, Wise RA. Chronic productive cough is associated with death in smokers with early COPD. *COPD*. 2014 Aug;11(4):451-8.

Page 13, right column, end of first paragraph insert statement and reference: A minimally clinical important differences (MCID) during rehabilitation of -0.4 for the CCQ has been identified⁶¹⁷.

Reference 617: Kon SS, Dilaver D, Mittal M, Nolan CM, Clark AL, Canavan JL, et al. The Clinical COPD Questionnaire: response to pulmonary rehabilitation and minimal clinically important difference. *Thorax*. 2014 Sep;69(9):793-8.

Page 25, left column, paragraph 1, insert statement and reference: Withdrawal of inhaled corticosteroids, in COPD patients at low risk of exacerbation, can be safe provided that patients are left on maintenance treatment with long-acting bronchodilators⁶¹⁸.

Reference 618: Rossi A, Guerriero M, Corrado A; OPTIMO/AIPO Study Group. Withdrawal of inhaled corticosteroids can be safe in COPD patients at low risk of exacerbation: a real-life study on the appropriateness of treatment in moderate COPD patients (OPTIMO). *Respir Res*. 2014 Jul 8;15:77.

Page 25, left column, paragraph 3, insert statement and reference: An inhaled corticosteroid/long-acting beta₂-agonist combination given once daily does not show relevant differences regarding efficacy compared to twice daily⁶¹⁹.

Reference 619: Dransfield MT, Feldman G, Korenblat P, LaForce CF, Locantore N, Pistolesi M, et al. Efficacy and safety of once-daily fluticasone furoate/vilanterol (100/25 mcg) versus twice-daily fluticasone propionate/salmeterol (250/50 mcg) in COPD patients. *Respir Med*. 2014 Aug;108(8):1171-9.

Page 25, left column, paragraph 4, insert statement and reference: However, systemic corticosteroids for treating acute exacerbations have been shown to improve symptoms, lung function, reduce rate of treatment failure, and shorten length of hospital stay. The effect of preventing a subsequent exacerbation has been shown in a pooled data analysis and it was demonstrated that systemic corticosteroids when being used to treat acute exacerbations can reduce 30-day readmission rates due to recurrent exacerbations.

Reference 620: Criner GJ, Bourbeau J, Diekemper RL, Ouellette DR, Goodridge D, Hernandez P, et al. Prevention of acute exacerbations of COPD: American College of Chest Physicians and Canadian Thoracic Society Guideline. *Chest*. 2015 Apr;147(4):894-2.

Page 28, left column, paragraph 3 insert statement and reference: Depressed mood is a risk factor for non-completion of rehabilitation programs⁶²¹.

Reference 622: Busch AM, Scott-Sheldon LA, Pierce J, Chattillion EA, Cunningham K, Buckley ML, et al. Depressed mood predicts pulmonary rehabilitation completion among women, but not men. *Respir Med*. 2014 Jul;108(7):1007-13.

B. References that provided confirmation or update of previous recommendations

Page 5, left column, paragraph 2, insert reference: **Reference 614:** She L Yang P, Wang Y, Qin X, Fan L Wang Y, Gao G, Luo G, et al. Chinese water-pipe smoking and the risk of capo. *Chest*. 2014 Oct;146(4):924-31.

Page 5, left column, paragraph 4, insert reference: **Reference 615:** Ramirez-Venegas A, Sansores RH, Quintana-Carrillo RH, Velazquez-Uncal M, Hernandez-

Zenteno RJ, Sanchez-Romero C, et al. FEV1 decline in patients with chronic obstructive pulmonary disease associated with biomass exposure. *Am J Respir Crit Care Med*. 2014 Nov 1;190(9):996-1002.

Page 40, right column last paragraph insert: However, clinically useful biomarkers for COPD patients in stable condition have yet to be identified.

Page 25, right column, second paragraph, insert reference:

Reference 621: Yu T, Fain K, Boyd CM, Singh S, Weiss CO, Li T, et al. Benefits and harms of roflumilast in moderate to severe COPD. *Thorax*. 2014 Jul;69(7):616-22.

Page 40, right column, paragraph 3, insert reference:

Reference 623: Konecny T, Park JY, Somers KR, Konecny D, Orban M, Soucek F, et al. Relation of chronic obstructive pulmonary disease to atrial and ventricular arrhythmias. *Am J Cardiol*. 2014 Jul 15;114(2):272-7.

Page 42, right column, paragraph 3, insert reference:

Reference 624: Walters JA, Tan DJ, White CJ, Wood-Baker R. Different durations of corticosteroid therapy for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2014 Dec 10;12:CD006897.

Page 45, right column, paragraph 2, insert reference:

Reference 625: Ingebrigtsen TS, Marott JL, Nordestgaard BG, Lange P, Hallas J, Vestbo J. Statin use and exacerbations in individuals with chronic obstructive pulmonary disease. *Thorax*. 2015 Jan;70(1):33-40.

C. Inserts related to tables/figures and special topics covered by the Committee

Page 20, key points, insert: Non-surgical bronchoscopic lung volume reduction techniques should not be used outside clinical trials until more data are available.

Page 22, Table 3-3, insert under Long-acting beta2-agonists: Olodaterol 5 mcg (SMI), 24 hours

Page 22, Table 3-3, insert under Combination long-acting beta2-agonist plus anticholinergic in one inhaler: Olodaterol/tiotropium 5mcg/5mcg (SMI), 24 hours

GLOBAL STRATEGY FOR THE DIAGNOSIS, MANAGEMENT, AND PREVENTION OF COPD

INTRODUCTION

Much has changed in the 10 years since the first GOLD report, *Global Strategy for the Diagnosis, Management, and Prevention of COPD*, was published. This major revision builds on the strengths from the original recommendations and incorporates new knowledge.

One of the strengths was the treatment objectives. These have stood the test of time, but are now organized into two groups: objectives that are directed towards immediately relieving and reducing the impact of symptoms, and objectives that reduce the risk of adverse health events that may affect the patient at some point in the future. (Exacerbations are an example of such events.) This emphasizes the need for clinicians to maintain a focus on both the short-term and long-term impact of COPD on their patients.

A second strength of the original strategy was the simple, intuitive system for classifying COPD severity. This was based upon the FEV₁ and was called a staging system because it was believed, at the time, that the majority of patients followed a path of disease progression in which the severity of the disease tracked the severity of the airflow limitation. Much is now known about the characteristics of patients in the different GOLD stages – for example, their level of risk of exacerbations, hospitalization, and death. However at an individual patient level, the FEV₁ is an unreliable marker of the severity of breathlessness, exercise limitation, and health status impairment. This report retains the GOLD classification system because it is a predictor of future adverse events, but the term “Stage” is now replaced by “Grade.”

At the time of the original report, improvement in both symptoms and health status was a GOLD treatment objective, but symptoms assessment did not have a direct relation to the choice of management, and health status measurement was a complex process largely confined to clinical studies. Now, there are simple and reliable questionnaires designed for use in routine daily clinical practice. These are available in many languages. These developments have enabled a new assessment system to be developed that draws together a measure of the impact of the patient’s symptoms and an assessment of the patient’s risk of having a serious adverse health event in the future. In turn, this new assessment system has led

to the construction of a new approach to management— one that matches assessment to treatment objectives. The new management approach can be used in any clinical setting anywhere in the world and moves COPD treatment towards individualized medicine – matching the patient’s therapy more closely to his or her needs.

BACKGROUND

Chronic Obstructive Pulmonary Disease (COPD), the fourth leading cause of death in the world¹, represents an important public health challenge that is both preventable and treatable. COPD is a major cause of chronic morbidity and mortality throughout the world; many people suffer from this disease for years, and die prematurely from it or its complications. Globally, the COPD burden is projected to increase in coming decades because of continued exposure to COPD risk factors and aging of the population².

In 1998, with the cooperation of the National Heart, Lung, and Blood Institute, NIH and the World Health Organization, the Global Initiative for Chronic Obstructive Lung Disease (GOLD) was implemented. Its goals were to increase awareness of the burden of COPD and to improve prevention and management of COPD through a concerted worldwide effort of people involved in all facets of health care and health care policy. An important and related goal was to encourage greater research interest in this highly prevalent disease.

In 2001, GOLD released its first report, *Global Strategy for the Diagnosis, Management, and Prevention of COPD*. This report was not intended to be a comprehensive textbook on COPD, but rather to summarize the current state of the field. It was developed by individuals with expertise in COPD research and patient care and was based on the best-validated concepts of COPD pathogenesis at that time, along with available evidence on the most appropriate management and prevention strategies. It provided state-of-the-art information about COPD for pulmonary specialists and other interested physicians and served as a source document for the production of various communications for other audiences, including an Executive Summary³, a Pocket Guide for Health Care Professionals, and a Patient Guide.

Immediately following the release of the first GOLD report in 2001, the GOLD Board of Directors appointed a Science Committee, charged with keeping the GOLD documents up-to-date by reviewing published research, evaluating the impact of this research on the management

recommendations in the GOLD documents, and posting yearly updates of these documents on the GOLD Website. The first update to the GOLD report was posted in July 2003, based on publications from January 2001 through December 2002. A second update appeared in July 2004, and a third in July 2005, each including the impact of publications from January through December of the previous year. In January 2005, the GOLD Science Committee initiated its work to prepare a comprehensively updated version of the GOLD report; it was released in 2006. The methodology used to create the annual updated documents, and the 2006 revision, appears at the front of each volume.

During the period from 2006 to 2010, again annual updated documents were prepared and released on the GOLD Website, along with the methodology used to prepare the documents and the list of published literature reviewed to examine the impact on recommendations made in the annual updates. In 2009, the GOLD Science Committee recognized that considerable new information was available particularly related to diagnosis and approaches to management of COPD that warranted preparation of a significantly revised report. The work on this new revision was implemented in mid-2009 while at the same time the Committee prepared the 2010 update.

METHODOLOGY

In September 2009 and in May and September 2010 while preparing the annual updated reports (<http://www.goldcopd.org>), Science Committee members began to identify the literature that impacted on major recommendations, especially for COPD diagnosis and management. Committee members were assigned chapters to review for proposed modifications and soon reached consensus that the report required significant change to reach the target audiences – the general practitioner and the individuals in clinics around the world who first see patients who present with respiratory symptoms that could lead to a diagnosis of COPD. In the summer of 2010 a writing committee was established to produce an outline of proposed chapters, which was first presented in a symposium for the European Respiratory Society in Barcelona, 2010. The writing committee considered recommendations from this session throughout fall 2010 and spring 2011. During this period the GOLD Board of Directors and GOLD National Leaders were provided summaries of the major new directions recommended. During the summer of 2011 the document was circulated for review to GOLD National Leaders, and other COPD opinion leaders in a variety of countries. The names of the individuals who submitted reviews appear in the front of this report. In September 2011 the GOLD Science Committee reviewed the comments and made final recommendations. The report was launched during a symposium hosted by the Asian Pacific Society of Respirology in November 2011.

NEW ISSUES PRESENTED IN THIS REPORT

1. This document has been considerably shortened in length by limiting to Chapter 1 the background information on COPD. Readers who wish to access more comprehensive information about the pathophysiology of COPD are referred to a variety of excellent textbooks that have appeared in the last decade.
2. Chapter 2 includes information on diagnosis and assessment of COPD. The definition of COPD has not been significantly modified but has been reworded for clarity.
3. Assessment of COPD is based on the patient's level of symptoms, future risk of exacerbations, the severity of the spirometric abnormality, and the identification of comorbidities. Whereas spirometry was previously used to support a diagnosis of COPD, spirometry is now required to make a confident diagnosis of COPD.
4. The spirometric classification of airflow limitation is divided into four Grades (GOLD 1, Mild; GOLD 2, Moderate; GOLD 3, Severe; and GOLD 4, Very Severe) using the fixed ratio, postbronchodilator $FEV_1/FVC < 0.70$, to define airflow limitation. It is recognized that use of the fixed ratio (FEV_1/FVC) may lead to more frequent diagnoses of COPD in older adults with mild COPD as the normal process of aging affects lung volumes and flows, and may lead to under-diagnosis in adults younger than 45 years. The concept of staging has been abandoned as a staging system based on FEV_1 alone was inadequate and the evidence for an alternative staging system does not exist. The most severe spirometric Grade, GOLD 4, does not include reference to respiratory failure as this seemed to be an arbitrary inclusion.
5. A new chapter (Chapter 3) on therapeutic approaches has been added. This includes descriptive information on both pharmacologic and non-pharmacologic therapies, identifying adverse effects, if any.
6. Management of COPD is presented in three chapters: Management of Stable COPD (Chapter 4); Management of COPD Exacerbations (Chapter 5); and COPD and Comorbidities (Chapter 6), covering both management of comorbidities in patients with COPD and of COPD in patients with comorbidities.
7. In Chapter 4, Management of Stable COPD, recommended approaches to both pharmacologic and non-pharmacologic treatment of COPD are presented. The chapter begins with the importance of identification and reduction of risk factors. Cigarette smoke continues to be

identified as the most commonly encountered risk factor for COPD and elimination of this risk factor is an important step toward prevention and control of COPD. However, more data are emerging to recognize the importance of other risk factors for COPD that should be taken into account where possible. These include occupational dusts and chemicals, and indoor air pollution from biomass cooking and heating in poorly ventilated dwellings – the latter especially among women in developing countries.

8. In previous GOLD documents, recommendations for management of COPD were based solely on spirometric category. However, there is considerable evidence that the level of FEV₁ is a poor descriptor of disease status and for this reason the management of stable COPD based on a strategy considering both disease impact (determined mainly by symptom burden and activity limitation) and future risk of disease progression (especially of exacerbations) is recommended.

9. Chapter 5, Management of Exacerbations, presents a revised definition of a COPD exacerbation.

10. Chapter 6, Comorbidities and COPD, focuses on cardiovascular diseases, osteoporosis, anxiety and depression, lung cancer, infections, and metabolic syndrome and diabetes.

11. APPENDIX: The report Diagnosis of Diseases of Chronic Airflow Limitation: Asthma, COPD, and Asthma-COPD Overlap Syndrome (ACOS) has been added.

LEVELS OF EVIDENCE

Levels of evidence are assigned to management recommendations where appropriate. Evidence levels are indicated in boldface type enclosed in parentheses after the relevant statement e.g., (**Evidence A**). The methodological issues concerning the use of evidence from meta-analyses were carefully considered. This evidence level scheme (**Table A**) has been used in previous GOLD reports, and was in use throughout the preparation of this document⁴.

Table A Description of Levels of Evidence

Evidence Category	Sources of Evidence	Definition
A	Randomized controlled trials (RCTs). Robust body of data.	Evidence is from endpoints of well-designed RCTs that provide a consistent pattern of findings in the population for which the recommendation is made. Category A requires substantial numbers of studies involving substantial numbers of participants.
B	Randomized controlled trials (RCTs). Limited body of data.	Evidence is from endpoints of intervention studies that include only a limited number of patients, posthoc or subgroup analysis of RCTs, or meta-analysis of RCTs. In general, Category B pertains when few randomized trials exist, they are small in size, they were undertaken in a population that differs from the target population of the recommendation, or the results are somewhat inconsistent.
C	Nonrandomized trials. Observational studies.	Evidence is from outcomes of uncontrolled or nonrandomized trials or from observational studies
D	Panel Consensus Judgment.	This category is used only in cases where the provision of some guidance was deemed valuable but the clinical literature addressing the subject was deemed insufficient to justify placement in one of the other categories. The Panel Consensus is based on clinical experience or knowledge that does not meet the above-listed criteria

CHAPTER

1

DEFINITION AND OVERVIEW


COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

CHAPTER 1: DEFINITION AND OVERVIEW

KEY POINTS:

- *Chronic Obstructive Pulmonary Disease (COPD), a common preventable and treatable disease, is characterized by persistent airflow limitation that is usually progressive and associated with an enhanced chronic inflammatory response in the airways and the lung to noxious particles or gases. Exacerbations and comorbidities contribute to the overall severity in individual patients.*
- COPD is a leading cause of morbidity and mortality worldwide and results in an economic and social burden that is both substantial and increasing.
- Inhaled cigarette smoke and other noxious particles such as smoke from biomass fuels cause lung inflammation, a normal response that appears to be modified in patients who develop COPD. This chronic inflammatory response may induce parenchymal tissue destruction (resulting in emphysema), and disrupt normal repair and defense mechanisms (resulting in small airway fibrosis). These pathological changes lead to air trapping and progressive airflow limitation, and in turn to breathlessness and other characteristic symptoms of COPD.

Many previous definitions of COPD have emphasized the terms “emphysema” and “chronic bronchitis,” which are not included in the definition used in this or earlier GOLD reports. Emphysema, or destruction of the gas-exchanging surfaces of the lung (alveoli), is a pathological term that is often (but incorrectly) used clinically and describes



only one of several structural abnormalities present in patients with COPD. Chronic bronchitis, or the presence of cough and sputum production for at least 3 months in each of two consecutive years, remains a clinically and epidemiologically useful term. However, it is important to recognize that chronic cough and sputum production (chronic bronchitis) is an independent disease entity that may precede or follow the development of airflow limitation and may be associated with development and/or acceleration of fixed airflow limitation. Chronic bronchitis also exists in patients with normal spirometry.

DEFINITION

Chronic Obstructive Pulmonary Disease (COPD), a common preventable and treatable disease, is characterized by persistent airflow limitation that is usually progressive and associated with an enhanced chronic inflammatory response in the airways and the lung to noxious particles or gases. Exacerbations and comorbidities contribute to the overall severity in individual patients.

The chronic airflow limitation characteristic of COPD is caused by a mixture of small airways disease (obstructive bronchiolitis) and parenchymal destruction (emphysema), the relative contributions of which vary from person to person (**Figure 1.1**). Chronic inflammation causes structural changes and narrowing of the small airways. Destruction of the lung parenchyma, also by inflammatory processes, leads to the loss of alveolar attachments to the small airways and decreases lung elastic recoil; in turn, these changes diminish the ability of the airways to remain open during expiration. Airflow limitation is best measured by spirometry, as this is the most widely available, reproducible test of lung function.

BURDEN OF COPD

COPD is a leading cause of morbidity and mortality worldwide and results in an economic and social burden that is both substantial and increasing^{2,5}. COPD prevalence, morbidity, and mortality vary across countries and across different groups within countries. COPD is the result of cumulative exposures over decades. Often, the prevalence of COPD is directly related to the prevalence of tobacco smoking, although in many countries, outdoor, occupational and indoor air pollution – the latter resulting from the burning of wood and other biomass fuels – are major COPD risk factors⁶. The prevalence and burden of COPD are projected to increase in the coming decades due to continued exposure to COPD risk factors and the changing age structure of the world’s population (with more people living longer and therefore expressing the long-term effects of exposure to COPD risk factors)⁵. Information on the burden of COPD can be found on international

Websites such as those of the **World Health Organization (WHO)** (<http://www.who.int>) and the **World Bank/WHO Global Burden of Disease Study** (http://www.who.int/topics/global_burden_of_disease). Aging itself is a risk factor for COPD and aging of the airways and parenchyma mimic some of the structural changes associated with COPD⁷.

Prevalence

Existing COPD prevalence data show remarkable variation due to differences in survey methods, diagnostic criteria, and analytic approaches⁸. The lowest estimates of prevalence are those based on self-reporting of a doctor diagnosis of COPD or equivalent condition. For example, most national data show that less than 6% of the adult population has been told that they have COPD⁸. This likely reflects the widespread under-recognition and under-diagnosis of COPD⁹.

Despite the complexities, data are emerging that enable some conclusions to be drawn regarding COPD prevalence, not least because of increasing data quality control. A systematic review and meta-analysis of studies carried out in 28 countries between 1990 and 2004⁸, and an additional study from Japan¹⁰, provide evidence that the prevalence of COPD is appreciably higher in smokers and ex-smokers than in nonsmokers, in those over 40 years of age than those under 40, and in men than in women. The Latin American Project for the Investigation of Obstructive Lung Disease (PLATINO)¹¹ examined the prevalence of post-bronchodilator airflow limitation among persons over age 40 in five major Latin American cities, each in a different country – Brazil, Chile, Mexico, Uruguay, and Venezuela. In each country, the prevalence of COPD increased steeply with age, with the highest prevalence among those over age 60, ranging in the total population from a low of 7.8% in Mexico City, Mexico to a high of 19.7% in Montevideo, Uruguay. In all cities/countries the prevalence was appreciably higher in men than in women¹¹, which contrasts with findings from European cities such as Salzburg¹². The Burden of Obstructive Lung Diseases program (BOLD) has carried out surveys in several parts of the world¹³ and has documented more severe disease than previously found and a substantial prevalence (3-11%) of COPD among never-smokers.

Morbidity

Morbidity measures traditionally include physician visits, emergency department visits, and hospitalizations. Although COPD databases for these outcome parameters are less readily available and usually less reliable than mortality databases, the limited data available indicate that morbidity due to COPD increases with age¹⁰⁻¹². Morbidity from COPD may be affected by other comorbid chronic conditions (e.g., cardiovascular disease, musculoskeletal

impairment, diabetes mellitus) that are related to COPD and may have an impact on the patient's health status, as well as interfere with COPD management.

Mortality

The World Health Organization publishes mortality statistics for selected causes of death annually for all WHO regions; additional information is available from the **WHO Evidence for Health Policy Department** (<http://www.who.int/evidence>). Data must be interpreted cautiously, however, because of inconsistent use of terminology for COPD. In the 10th revision of the ICD, deaths from COPD or chronic airways obstruction are included in the broad category of "COPD and allied conditions" (ICD-10 codes J42-46).

Under-recognition and under-diagnosis of COPD still affect the accuracy of mortality data^{14,15}. Although COPD is often a primary cause of death, it is more likely to be listed as a contributory cause of death or omitted from the death certificate entirely^{16,504}. However, it is clear that COPD is one of the most important causes of death in most countries. The Global Burden of Disease Study projected that COPD, which ranked sixth as a cause of death in 1990, will become the third leading cause of death worldwide by 2020; a newer projection estimated COPD will be the fourth leading cause of death in 2030⁵. This increased mortality is mainly driven by the expanding epidemic of smoking, reduced mortality from other common causes of death (e.g. ischemic heart disease, infectious diseases), and aging of the world population.

Economic Burden

COPD is associated with significant economic burden. In the European Union, the total direct costs of respiratory disease are estimated to be about 6% of the total health care budget, with COPD accounting for 56% (38.6 billion Euros) of this cost of respiratory disease¹⁷. In the United States the estimated direct costs of COPD are \$29.5 billion and the indirect costs \$20.4 billion¹⁸. COPD exacerbations account for the greatest proportion of the total COPD burden on the health care system. Not surprisingly, there is a striking direct relationship between the severity of COPD and the cost of care, and the distribution of costs changes as the disease progresses. For example, hospitalization and ambulatory oxygen costs soar as COPD severity increases. Any estimate of direct medical expenditures for home care under-represents the true cost of home care to society, because it ignores the economic value of the care provided to those with COPD by family members.

In developing countries, direct medical costs may be less important than the impact of COPD on workplace and home productivity. Because the health care sector might not provide long-term supportive care services for severely

disabled individuals, COPD may force two individuals to leave the workplace—the affected individual and a family member who must now stay home to care for the disabled relative. Since human capital is often the most important national asset for developing countries, the indirect costs of COPD may represent a serious threat to their economies.

Social Burden

Since mortality offers a limited perspective on the human burden of a disease, it is desirable to find other measures of disease burden that are consistent and measurable across nations. The authors of the Global Burden of Disease Study designed a method to estimate the fraction of mortality and disability attributable to major diseases and injuries using a composite measure of the burden of each health problem, the Disability-Adjusted Life Year (DALY)^{2,19,20}. The DALYs for a specific condition are the sum of years lost because of premature mortality and years of life lived with disability, adjusted for the severity of disability. In 1990, COPD was the twelfth leading cause of DALYs lost in the world, responsible for 2.1% of the total. According to the projections, COPD will be the seventh leading cause of DALYs lost worldwide in 2030⁵.

FACTORS THAT INFLUENCE DISEASE DEVELOPMENT AND PROGRESSION

Although cigarette smoking is the best-studied COPD risk factor, it is not the only one and there is consistent evidence from epidemiologic studies that nonsmokers may also develop chronic airflow limitation^{21,24}. Much of the evidence concerning risk factors for COPD comes from cross-sectional epidemiological studies that identify associations rather than cause-and-effect relationships. Although several longitudinal studies of COPD have followed groups and populations for up to 20 years²⁵, none has monitored the progression of the disease through its entire course, or has included the pre- and perinatal periods which may be important in shaping an individual's future COPD risk. Thus, current understanding of risk factors for COPD is in many respects still incomplete.

COPD results from a gene-environment interaction. Among people with the same smoking history, not all will develop COPD due to differences in genetic predisposition to the disease, or in how long they live. Risk factors for COPD may also be related in more complex ways. For example, gender may influence whether a person takes up smoking or experiences certain occupational or environmental exposures; socioeconomic status may be linked to a child's birth weight (as it impacts on lung growth and development

and in turn on susceptibility to develop the disease); and longer life expectancy will allow greater lifetime exposure to risk factors. Understanding the relationships and interactions among risk factors requires further investigation.

Genes

The genetic risk factor that is best documented is a severe hereditary deficiency of alpha-1 antitrypsin²⁶, a major circulating inhibitor of serine proteases. Although alpha-1 antitrypsin deficiency is relevant to only a small part of the world's population, it illustrates the interaction between genes and environmental exposures leading to COPD.

A significant familial risk of airflow limitation has been observed in smoking siblings of patients with severe COPD²⁷, suggesting that genetic together with environmental factors could influence this susceptibility. Single genes such as the gene encoding matrix metalloproteinase 12 (*MMP12*) have been related to decline in lung function²⁸. Although several genome-wide association studies indicate a role of the gene for the alpha-nicotinic acetylcholine receptor as well as the hedge-hog interacting protein gene and possibly one or two others, there remains a discrepancy between findings from analyses of COPD and lung function as well as between genome-wide association study analyses and candidate gene analyses²⁹⁻³³.

Age and Gender

Age is often listed as a risk factor for COPD. It is unclear if healthy aging as such leads to COPD or if age reflects the sum of cumulative exposures throughout life. In the past, most studies showed that COPD prevalence and mortality were greater among men than women but data from developed countries^{18,34} show that the prevalence of the disease is now almost equal in men and women, probably reflecting the changing patterns of tobacco smoking. Some studies have even suggested that women are more susceptible to the effects of tobacco smoke than men³⁵⁻³⁸.

Lung Growth and Development

Lung growth is related to processes occurring during gestation, birth, and exposures during childhood and adolescence^{39,40}. Reduced maximal attained lung function (as measured by spirometry) may identify individuals who are at increased risk for the development of COPD⁴¹. Any factor that affects lung growth during gestation and childhood has the potential for increasing an individual's risk of developing COPD. For example, a large study and meta-analysis confirmed a positive association between birth weight and FEV₁ in adulthood⁴², and several studies have found an effect of early childhood lung infections.

4 DEFINITION AND OVERVIEW

A study found that factors in early life termed “childhood disadvantage factors” were as important as heavy smoking in predicting lung function in early adult life⁴³.

Exposure to Particles

Across the world, cigarette smoking is the most commonly encountered risk factor for COPD. Cigarette smokers have a higher prevalence of respiratory symptoms and lung function abnormalities, a greater annual rate of decline in FEV₁, and a greater COPD mortality rate than nonsmokers⁴⁴. Other types of tobacco (e.g., pipe, cigar, water pipe^{45,614}) and marijuana⁴⁶ are also risk factors for COPD^{47,48}. Passive exposure to cigarette smoke (also known as environmental tobacco smoke or ETS) may also contribute to respiratory symptoms⁴⁹ and COPD⁵⁰ by increasing the lung’s total burden of inhaled particles and gases^{51,52}. Smoking during pregnancy may also pose a risk for the fetus, by affecting lung growth and development in utero and possibly the priming of the immune system^{53,54}.

Occupational exposures, including organic and inorganic dusts and chemical agents and fumes, are an underappreciated risk factor for COPD⁵⁵⁻⁵⁷. An analysis of the large U.S. population-based NHANES III survey of almost 10,000 adults aged 30-75 years estimated the fraction of COPD attributable to work was 19.2% overall, and 31.1% among never-smokers⁵⁸. These estimates are consistent with a statement published by the American Thoracic Society that concluded that occupational exposures account for 10-20% of either symptoms or functional impairment consistent with COPD⁵⁹. The risk from occupational exposures in less regulated areas of the world is likely to be much higher than reported in studies from Europe and North America.

Wood, animal dung, crop residues, and coal, typically burned in open fires or poorly functioning stoves, may lead to very high levels of indoor air pollution. Evidence continues to grow that indoor pollution from biomass cooking and heating in poorly ventilated dwellings is an important risk factor for COPD^{60-66,615}. Almost 3 billion people worldwide use biomass and coal as their main source of energy for cooking, heating, and other household needs, so the population at risk worldwide is very large^{63,67}.

High levels of urban air pollution are harmful to individuals with existing heart or lung disease. The role of outdoor air pollution in causing COPD is unclear, but appears to be small when compared with that of cigarette smoking. It has also been difficult to assess the effects of single pollutants in long-term exposure to atmospheric pollution. However, air pollution from fossil fuel combustion, primarily from motor vehicle emissions in cities, is associated with decrements of respiratory function⁶⁸. The relative effects of short-term, high-peak exposures and long-term, low-level exposures are yet to be resolved.

Socioeconomic Status

Poverty is clearly a risk factor for COPD but the components of poverty that contribute to this are unclear. There is strong evidence that the risk of developing COPD is inversely related to socioeconomic status⁶⁹. It is not clear, however, whether this pattern reflects exposures to indoor and outdoor air pollutants, crowding, poor nutrition, infections, or other factors that are related to low socioeconomic status.

Asthma/Bronchial Hyperreactivity

Asthma may be a risk factor for the development of COPD, although the evidence is not conclusive. In a report from a longitudinal cohort of the Tucson Epidemiological Study of Airway Obstructive Disease, adults with asthma were found to have a twelve-fold higher risk of acquiring COPD over time than those without asthma, after adjusting for smoking⁷⁰. Another longitudinal study of people with asthma found that around 20% of subjects developed irreversible airflow limitation and reduced transfer coefficient⁷¹, and in a longitudinal study self-reported asthma was associated with excess loss of FEV₁ in the general population⁷². In the European Community Respiratory Health Survey, bronchial hyperresponsiveness was second only to cigarette smoking as the leading risk factor for COPD, responsible for 15% of the population attributable risk (smoking had a population attributable risk of 39%)⁷³. The pathology of chronic airflow limitation in asthmatic nonsmokers and non-asthmatic smokers is markedly different, suggesting that the two disease entities may remain different even when presenting with similarly reduced lung function⁷⁴. However, clinically separating asthma from COPD may not be easy.

Bronchial hyperreactivity can exist without a clinical diagnosis of asthma and has been shown to be an independent predictor of COPD in population studies⁷⁵ as well as an indicator of risk of excess decline in lung function in patients with mild COPD⁷⁶.

Chronic Bronchitis

In the seminal study by Fletcher and coworkers, chronic bronchitis was not associated with decline in lung function⁷⁷. However, subsequent studies have found an association between mucus hypersecretion and FEV₁ decline⁷⁸, and cough and sputum production is associated with increased mortality in mild to moderate COPD⁶¹⁶. In younger adults who smoke the presence of chronic bronchitis is associated with an increased likelihood of developing COPD^{79,80}.

Infections

A history of severe childhood respiratory infection has been associated with reduced lung function and increased respiratory symptoms in adulthood^{39,73}. Susceptibility to infections plays a role in exacerbations of COPD but the effect on the development of the disease is less clear.

PATHOLOGY, PATHOGENESIS AND PATHOPHYSIOLOGY

HIV infection has been shown to accelerate the onset of smoking-related emphysema⁸¹. Tuberculosis has been found to be a risk factor for COPD^{82,83}. In addition, tuberculosis is both a differential diagnosis to COPD and a potential comorbidity^{83,84}.

Inhaled cigarette smoke and other noxious particles such as smoke from biomass fuels cause lung inflammation, a normal response that appears to be modified in patients who develop COPD. This chronic inflammatory response may induce parenchymal tissue destruction (resulting in emphysema), and disrupt normal repair and defense mechanisms (resulting in small airway fibrosis). These pathological changes lead to air trapping and progressive airflow limitation. A brief overview follows of the pathologic changes in COPD, their cellular and molecular mechanisms, and how these underlie physiologic abnormalities and symptoms characteristic of the disease⁸⁵.

Pathology

Pathological changes characteristic of COPD are found in the airways, lung parenchyma, and pulmonary vasculature⁸⁶. The pathological changes include chronic inflammation, with increased numbers of specific inflammatory cell types in different parts of the lung, and structural changes resulting from repeated injury and repair. In general, the inflammatory and structural changes in the airways increase with disease severity and persist on smoking cessation.

Pathogenesis

The inflammation in the respiratory tract of COPD patients appears to be a modification of the inflammatory response of the respiratory tract to chronic irritants such as cigarette smoke. The mechanisms for this amplified inflammation are not yet understood but may be genetically determined. Patients can clearly develop COPD without smoking, but the nature of the inflammatory response in these patients is unknown. Oxidative stress and an excess of proteinases in the lung further modify lung inflammation. Together, these mechanisms lead to the characteristic pathological changes in COPD. Lung inflammation persists after smoking cessation through unknown mechanisms, although autoantigens and persistent microorganisms may play a role⁸⁷.

Oxidative Stress. Oxidative stress may be an important amplifying mechanism in COPD⁸⁸. Biomarkers of oxidative stress (e.g., hydrogen peroxide, 8-isoprostane) are increased in the exhaled breath condensate, sputum, and systemic circulation of COPD patients. Oxidative stress is further increased in exacerbations. Oxidants are generated by cigarette smoke and other inhaled particulates, and released from activated inflammatory cells such as macrophages and neutrophils. There may also be a reduction in endogenous antioxidants in COPD patients as a result of reduction in a transcription factor called Nrf2 that regulates many antioxidant genes⁸⁹.

Protease-Antiprotease Imbalance. There is compelling evidence for an imbalance in the lungs of COPD patients between proteases that break down connective tissue components and antiproteases that protect against this. Several proteases, derived from inflammatory cells and epithelial cells, are increased in COPD patients. There is increasing evidence that they may interact with each other. Protease-mediated destruction of elastin, a major connective tissue component in lung parenchyma, is believed to be an important feature of emphysema and is likely to be irreversible.

Inflammatory Cells. COPD is characterized by a specific pattern of inflammation involving increased numbers of CD8+ (cytotoxic) Tc1 lymphocytes present only in smokers that develop the disease⁸⁵. These cells, together with neutrophils and macrophages, release inflammatory mediators and enzymes and interact with structural cells in the airways, lung parenchyma and pulmonary vasculature⁹⁰.

Inflammatory Mediators. The wide variety of inflammatory mediators that have been shown to be increased in COPD patients⁹¹ attract inflammatory cells from the circulation (chemotactic factors), amplify the inflammatory process (proinflammatory cytokines), and induce structural changes (growth factors)⁹².

Differences in Inflammation Between COPD and Asthma.

Although both COPD and asthma are associated with chronic inflammation of the respiratory tract, there are differences in the inflammatory cells and mediators involved in the two diseases, which in turn account for differences in physiological effects, symptoms, and response to therapy⁷⁴. Some patients with COPD have features consistent with asthma and may have a mixed inflammatory pattern with increased eosinophils.

Pathophysiology

There is now a good understanding of how the underlying disease process in COPD leads to the characteristic

6 DEFINITION AND OVERVIEW

physiologic abnormalities and symptoms. For example, inflammation and narrowing of peripheral airways leads to decreased FEV₁. Parenchymal destruction due to emphysema also contributes to airflow limitation and leads to decreased gas transfer.

Airflow Limitation and Air Trapping. The extent of inflammation, fibrosis, and luminal exudates in small airways is correlated with the reduction in FEV₁ and FEV₁/FVC ratio, and probably with the accelerated decline in FEV₁ characteristic of COPD⁹⁰. This peripheral airway obstruction progressively traps air during expiration, resulting in hyperinflation. Although emphysema is more associated with gas exchange abnormalities than with reduced FEV₁, it does contribute to gas trapping during expiration. This is especially so as alveolar attachments to small airways are destroyed when the disease becomes more severe. Hyperinflation reduces inspiratory capacity such that functional residual capacity increases, particularly during exercise (dynamic hyperinflation), resulting in increased dyspnea and limitation of exercise capacity. These factors contribute to impairment of the intrinsic contractile properties of respiratory muscles; this results in upregulation of local pro-inflammatory cytokines. It is thought that hyperinflation develops early in the disease and is the main mechanism for exertional dyspnea^{93,94}. Bronchodilators acting on peripheral airways reduce air trapping, thereby reducing lung volumes and improving symptoms and exercise capacity⁹³.

Gas Exchange Abnormalities. Gas exchange abnormalities result in hypoxemia and hypercapnia, and have several mechanisms in COPD. In general, gas transfer for oxygen and carbon dioxide worsens as the disease progresses. Reduced ventilation may also be due to reduced ventilatory drive. This may lead to carbon dioxide retention when it is combined with reduced ventilation due to a high work of breathing because of severe obstruction and hyperinflation coupled with ventilatory muscle impairment. The abnormalities in alveolar ventilation and a reduced pulmonary vascular bed further worsen the V_A/Q abnormalities⁹⁵.

Mucus Hypersecretion. Mucus hypersecretion, resulting in a chronic productive cough, is a feature of chronic bronchitis and is not necessarily associated with airflow limitation. Conversely, not all patients with COPD have symptomatic mucus hypersecretion. When present, it is due to an increased number of goblet cells and enlarged submucosal glands in response to chronic airway irritation by cigarette smoke and other noxious agents. Several mediators and proteases stimulate mucus hypersecretion and many of them exert their effects through the activation of epidermal growth factor receptor (EGFR)⁹⁶.

Pulmonary Hypertension. Pulmonary hypertension may develop late in the course of COPD and is due mainly to hypoxic vasoconstriction of small pulmonary arteries, eventually resulting in structural changes that include intimal hyperplasia and later smooth muscle hypertrophy/hyperplasia⁹⁷. There is an inflammatory response in vessels similar to that seen in the airways and evidence of endothelial cell dysfunction. The loss of the pulmonary capillary bed in emphysema may also contribute to increased pressure in the pulmonary circulation. Progressive pulmonary hypertension may lead to right ventricular hypertrophy and eventually to right-side cardiac failure.

Exacerbations. Exacerbations of respiratory symptoms often occur in patients with COPD, triggered by infection with bacteria or viruses (which may coexist), environmental pollutants, or unknown factors. Patients with bacterial and viral episodes have a characteristic response with increased inflammation. During respiratory exacerbations there is increased hyperinflation and gas trapping, with reduced expiratory flow, thus accounting for the increased dyspnea⁹⁸. There is also worsening of V_A/Q abnormalities, which can result in hypoxemia⁹⁹. Other conditions (pneumonia, thromboembolism, and acute cardiac failure) may mimic or aggravate an exacerbation of COPD.

Systemic Features. It is increasingly recognized that many patients with COPD have comorbidities that have a major impact on quality of life and survival¹⁰⁰. Airflow limitation and particularly hyperinflation affect cardiac function and gas exchange¹⁰¹. Inflammatory mediators in the circulation may contribute to skeletal muscle wasting and cachexia, and may initiate or worsen comorbidities such as ischemic heart disease, heart failure, osteoporosis, normocytic anemia, diabetes, metabolic syndrome, and depression.

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

CHAPTER

2

DIAGNOSIS AND ASSESSMENT

CHAPTER 2: DIAGNOSIS AND ASSESSMENT

KEY POINTS:

- A clinical diagnosis of COPD should be considered in any patient who has dyspnea, chronic cough or sputum production, and a history of exposure to risk factors for the disease.
- Spirometry is required to make the diagnosis in this clinical context; the presence of a post-bronchodilator $FEV_1/FVC < 0.70$ confirms the presence of persistent airflow limitation and thus of COPD.
- The goals of COPD assessment are to determine the severity of the disease, including the severity of airflow limitation, the impact on the patient's health status, and the risk of future events (such as exacerbations, hospital admissions, or death), in order to guide therapy.
- Comorbidities occur frequently in COPD patients, including cardiovascular disease, skeletal muscle dysfunction, metabolic syndrome, osteoporosis, depression, and lung cancer. Given that they can occur in patients with mild, moderate and severe airflow limitation and influence mortality and hospitalizations independently, comorbidities should be actively looked for, and treated appropriately if present.
- Non-surgical bronchoscopic lung volume reduction techniques should not be used outside clinical trials until more data are available.

DIAGNOSIS

A clinical diagnosis of COPD should be considered in any patient who has dyspnea, chronic cough or sputum production, and a history of exposure to risk factors for the disease (**Table 2.1**). Spirometry is required to make the diagnosis in this clinical context⁵⁰⁵; the presence of a post-bronchodilator $FEV_1/FVC < 0.70$ confirms the presence of persistent airflow limitation and thus of COPD.

The spirometric criterion for airflow limitation remains a post-bronchodilator fixed ratio of $FEV_1/FVC < 0.70$. This criterion is simple, independent of reference values, and has been used in numerous clinical trials forming the evidence base from which most of our treatment recommendations are drawn. Diagnostic simplicity and consistency are key for the busy non-specialist clinician.

While post-bronchodilator spirometry is required for the

diagnosis and assessment of severity of COPD, the degree of reversibility of airflow limitation (e.g., measuring FEV_1 before and after bronchodilator or corticosteroids) is no longer recommended⁵⁰⁶. The degree of reversibility has

Table 2.1. Key Indicators for Considering a Diagnosis of COPD

Consider COPD, and perform spirometry, if any of these indicators are present in an individual over age 40. These indicators are not diagnostic themselves, but the presence of multiple key indicators increases the probability of a diagnosis of COPD. Spirometry is required to establish a diagnosis of COPD.

Dyspnea that is: Progressive (worsens over time).
Characteristically worse with exercise.
Persistent.

Chronic cough: May be intermittent and may be unproductive.

Chronic sputum production:
Any pattern of chronic sputum production may indicate COPD.

History of exposure to risk factors:
Tobacco smoke (including popular local preparations).
Smoke from home cooking and heating fuels.
Occupational dusts and chemicals.

Family history of COPD

never been shown to add to the diagnosis, differential diagnosis with asthma, or to predicting the response to long-term treatment with bronchodilators or corticosteroids. The role of screening spirometry in the general population is controversial. Both FEV_1 and FVC predict all-cause mortality independent of tobacco smoking, and abnormal lung function identifies a subgroup of smokers at increased risk for lung cancer. This has been the basis of an argument that screening spirometry should be employed as a global health assessment tool^{102,103}. However, there are no data to indicate that screening spirometry is effective in directing management decisions or in improving COPD outcomes in patients who are identified before the development of significant symptoms¹⁰⁴. Thus, GOLD advocates active case finding⁵⁸⁴ but not screening spirometry.

The use of the fixed FEV_1/FVC ratio to define airflow limitation will result in more frequent diagnosis of COPD in the elderly¹⁰⁵, and less frequent diagnosis in adults younger than 45 years¹⁰⁶, especially of mild disease, compared to using a cutoff based on the lower limit of normal (LLN) values for FEV_1/FVC . These LLN values are based on

the normal distribution and classify the bottom 5% of the healthy population as abnormal. From a scientific perspective it is difficult to determine which of these criteria is correct to diagnose COPD¹⁰⁷, and no studies exist comparing clinical diagnosis based on the two approaches. However, LLN values are highly dependent on the choice of valid reference equations using post-bronchodilator FEV₁, and neither longitudinal studies validating the use of the LLN nor studies using reference equations in populations where smoking is not the major cause of COPD are available. The risk of misdiagnosis and over-treatment of individual patients using the fixed ratio as a diagnostic criterion is limited, as spirometry is only one parameter for establishing the clinical diagnosis of COPD, the others being symptoms and risk factors.

Symptoms

The characteristic symptoms of COPD are chronic and progressive dyspnea, cough, and sputum production that can be variable from day-to-day^{507,508}. Chronic cough and sputum production may precede the development of airflow limitation by many years. Individuals, particularly those exposed to COPD risk factors, who present with these symptoms should be examined to search for an underlying cause(s) and appropriate interventions taken. Conversely, significant airflow limitation may develop without chronic cough and sputum production. Although COPD is defined on the basis of airflow limitation, in practice the decision to seek medical help (and so permit the diagnosis to be made) is usually determined by the impact of a symptom on a patient's daily life. A person may seek medical attention either because of chronic symptoms or because of a first exacerbation.

Dyspnea. Dyspnea, a cardinal symptom of COPD, is a major cause of disability and anxiety associated with the disease. Typical COPD patients describe their dyspnea as a sense of increased effort to breathe, heaviness, air hunger, or gasping¹⁰⁸. However, the terms used to describe dyspnea vary both by individual and by culture¹⁰⁹.

Cough. Chronic cough, often the first symptom of COPD to develop¹¹⁰, is frequently discounted by the patient as an expected consequence of smoking and/or environmental exposures. Initially, the cough may be intermittent, but later is present every day, often throughout the day. The chronic cough in COPD may be unproductive¹¹¹. In some cases, significant airflow limitation may develop without the presence of a cough. **Table 2.2** lists some of the other causes of chronic cough.

Sputum production. COPD patients commonly raise small quantities of tenacious sputum after coughing bouts.

Regular production of sputum for 3 or more months in 2 consecutive years (in the absence of any other conditions that may explain it) is the epidemiological definition of chronic bronchitis¹¹², but this is a somewhat arbitrary definition that does not reflect the range of sputum production in COPD patients. Sputum production is often difficult to evaluate because patients may swallow sputum rather than expectorate it, a habit subject to significant cultural and gender variation. Patients producing large volumes of sputum may have underlying bronchiectasis. The presence of purulent sputum reflects an increase

Table 2.2. Causes of Chronic Cough	
Intrathoracic	<ul style="list-style-type: none"> • Chronic obstructive pulmonary disease • Asthma • Lung cancer • Tuberculosis • Bronchiectasis • Left heart failure • Interstitial lung disease • Cystic fibrosis • Idiopathic cough
Extrathoracic	<ul style="list-style-type: none"> • Chronic allergic rhinitis • Upper Airway Cough Syndrome (UACS) • Gastroesophageal reflux • Medication (e.g., ACE inhibitors)

in inflammatory mediators¹¹³, and its development may identify the onset of a bacterial exacerbation¹¹⁴.

Wheezing and Chest Tightness. Wheezing and chest tightness are nonspecific symptoms that may vary between days, and over the course of a single day. Audible wheeze may arise at a laryngeal level and need not be accompanied by auscultatory abnormalities. Alternatively, widespread inspiratory or expiratory wheezes can be present on listening to the chest. Chest tightness often follows exertion, is poorly localized, is muscular in character, and may arise from isometric contraction of the intercostal muscles. An absence of wheezing or chest tightness does not exclude a diagnosis of COPD, nor does the presence of these symptoms confirm a diagnosis of asthma.

Additional Features in Severe Disease. Fatigue, weight loss and anorexia are common problems in patients with severe and very severe COPD¹¹⁵. They are prognostically important¹¹⁶ and can also be a sign of other diseases (e.g., tuberculosis, lung cancer), and therefore should always be investigated. Cough syncope occurs due to rapid increases in intrathoracic pressure during prolonged attacks of coughing. Coughing spells may also cause rib fractures, which are sometimes asymptomatic. Ankle swelling may be the only symptomatic pointer to the development of cor pulmonale. Symptoms of depression and/or anxiety merit

specific enquiry in the clinical history because they are common in COPD¹¹⁷ and are associated with increased risk of exacerbations and poorer health status.

Medical History

A detailed medical history of a new patient known or thought to have COPD should assess:

- *Patient's exposure to risk factors*, such as smoking and occupational or environmental exposures
- *Past medical history*, including asthma, allergy, sinusitis, or nasal polyps; respiratory infections in childhood; other respiratory diseases
- *Family history of COPD or other chronic respiratory disease*
- *Pattern of symptom development*: COPD typically develops in adult life and most patients are conscious of increased breathlessness, more frequent or prolonged "winter colds," and some social restriction for a number of years before seeking medical help
- *History of exacerbations or previous hospitalizations for respiratory disorder*: Patients may be aware of periodic worsening of symptoms even if these episodes have not been identified as exacerbations of COPD
- *Presence of comorbidities*, such as heart disease, osteoporosis, musculoskeletal disorders, and malignancies that may also contribute to restriction of activity¹¹⁸
- *Impact of disease on patient's life*, including limitation of activity, missed work and economic impact, effect on family routines, feelings of depression or anxiety, well being and sexual activity
- *Social and family support available to the patient*
- *Possibilities for reducing risk factors, especially smoking cessation*

Physical Examination

Although an important part of patient care, a physical examination is rarely diagnostic in COPD. Physical signs of airflow limitation are usually not present until significant impairment of lung function has occurred^{119,120}, and their detection has a relatively low sensitivity and specificity. A number of physical signs may be present in COPD, but their absence does not exclude the diagnosis.

Spirometry

Spirometry is the most reproducible and objective measurement of airflow limitation available. Peak expiratory flow measurement alone cannot be reliably used as the only diagnostic test, despite its good sensitivity, because of its weak specificity¹²¹. Good quality spirometric measurement is possible in any health care setting and all health care workers who care for COPD patients should

have access to spirometry. **Table 2.3** summarizes some of the factors needed to achieve accurate test results.

Spirometry should measure the volume of air forcibly exhaled from the point of maximal inspiration (forced vital capacity, FVC) and the volume of air exhaled during the first second of this maneuver (forced expiratory volume in one second, FEV₁), and the ratio of these two measurements (FEV₁/FVC) should be calculated. The ratio between FEV₁ and slow vital capacity (VC), FEV₁/VC, is sometimes measured instead of the FEV₁/FVC ratio. This will often lead to lower values of the ratio, especially in pronounced airflow limitation; however, the cut-off point of 0.7 should still be applied. Spirometry measurements are evaluated by comparison with reference values¹²² based on age, height, sex, and race.

Table 2.3. Considerations in Performing Spirometry

Preparation

- Spirometers need calibration on a regular basis.
- Spirometers should produce hard copy or have a digital display of the expiratory curve to permit detection of technical errors or have an automatic prompt to identify an unsatisfactory test and the reason for it.
- The supervisor of the test needs training in its effective performance.
- Maximal patient effort in performing the test is required to avoid underestimation of values and hence errors in diagnosis and management.

Bronchodilation

- Possible dosage protocols are 400 mcg beta₂-agonist, 160 mcg anticholinergic, or the two combined¹²². FEV₁ should be measured 10-15 minutes after a short-acting beta₂-agonist is given, or 30-45 minutes after a short-acting anticholinergic or a combination.

Performance

- Spirometry should be performed using techniques that meet published standards¹²³.
- The expiratory volume/time traces should be smooth and free from irregularities.
- The recording should go on long enough for a volume plateau to be reached, which may take more than 15 seconds in severe disease.
- Both FVC and FEV₁ should be the largest value obtained from any of 3 technically satisfactory curves and the FVC and FEV₁ values in these three curves should vary by no more than 5% or 150 ml, whichever is greater.
- The FEV₁/FVC ratio should be taken from the technically acceptable curve with the largest sum of FVC and FEV₁.


Evaluation

- Spirometry measurements are evaluated by comparison of the results with appropriate reference values based on age, height, sex, and race.
- The presence of a postbronchodilator FEV₁/FVC < 0.70 confirms the presence of airflow limitation.

ASSESSMENT OF DISEASE

Figure 2.1A shows a normal spirometry tracing; **Figure 2.1B** a spirometry tracing typical of a patient with obstructive disease. Patients with COPD typically show a decrease in both FEV₁ and FVC.

The goals of COPD assessment are to determine the severity of the disease, its impact on the patient's health status and the risk of future events (such as exacerbations, hospital



admissions or death), in order to, eventually, guide therapy.

To achieve these goals, COPD assessment must consider the following aspects of the disease separately:

- Current level of patient's symptoms
- Severity of the spirometric abnormality
- Exacerbation risk
- Presence of comorbidities

Assessment of Symptoms

In the past, COPD was viewed as a disease largely characterized by breathlessness. A simple measure of breathlessness such as the Modified British Medical Research Council (mMRC) Questionnaire (**Table 2.4**) was considered adequate for assessment of symptoms, as the mMRC relates well to other measures of health status⁹¹ and predicts future mortality risk⁹². However, it is now recognized that COPD has multiple symptomatic effects¹⁵¹. For this reason, a comprehensive symptom assessment is recommended rather than just a measure of breathlessness.

The most comprehensive disease-specific health-related quality of life or health status questionnaires such as the CRQ²³⁶ and SGRQ³⁴⁷ are too complex to use in routine practice, but two shorter comprehensive measures (COPD Assessment Test, CAT and COPD Control Questionnaire, CCQ) have been developed and are suitable.

COPD Assessment Test (CAT). The COPD Assessment Test is an 8-item unidimensional measure of health status impairment in COPD¹²⁴. It was developed to be applicable worldwide and validated translations are available in a wide range of languages. The score ranges from 0-40, correlates very closely with the SGRQ, and has been extensively documented in numerous publications⁵⁴⁸ (<http://www.catestonline.org>).

COPD Control Questionnaire (CCQ). The COPD Control

Table 2.4. Modified Medical Research Council Questionnaire for Assessing the Severity of Breathlessness

PLEASE TICK IN THE BOX THAT APPLIES TO YOU (ONE BOX ONLY)

- | | |
|--|--------------------------|
| mMRC Grade 0. I only get breathless with strenuous exercise. | <input type="checkbox"/> |
| mMRC Grade 1. I get short of breath when hurrying on the level or walking up a slight hill. | <input type="checkbox"/> |
| mMRC Grade 2. I walk slower than people of the same age on the level because of breathlessness, or I have to stop for breath when walking on my own pace on the level. | <input type="checkbox"/> |
| mMRC Grade 3. I stop for breath after walking about 100 meters or after a few minutes on the level. | <input type="checkbox"/> |
| mMRC Grade 4. I am too breathless to leave the house or I am breathless when dressing or undressing. | <input type="checkbox"/> |

Questionnaire is a 10 item self-administered questionnaire developed to measure clinical control in patients with COPD^{509, 510}. Although the concept of "control" in COPD remains controversial, the CCQ is short and easy to administer. It is reliable and responsive, is available in a range of languages, and has been validated (<http://www.ccq.nl>). A minimally clinically important differences (MCID) during rehabilitation of -0.4 for the CCQ has been identified⁶¹⁷.


Choice of Cut Points

The CAT and CCQ provide a measure of the symptomatic impact of COPD but do not categorize patients into lower and higher symptoms for the purpose of treatment. The SGRQ is the most widely documented comprehensive measure; scores less than 25 are uncommon in diagnosed COPD patients^{131,549} and scores ≥ 25 are very uncommon

Table 2.5. Classification of Severity of Airflow Limitation in COPD (Based on Post-Bronchodilator FEV₁)

In patients with FEV ₁ /FVC < 0.70:		
GOLD 1:	Mild	FEV ₁ ≥ 80% predicted
GOLD 2:	Moderate	50% ≤ FEV ₁ < 80% predicted
GOLD 3:	Severe	30% ≤ FEV ₁ < 50% predicted
GOLD 4:	Very Severe	FEV ₁ < 30% predicted

Figure 2.2. Relationship Between Health-related Quality of Life, Post-bronchodilator FEV₁ and GOLD Spirometric Classification (Adapted from Jones¹²⁷)



in healthy persons⁵⁴⁹. In clinical trials of long-acting bronchodilator medications^{201,210, 516, 550-555}, the baseline weighted mean SGRQ score was 44, and one standard deviation below the mean was 26. Therefore, it is recommended that a symptom score equivalent to SGRQ score ≥ 25 should be used as the cut-point for considering regular treatment for symptoms including breathlessness, particularly since this corresponds to the range of severity seen in patients recruited to the trials that provide the evidence base for treatment recommendations. The equivalent cut-point for the CAT is 10^{123, 554}. The equivalent cut-point for the CCQ has yet to be finally determined, but appears to be in the range 1.0 - 1.5.

An equivalent mMRC score cannot be calculated because a simple breathlessness cut-point cannot equate to a comprehensive symptom score cut-point. The great majority of patients with an SGRQ of 25 or more will have an mMRC of 2 or more; however patients with mMRC < 2 may also have a number of other COPD symptoms. While use of an mMRC ≥ 2 as a cut-point

may be adequate for breathlessness assessment, it will also categorize a number of patients with symptoms other than breathlessness as having “few symptoms.” For this reason, the use of a comprehensive symptom assessment is recommended. However, because use of the mMRC is still widespread, an mMRC of ≥ 2 is still included as a cut-point for separating “less breathlessness” from “more breathlessness.” However, users are cautioned that assessment of other symptoms is required^{554,555}.

Spirometric Assessment

Table 2.5 shows the classification of airflow limitation severity in COPD. Specific spirometric cut-points are used for purposes of simplicity. Spirometry should be performed after the administration of an adequate dose of a short-acting inhaled bronchodilator in order to minimize variability.

However, there is only a weak correlation between FEV₁, symptoms and impairment of a patient’s health-related quality of life. This is illustrated in **Figure 2.2** in which health-related quality of life is plotted against post-bronchodilator FEV₁^{126,127} with the GOLD spirometric classification superimposed. The figure illustrates that, within any given category, patients may have anything between relatively well preserved to very poor health status. For this reason, formal symptomatic assessment is also required.

Assessment of Exacerbation Risk

An exacerbation of COPD is defined as an acute event characterized by a worsening of the patient’s respiratory symptoms that is beyond normal day-to-day variations and leads to a change in medication¹²⁸⁻¹³⁰. The rate at which exacerbations occur varies greatly between patients^{131,512}. The best predictor of having frequent exacerbations (2 or more exacerbations per year) is a history of previous treated events¹³². In addition, worsening airflow limitation is associated with an increasing prevalence of exacerbations and risk of death. Hospitalization for a COPD exacerbation is associated with a poor prognosis with increased risk of death⁵⁵⁶.

A large body of data has been accumulated in patients^{131,152} classified using GOLD spirometric grading systems. These show an increase in risk of exacerbations, hospitalization and death with worsening of airflow limitation. The data in **Table 2.6** are derived from prospectively collected data from large medium-term clinical trials¹³²⁻¹³⁴. They are not precise estimates that apply to each patient, but they illustrate clearly the increased risk of exacerbations and death between spirometric levels. Roughly, although up to

20% of GOLD 2 (Moderate airflow limitation) patients may experience frequent exacerbations requiring treatment with antibiotics and/or systemic corticosteroids¹³², the risk of exacerbations significantly increases in GOLD 3 (Severe) and GOLD 4 (Very Severe). Since exacerbations increase the decline in lung function, deterioration in health status and risk of death, the assessment of exacerbation risk can also be seen as an assessment of the risk of poor outcomes in general.

Table 2.6: RISK IN COPD: Placebo-limb data from TORCH^{134*}, Uplift^{133†} and Eclipse^{132‡}

GOLD spirometric level	Exacerbations (per year)*†‡	Hospitalizations (per year)* ‡	3-year Mortality*†
GOLD 1: Mild	?	?	?
GOLD 2: Moderate	0.7 – 0.9	0.11 – 0.2	11%*†
GOLD 3: Severe	1.1 – 1.3	0.25 – 0.3	15%*
GOLD 4: Very severe	1.2 – 2.0	0.4 – 0.54	24%*


*Toward a Revolution in COPD Health (TORCH) study¹³⁴
 † Understanding Potential Long-Term Impacts on Function with Tiotropium (UPLIFT) study¹³³
 ‡ Evaluation of COPD Longitudinally to Identify Predictive Surrogate Endpoints (ECLIPSE) study¹³²

Assessment of Comorbidities

Because COPD often develops in long-time smokers in middle age, patients frequently have a variety of other diseases related to either smoking or aging¹³⁵. COPD itself also has significant extrapulmonary (systemic) effects including weight loss, nutritional abnormalities and skeletal muscle dysfunction. The latter is characterized by both sarcopenia (loss of muscle cells) and abnormal function of the remaining cells¹³⁶. Its causes are likely multifactorial (inactivity, poor diet, inflammation, hypoxia) and it can contribute to exercise intolerance and poor health status in patients with COPD. Importantly, skeletal muscle dysfunction is a remediable source of exercise intolerance¹³⁷.

Comorbidities that occur frequently in COPD patients include cardiovascular disease, skeletal muscle dysfunction, metabolic syndrome, osteoporosis, depression and lung cancer. The existence of COPD may actually increase the risk for other diseases; this is particularly striking for COPD and lung cancer¹³⁸⁻¹⁴¹. Whether this association is due to common risk factors (e.g., smoking), involvement of susceptibility genes, or impaired clearance of carcinogens is not clear. Comorbidities can occur in patients with mild, moderate or severe airflow limitation¹³¹, influence mortality

and hospitalizations independently¹⁴², and deserve specific treatment. Therefore, comorbidities should be looked for routinely, and treated appropriately, in any patient with COPD. The guidelines for the diagnosis, assessment of severity, and management of individual comorbidities in



patients with COPD are the same as for all other patients. A more detailed description of the management of COPD and comorbidities is given in Chapter 6.

Combined COPD Assessment

An understanding of the impact of COPD on an individual patient combines the symptomatic assessment with the patient's spirometric classification and/or risk of exacerbations. This approach to combined assessment is illustrated in **Figure 2.3**.

As detailed above, the CAT is recommended as a comprehensive measure of symptoms, with a CAT score ≥ 10 indicating a high level of symptoms. Comprehensive assessment of the symptomatic impact of the disease is preferred, but in its absence mMRC scores provide an assessment of the impact of dyspnea. It is unnecessary and possibly confusing to use more than one scale.

There are three methods of assessing exacerbation risk. One is a population-based method using the GOLD spirometric classification (**Table 2.5**), with GOLD 3 or GOLD 4 categories indicating high risk. The second based on the individual patient's history of exacerbations¹³²,

with two or more exacerbations in the preceding year indicating high risk. The third is a history of hospitalization due to an exacerbation in the preceding year. (If there is a discrepancy between these criteria, the assessment pointing to the highest risk should be used.) To use **Figure 2.3**, first assess symptoms with the CAT scale (or dyspnea with the mMRC) and determine if the patient belongs to the boxes on the left side – Less Symptoms (CAT < 10) or Less Breathlessness (mMRC grade 0-1); or belongs to boxes on the right side - More Symptoms (CAT ≥ 10) or More Breathlessness (mMRC grade ≥ 2).

Next assess the risk of exacerbations to determine if the patient belongs to the lower part of the box – Low Risk – or the upper part of the box – High Risk. This can be done by one of three methods: (1) use spirometry to determine the GOLD grade of airflow limitation (GOLD 1 and GOLD 2 categories indicate Low Risk, while GOLD 3 and GOLD 4 indicate High Risk); (2) assess the number of exacerbations the patient has had within the previous 12 months (0 or 1 indicates Low Risk, while 2 or more exacerbations indicates High Risk); (3) determine whether the patient has had one or more hospitalization in the previous year for a COPD exacerbation. In some patients, these three ways of assessing risk of exacerbations will not lead to the same level of risk; in this case, the risk should be determined by the method indicating High Risk.

Example: Imagine a patient with a CAT score of 18, FEV₁ of 55% of predicted, and a history of 3 exacerbations within the last 12 months. Symptom assessment using CAT shows that the patient is More Symptomatic (CAT ≥ 10) and is therefore either Group B or Group D. Spirometry indicates Low Risk as the patient is GOLD 2 (Moderate airflow limitation) but as the patient had 3 exacerbations within the last 12 months this indicates High Risk and outweighs the lower risk assessment based on spirometry. The patient therefore belongs in Group D.

The groups can be summarized as follows:

- **Patient Group A – Low Risk, Less Symptoms**
Typically GOLD 1 or GOLD 2 (Mild or Moderate airflow limitation); and/or 0-1 exacerbation per year *and no* hospitalization for exacerbation; *and* CAT score < 10 *or* mMRC grade 0-1
- **Patient Group B – Low Risk, More Symptoms**
Typically GOLD 1 or GOLD 2 (Mild or Moderate airflow limitation); and/or 0-1 exacerbation per year *and no* hospitalization for exacerbation; *and* CAT score ≥ 10 *or* mMRC grade ≥ 2
- **Patient Group C – High Risk, Less Symptoms**
Typically GOLD 3 or GOLD 4 (Severe or Very Severe airflow limitation); and/or ≥ 2 exacerbations per year *or* ≥ 1 with hospitalization for exacerbation; *and* CAT

score < 10 *or* mMRC grade 0-1

- **Patient Group D – High Risk, More Symptoms**
Typically GOLD 3 or GOLD 4 (Severe or Very Severe airflow limitation); and/or ≥ 2 exacerbations per year *or* ≥ 1 with hospitalization for exacerbation; *and* CAT score ≥ 10 *or* mMRC grade ≥ 2

Evidence to support this classification system includes:

- Patients with a high risk of exacerbations tend to be in GOLD categories 3 and 4 (Severe or Very Severe airflow limitation, **Figure 2.3**) and can be identified quite reliably from their own past history¹³².
- Higher exacerbation rates are associated with faster loss of FEV₁¹⁴³ and greater worsening of health status¹⁴⁴.
- Hospitalization for a COPD exacerbation is associated with a poor prognosis⁵⁵⁶.
- CAT scores ≥ 10 are associated with significantly impaired health status¹⁴⁵.

Even in the absence of frequent exacerbations, patients in GOLD categories 3 and 4 may be at greater risk of hospital admission and death (**Figure 2.3**). These important increased risks form the rationale for including such patients in the “High Risk” groups.

This approach, combined with an assessment of potential comorbidities, reflects the complexity of COPD better than the unidimensional analysis of airflow limitation previously used for staging the disease and forms the basis of the guide to individualized management provided in Chapter 4.

Additional Investigations

The following additional investigations may be considered as part of the diagnosis and assessment of COPD:

Imaging. A chest X-ray is not useful to establish a diagnosis in COPD, but it is valuable in excluding alternative diagnoses and establishing the presence of significant comorbidities such as concomitant respiratory (pulmonary fibrosis, bronchiectasis, pleural diseases), skeletal (e.g., kyphoscoliosis), and cardiac diseases (e.g., cardiomegaly). Radiological changes associated with COPD include signs of lung hyperinflation (flattened diaphragm on the lateral chest film, and an increase in the volume of the retrosternal air space), hyperlucency of the lungs, and rapid tapering of the vascular markings. Computed tomography (CT) of the chest is not routinely recommended. However, when there is doubt about the diagnosis of COPD, CT scanning might help in the differential diagnosis where concomitant diseases are

present. In addition, if a surgical procedure such as lung volume reduction is contemplated, a chest CT scan is necessary since the distribution of emphysema is one of the most important determinants of surgical suitability¹⁴⁶.

Lung Volumes and Diffusing Capacity. COPD patients exhibit gas trapping (a rise in residual volume) from early in the disease, and as airflow limitation worsens static hyperinflation (an increase in total lung capacity) occurs. These changes can be documented by body plethysmography, or less accurately by helium dilution lung volume measurement. These measurements help characterize the severity of COPD but are not essential to patient management. Measurement of diffusing capacity (DL_{CO}) provides information on the functional impact of emphysema in COPD and is often helpful in patients with breathlessness that may seem out of proportion with the degree of airflow limitation.

Oximetry and Arterial Blood Gas Measurement. Pulse oximetry can be used to evaluate a patient's oxygen saturation and need for supplemental oxygen therapy. Pulse oximetry should be used to assess all stable patients with $FEV_1 < 35\%$ predicted or with clinical signs suggestive of respiratory failure or right heart failure. If peripheral saturation is $< 92\%$ arterial blood gases should be assessed¹⁴⁷.

Alpha-1 Antitrypsin Deficiency Screening. The World Health Organization recommends that COPD patients from areas with a particularly high prevalence of alpha-1 antitrypsin deficiency should be screened for this genetic disorder¹⁴⁸. However, the typical patient tends to present at a younger age (< 45 years) with lower lobe emphysema. Family members can be identified and family screening is useful for appropriate counseling. A serum concentration of alpha-1 antitrypsin below 15-20% of the normal value is highly suggestive of homozygous alpha-1 antitrypsin deficiency.

Exercise Testing. Objectively measured exercise impairment, assessed by a reduction in self-paced walking distance¹⁴⁹ or during incremental exercise testing in a laboratory¹⁵⁰, is a powerful indicator of health status impairment and predictor of prognosis¹⁵¹; exercise capacity may fall in the year before death⁵⁵⁷. Walking tests can be useful for assessing disability and are used to assess the effectiveness of pulmonary rehabilitation. Both the paced shuttle walk tests^{152,153} and the unpaced 6-minute walk test can be used¹⁵⁴. As the course length has a substantial impact on the distance walked, existing reference equations established for a 30 m course cannot be applied to predict the distance achieved on shorter courses⁵⁸⁵. Laboratory testing using cycle or treadmill ergometry can

identify co-existing or alternative conditions, e.g., cardiac diagnoses. Monitoring of physical activity may be more relevant regarding prognosis than evaluating exercise capacity¹⁵⁵. This can be done using accelerometers or multisensor instruments.

Composite Scores. Several variables including FEV_1 , exercise tolerance assessed by walking distance or peak oxygen consumption, weight loss, and reduction in arterial oxygen tension identify patients at increased risk for mortality. A relatively simple approach to identifying disease severity using a combination of most of the above variables has been proposed. The BODE method gives a composite score (Body mass index, Obstruction, Dyspnea, and Exercise) that is a better predictor of subsequent survival than any component singly¹⁵⁶, and its properties as a measurement tool are under investigation. Simpler alternatives not including an exercise test have been suggested but all these approaches need validation across a wide range of disease severities and in different clinical settings to confirm that they are suitable for routine clinical use^{157,158}.

DIFFERENTIAL DIAGNOSIS

In some patients with chronic asthma, a clear distinction from COPD is not possible using current imaging and physiological testing techniques, and it is assumed that asthma and COPD coexist in these patients. In these cases, current management will include use of anti-inflammatory drugs and other treatments need to be individualized. Other potential diagnoses are usually easier to distinguish from COPD (**Table 2.7**).

Table 2.7. COPD and its Differential Diagnoses

Diagnosis	Suggestive Features
COPD	Onset in mid-life. Symptoms slowly progressive. History of tobacco smoking or exposure to other types of smoke.
Asthma	Onset early in life (often childhood). Symptoms vary widely from day to day. Symptoms worse at night/early morning. Allergy, rhinitis, and/or eczema also present. Family history of asthma.
Congestive Heart Failure	Chest X-ray shows dilated heart, pulmonary edema. Pulmonary function tests indicate volume restriction, not airflow limitation.
Bronchiectasis	Large volumes of purulent sputum. Commonly associated with bacterial infection. Chest X-ray/CT shows bronchial dilation, bronchial wall thickening.
Tuberculosis	Onset all ages. Chest X-ray shows lung infiltrate. Microbiological confirmation. High local prevalence of tuberculosis.
Obliterative Bronchiolitis	Onset at younger age, nonsmokers. May have history of rheumatoid arthritis or acute fume exposure. Seen after lung or bone marrow transplantation. CT on expiration shows hypodense areas.
Diffuse Panbronchiolitis	Predominantly seen in patients of Asian descent. Most patients are male and nonsmokers. Almost all have chronic sinusitis. Chest X-ray and HRCT show diffuse small centrilobular nodular opacities and hyperinflation.
<p><i>These features tend to be characteristic of the respective diseases, but are not mandatory. For example, a person who has never smoked may develop COPD (especially in the developing world where other risk factors may be more important than cigarette smoking); asthma may develop in adult and even in elderly patients.</i></p>	

CHAPTER

3

***THERAPEUTIC
OPTIONS***

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

CHAPTER 3: THERAPEUTIC OPTIONS

KEY POINTS:

- In patients who smoke, smoking cessation is very important. Pharmacotherapy and nicotine replacement reliably increase long-term smoking abstinence rates.
- Appropriate pharmacologic therapy can reduce COPD symptoms, reduce the frequency and severity of exacerbations, and improve health status and exercise tolerance.
- To date, none of the existing medications for COPD has been shown conclusively to modify the long-term decline in lung function.
- Each pharmacological treatment regimen needs to be patient-specific, guided by severity of symptoms, risk of exacerbations, drug availability, and the patient's response.
- Influenza and pneumococcal vaccination should be offered to every COPD patient; they appear to be more effective in older patients and those with more severe disease or cardiac comorbidity.
- All patients who get short of breath when walking on their own pace on level ground should be offered rehabilitation; it can improve symptoms, quality of life, and physical and emotional participation in everyday activities.
- Non-surgical bronchoscopic lung volume reduction techniques should not be used outside clinical trials until more data are available.

SMOKING CESSATION

Smoking cessation is the intervention with the greatest capacity to influence the natural history of COPD. Evaluation of the smoking cessation component in a long-term, multicenter study indicates that if effective resources and time are dedicated to smoking cessation, 25% long-term quit rates can be achieved¹⁵⁹.

Pharmacotherapies for Smoking Cessation

Nicotine Replacement Products. Nicotine replacement therapy in any form (nicotine gum, inhaler, nasal spray, transdermal patch, sublingual tablet, or lozenge) reliably increases long-term smoking abstinence rates¹⁶⁰⁻¹⁶² and is significantly more effective than placebo. Patients need to be informed about the proper use of these products to optimize efficacy. Medical contraindications to nicotine replacement therapy include unstable coronary artery disease, untreated peptic ulcer disease, and recent myocardial infarction or stroke¹⁶³. Continuous chewing of nicotine gum produces secretions that are swallowed rather than absorbed through

the buccal mucosa, results in little absorption, and can cause nausea. Acidic beverages, particularly coffee, juices, and soft drinks, interfere with the absorption of nicotine.

Pharmacologic. Varenicline¹⁶⁴, bupropion¹⁶⁵ and nortriptyline have been shown to increase long-term quit rates^{161,163,166}, but should always be used as one element in a supportive intervention program rather than on their own. Although more studies need to be conducted with these medications, a randomized controlled trial with counseling and support showed quit rates at one year of 30% with sustained-release bupropion alone and 35% with sustained-release bupropion plus nicotine patch¹⁶⁵. The effectiveness of the antihypertensive drug clonidine is limited by side effects¹⁶¹.

Table 3.1. Treating Tobacco Use and Dependence: A Clinical Practice Guideline—Major Findings and Recommendations¹⁶⁰

1. Tobacco dependence is a chronic condition that warrants repeated treatment until long-term or permanent abstinence is achieved.
2. Effective treatments for tobacco dependence exist and all tobacco users should be offered these treatments.
3. Clinicians and health care delivery systems must institutionalize the consistent identification, documentation, and treatment of every tobacco user at every visit.
4. Brief smoking cessation counseling is effective and every tobacco user should be offered such advice at every contact with health care providers.
5. There is a strong dose-response relation between the intensity of tobacco dependence counseling and its effectiveness.
6. Three types of counseling have been found to be especially effective: practical counseling, social support as part of treatment, and social support arranged outside of treatment.
7. First-line pharmacotherapies for tobacco dependence—varenicline, bupropion SR, nicotine gum, nicotine inhaler, nicotine nasal spray, and nicotine patch—are effective and at least one of these medications should be prescribed in the absence of contraindications.
8. Tobacco dependence treatments are cost effective relative to other medical and disease prevention interventions.

Recommendations for treating tobacco use and dependence are summarized in **Table 3.1**.

A five-step program for intervention (**Table 3.2**) provides a strategic framework helpful to health care providers interested in helping their patients stop smoking^{160,167-169}. Because tobacco dependence is a chronic disease¹⁶⁰, clinicians should recognize that relapse is common and reflects the chronic nature of dependence and addiction, not failure on the part of the clinician or the patient.

Counseling delivered by physicians and other health professionals significantly increases quit rates over self-initiated strategies¹⁷⁰ (**Evidence A**). Even a brief (3-minute)

period of counseling to urge a smoker to quit results in smoking cessation rates of 5-10%¹⁷¹. There is a strong dose-response relationship between counseling intensity and cessation success^{172,173}. Ways to intensify treatment include increasing the length of the treatment session, the number of treatment sessions, and the number of weeks over which the treatment is delivered. Sustained quit rates of 10.9% at 6 months have been achieved when clinician tutorials and feedback are linked to counseling sessions¹⁷⁴. With more complex interventions quit rates can reach 20-30%¹⁷². In a multicenter controlled clinical trial, a combination of physician advice, group support,

needs to be patient-specific as the relationship between severity of symptoms, airflow limitation, and severity of exacerbations will differ between patients. When treatment is given by the inhaled route, attention to effective drug delivery and training in inhaler technique is essential¹⁷⁷. The choice of inhaler device will depend on availability, cost, the prescribing physician, and the skills and ability of the patient. COPD patients may have problems with coordination and find it hard to use a metered-dose inhaler (MDI). It is essential to ensure that inhaler technique is correct and to re-check this at each visit.

Table 3.2. Brief Strategies to Help the Patient Willing to Quit^{160,167-169}

1. **ASK:** Systematically identify all tobacco users at every visit. *Implement an office-wide system that ensures that, for EVERY patient at EVERY clinic visit, tobacco-use status is queried and documented.*
2. **ADVISE:** Strongly urge all tobacco users to quit. *In a clear, strong, and personalized manner, urge every tobacco user to quit.*
3. **ASSESS:** Determine willingness to make a quit attempt. *Ask every tobacco user if he or she is willing to make a quit attempt at this time (e.g., within the next 30 days).*
4. **ASSIST:** Aid the patient in quitting. *Help the patient with a quit plan; provide practical counseling; provide intra-treatment social support; help the patient obtain extra-treatment social support; recommend use of approved pharmacotherapy except in special circumstances; provide supplementary materials.*
5. **ARRANGE:** Schedule follow-up contact. *Schedule follow-up contact, either in person or via telephone.*

PHARMACOLOGIC THERAPY FOR STABLE COPD

skills training, and nicotine replacement therapy achieved a quit rate of 35% at 1 year and a sustained quit rate of 22% at 5 years¹⁵⁹.

Overview of the Medications

Pharmacologic therapy for COPD is used to reduce symptoms, reduce the frequency and severity of exacerbations, and improve health status and exercise tolerance. To date, none of the existing medications for COPD has been conclusively shown to modify the long-term decline in lung function when this is tested as a primary or secondary outcome in clinical trials^{126,159,175,176}. *Post-hoc* evidence of such an effect with long-acting bronchodilators and/or inhaled corticosteroids^{133,143} requires confirmation in specifically designed trials.

The classes of medications commonly used in treating COPD are shown in **Table 3.3**. The choice within each class depends on the availability and cost of medication and the patient's response. Each treatment regimen

Alternative breath-activated or spacer devices are available. In general, particle deposition from dry powder inhalers (DPIs) will tend to be more central with the fixed airflow limitation and lower inspiratory flow rates in COPD^{178,179}. However, as has been shown in asthma, patients are also likely to find the use of some dry powder inhalers difficult. For the MDI, the addition of a large or small volume spacer often overcomes coordination problems, and improves lower airway deposition and clinical benefit. Many drugs are available as nebulizer solutions and, for patients who are severely overinflated and consequently may have very low inspiratory flow rates, there may be theoretical advantages of nebulizer use. However, there is little randomized trial evidence for their benefit over other devices, and use of nebulizers will often depend on local preference, availability and price. Benefit should be judged symptomatically, since changes in lung function may be small and within the limits of repeatability. Nebulized treatment should only be continued if the patient reports clear symptomatic benefit that cannot be achieved by simpler, cheaper, and more portable alternatives.

Bronchodilators

Medications that increase the FEV₁ or change other spirometric variables, usually by altering airway smooth muscle tone, are termed bronchodilators¹⁸⁰, since the improvements in expiratory flow reflect widening of the airways rather than changes in lung elastic recoil. Such medications improve emptying of the lungs, tend to reduce dynamic hyperinflation at rest and during exercise^{181,182}, and improve exercise performance. The extent of these changes, especially in severe and very severe patients, is not easily predictable from the improvement in FEV₁^{183,184}.

Dose-response relationships using FEV₁ as the outcome are relatively flat with all classes of bronchodilators¹⁸⁵⁻¹⁸⁸. Toxicity is also dose-related. Increasing the dose of either a beta₂-agonist or an anticholinergic by an order of magnitude, especially when given by a nebulizer, appears to provide subjective benefit in acute episodes¹⁸⁹ but is not necessarily helpful in stable disease¹⁹⁰.

Table 3.3. Formulations and Typical Doses of COPD Medications*

Drug	Inhaler (mcg)	Solution for Nebulizer (mg/ml)	Oral	Vials for Injection (mg)	Duration of Action (hours)
Beta₂-agonists					
<i>Short-acting</i>					
Fenoterol	100-200 (MDI)	1	0.05% (Syrup)		4-6
Levalbuterol	45-90 (MDI)	0.21, 0.42			6-8
Salbutamol (albuterol)	100, 200 (MDI & DPI)	5	5 mg (Pill), 0.024%(Syrup)	0.1, 0.5	4-6
Terbutaline	400, 500 (DPI)		2.5, 5 mg (Pill)		4-6
<i>Long-acting</i>					
Formoterol	4.5-12 (MDI & DPI)	0.01¶			12
Arformoterol		0.0075			12
Indacaterol	75-300 (DPI)				24
Olodaterol	5 (SMI)				24
Salmeterol	25-50 (MDI & DPI)				12
Tulobuterol			2 mg (transdermal)		24
Anticholinergics					
<i>Short-acting</i>					
Ipratropium bromide	20, 40 (MDI)	0.25-0.5			6-8
Oxipropium bromide	100 (MDI)	1.5			7-9
<i>Long-acting</i>					
Aclidinium bromide	322 (DPI)				12
Glycopyrronium bromide	44 (DPI)				24
Tiotropium	18 (DPI), 5 (SMI)				24
Umeclidinium	62.5 (DPI)				24
Combination short-acting beta₂-agonist plus anticholinergic in one inhaler					
Fenoterol/Ipratropium	200/80 (MDI)	1.25/0.5			6-8
Salbutamol/Ipratropium	100/20 (SMI)				6-8
Combination long-acting beta₂-agonist plus anticholinergic in one inhaler					
Formoterol/aclidinium	12/340 (DPI)				12
Indacaterol/ glycopyrronium	85/43 (DPI)				24
Olodaterol/tiotropium	5/5 (SMI)				24
Vilanterol/umeclidinium	25/62.5 (DPI)				24
Methylxanthines					
Aminophylline			200-600 mg (Pill)	240	Variable, up to 24
Theophylline (SR)			100-600 mg (Pill)		Variable, up to 24
Inhaled corticosteroids					
Beclomethasone	50-400 (MDI & DPI)	0.2-0.4			
Budesonide	100, 200, 400 (DPI)	0.20, 0.25, 0.5			
Fluticasone	50-500 (MDI & DPI)				
Combination long-acting beta₂-agonists plus corticosteroids in one inhaler					
Formoterol/beclomethasone	6/100 (MDI & DPI)				
Formoterol/budesonide	4.5/160 (MDI) 9/320 (DPI)				
Formoterol/mometasone	10/200, 10/400 (MDI)				
Salmeterol/Fluticasone	50/100, 250, 500 (DPI)				
Vilanterol/Fluticasone furoate	25/100 (DPI)				
Systemic corticosteroids					
Prednisone			5-60 mg (Pill)		
Methyl-prednisolone			4, 8, 16 mg (Pill)		
Phosphodiesterase-4 inhibitors					
Roflumilast			500 mcg (Pill)		24

22 THERAPEUTIC OPTIONS

MDI=metered dose inhaler; DPI=dry powder inhaler; SMI=soft mist inhaler

*Not all formulations are available in all countries; in some countries, other formulations may be available.

¶Formoterol nebulized solution is based on the unit dose vial containing 20 mcg in a volume of 2.0 ml

Bronchodilator medications are given on either an as-needed basis or a regular basis to prevent or reduce symptoms¹⁸⁵⁻¹⁸⁸ (**Evidence A**) (**Table 3.4**).

Beta₂-agonists. The principal action of beta₂-agonists is to relax airway smooth muscle by stimulating beta₂-adrenergic receptors, which increases cyclic AMP and produces functional antagonism to bronchoconstriction. The bronchodilator effects of short-acting beta₂-agonists usually wear off within 4 to 6 hours^{191,192}. Regular and as-needed use of short-acting beta-agonists improve FEV₁ and symptoms¹⁹³ (**Evidence B**). The use of high doses of short-acting beta₂-agonists on an as-needed basis in patients already treated with long-acting bronchodilators is not supported by evidence, may be limited by side effects, and cannot be recommended. For single-dose, as-needed

Table 3.4. Bronchodilators in Stable COPD

- Bronchodilator medications are central to symptom management in COPD.
- Inhaled therapy is preferred.
- The choice between beta₂-agonist, anticholinergic, theophylline, or combination therapy depends on availability and individual patient response in terms of symptom relief and side effects.
- Bronchodilators are prescribed on an as-needed or on a regular basis to prevent or reduce symptoms.
- Long-acting inhaled bronchodilators are convenient and more effective at producing maintained symptom relief than short-acting bronchodilators.
- Combining bronchodilators of different pharmacological classes may improve efficacy and decrease the risk of side effects compared to increasing the dose of a single bronchodilator.

use in COPD, there appears to be no advantage in using levalbuterol over conventional bronchodilators¹⁹⁴.

Long-acting inhaled beta₂-agonists show duration of action of 12 or more hours. Formoterol and salmeterol significantly improve FEV₁ and lung volumes, dyspnea, health-related quality of life and exacerbation rate¹⁹⁵⁻²⁰⁰ (**Evidence A**), but have no effect on mortality and rate of decline of lung function. A systematic review of trials of salmeterol and formoterol showed a significant reduction in the numbers of patients requiring treatment for exacerbations and the number requiring hospitalization⁵⁸⁶. Salmeterol reduces the rate of hospitalization¹⁹⁵ (**Evidence B**). Indacaterol is a once daily beta₂-agonist with a duration of action of 24 hours^{201,202}. The bronchodilator effect is significantly greater than that of formoterol and salmeterol, and similar to tiotropium (**Evidence A**). Indacaterol has significant effects on breathlessness, health status and exacerbation rate (**Evidence B**). Its safety profile is similar to placebo; in clinical trials a significant number of patients (24% vs 7%) experienced cough following the inhalation of indacaterol⁵¹³⁻⁵¹⁶.

Adverse effects. Stimulation of beta₂-adrenergic receptors can produce resting sinus tachycardia and has the potential to precipitate cardiac rhythm disturbances in susceptible patients, although these seem to have remarkably few clinical implications. Exaggerated somatic tremor is troublesome in some older patients treated with higher doses of beta₂-agonists, whatever the route of administration, and this limits the dose that can be tolerated. Although hypokalemia can occur, especially when treatment is combined with thiazide diuretics²⁰³, and oxygen consumption can be increased under resting conditions²⁰⁴, these metabolic effects show tachyphylaxis unlike the bronchodilator actions. Mild falls in PaO₂ can occur after administration of both short- and long-acting beta₂-agonists^{205,206} but the clinical significance of these changes is doubtful. Despite the concerns raised some years ago related to beta₂-agonists in the management of asthma, further detailed study has found no association between beta₂-agonist use and an accelerated loss of lung function or increased mortality in COPD.

Anticholinergics. The most important effect in COPD patients of anticholinergic medications, such as ipratropium, oxitropium and tiotropium bromide, appears to be blockage of acetylcholine's effect on muscarinic receptors. Current short-acting drugs block M2 and M3 receptors and modify transmission at the pre-ganglionic junction, although these effects appear less important in COPD²⁰⁷. The long-acting anticholinergic tiotropium has a pharmacokinetic selectivity for the M3 and M1 receptors²⁰⁸. The bronchodilating effect of short-acting inhaled anticholinergics lasts longer than that of short-acting beta₂-agonists, with some bronchodilator effect generally apparent up to 8 hours after administration¹⁹¹.

Among long-acting anticholinergics, aclidinium has a duration of at least 12 hours⁵⁵² whereas tiotropium and glycopyrronium have a duration of action of more than 24 hours²⁰⁹⁻²¹¹. Tiotropium reduces exacerbations and related hospitalizations, improves symptoms and health status^{212, 587} (**Evidence A**), and improves the effectiveness of pulmonary rehabilitation²¹³ (**Evidence B**). In a large, long-term clinical trial on patients with COPD, there was no effect of tiotropium added to other standard therapies on the rate of lung function decline and no evidence of cardiovascular risk²¹⁴. In another large trial, tiotropium was superior to salmeterol in reducing exacerbations although the difference was small^{215,517}. The long-acting anticholinergics aclidinium⁵⁸⁸ and glycopyrronium seem to have similar action on lung function and breathlessness as tiotropium, whereas far less data are available for other outcomes^{552, 558}.

Adverse effects. Anticholinergic drugs are poorly absorbed which limits the troublesome systemic effects seen with atropine²¹⁶. Extensive use of this class of inhaled agents

in a wide range of doses and clinical settings has shown them to be very safe. The main side effect is dryness of the mouth. Twenty-one days of inhaled tiotropium, 18 mcg/day as a dry powder, does not retard mucus clearance from the lungs¹⁴⁴. Although occasional prostatic symptoms have been reported, there are no data to prove a true causal relationship. Some patients using ipratropium report a bitter, metallic taste. An unexpected small increase in cardiovascular events in COPD patients regularly treated with ipratropium bromide has been reported and requires further investigation^{217,218}. Tiotropium delivered via the Respimat[®] soft mist inhaler was associated with a significantly increased risk of mortality compared with placebo in a meta-analysis⁵¹⁹; however, the findings of the TIOSPIR[®] trial showed that there was no difference in mortality or rates of exacerbation when comparing tiotropium in a dry-powder inhaler to the Respimat[®] inhaler⁵⁵⁹. Use of solutions with a face mask has been reported to precipitate acute glaucoma, probably by a direct effect of the solution on the eye.

Methylxanthines. Controversy remains about the exact effects of xanthine derivatives. They may act as nonselective phosphodiesterase inhibitors, but have also been reported to have a range of non-bronchodilator actions, the significance of which is disputed²²⁰⁻²²⁴. Data on duration of action for conventional, or even slow-release, xanthine preparations are lacking in COPD.

Theophylline, the most commonly used methylxanthine, is metabolized by cytochrome P450 mixed function oxidases. Clearance of the drug declines with age. Many other physiological variables and drugs modify theophylline metabolism. Changes in inspiratory muscle function have been reported in patients treated with theophylline²²⁰, but whether this reflects changes in spirometry or a primary effect on the muscle is not clear. All studies that have shown efficacy of theophylline in COPD were performed with slow-release preparations.

Theophylline is less effective and less well tolerated than inhaled long-acting bronchodilators²²⁵ and is not recommended if those drugs are available and affordable. However, there is evidence for a modest bronchodilator effect compared with placebo in stable COPD²²⁶ (**Evidence A**). There is also some evidence of symptomatic benefit compared to placebo²²⁷. Addition of theophylline to salmeterol produced a greater improvement in FEV₁ and breathlessness than salmeterol alone²²⁸ (**Evidence B**). Low-dose theophylline reduces exacerbations but does not improve post-bronchodilator lung function²²⁷ (**Evidence B**).

Adverse effects. Toxicity is dose-related, a particular problem with the xanthine derivatives because their therapeutic ratio is small and most of the benefit occurs only when near-toxic doses are given^{221,223,229}.

Methylxanthines are nonspecific inhibitors of all phosphodiesterase enzyme subsets, which explains their wide range of toxic effects. Problems include the development of atrial and ventricular arrhythmias (which can prove fatal) and grand mal convulsions (which can occur irrespective of prior epileptic history). Other side effects include headaches, insomnia, nausea, and heartburn, and these may occur within the therapeutic range of serum theophylline. These medications also have significant interactions with commonly used medications such as digitalis, coumadin, etc. Unlike the other bronchodilator classes, xanthine derivatives may involve a risk of overdose (either intentional or accidental).

Combination Bronchodilator Therapy. Combining bronchodilators with different mechanisms and durations of action may increase the degree of bronchodilation for equivalent or lesser side effects²³⁰. For example, a combination of a short-acting beta₂-agonist and an anticholinergic produces greater and more sustained improvements in FEV₁ than either drug alone and does not produce evidence of tachyphylaxis over 90 days of treatment^{191,231,232}. The combination of a beta₂-agonist, an anticholinergic, and/or theophylline may produce additional improvements in lung function^{191,224,228,231-235} and health status^{191,236}. Short-term combination therapy using formoterol and tiotropium has been shown to have a bigger impact on FEV₁ than the single components^{237,238} (**Evidence B**). Combinations of short-acting beta₂-agonists and anticholinergics are also superior compared to either medication alone in improving FEV₁ and symptoms²³¹ (**Evidence B**). Combinations of a long-acting beta₂-agonist and a long-acting anticholinergic have shown a significant increase in lung function whereas the impact on patient reported outcomes is still limited^{560,589}. There is still too little evidence to determine if a combination of long-acting bronchodilators is more effective than a long-acting anticholinergic alone for preventing exacerbations⁵⁶¹.

Corticosteroids

Inhaled Corticosteroids. The dose-response relationships and long-term safety of inhaled corticosteroids in COPD are not known. Only moderate to high doses have been used in long-term clinical trials. The efficacy and side effects of inhaled corticosteroids in asthma are dependent on the dose and type of corticosteroid²³⁹, but whether this is also the case in COPD is unclear. The effects of corticosteroids on pulmonary and systemic inflammation in patients with COPD are controversial, and their role in the management of stable COPD is limited to specific indications.

Regular treatment with inhaled corticosteroids improves symptoms, lung function, and quality of life, and reduces the frequency of exacerbations¹⁴⁴ in COPD patients with an FEV₁ < 60% predicted^{195,240-244} (**Evidence A**). Withdrawal

from treatment with inhaled corticosteroids may lead to exacerbations in some patients²⁴⁵, although in another study with severe and very severe COPD patients, inhaled corticosteroids could be gradually withdrawn over a three-month period without increasing the medium term risk of exacerbations, although lung function deteriorated significantly⁵⁹⁰. Withdrawal of inhaled corticosteroids, in COPD patients at low risk of exacerbation, can be safe provided that patients are left on maintenance treatment with long-acting bronchodilators⁶¹⁸. Regular treatment with inhaled corticosteroids does not modify the long-term decline of FEV₁ nor mortality in patients with COPD^{126,175,176,195,246,247,520} (**Evidence A**).

Adverse effects. Inhaled corticosteroid use is associated with higher prevalence of oral candidiasis, hoarse voice, and skin bruising^{126,175}. Treatment with inhaled corticosteroids is associated with an increased risk of pneumonia^{195,246-248,591}. While long-term treatment with triamcinolone acetonide is associated with an increased risk of reduced bone density, the evidence with other inhaled corticosteroids is controversial. One long-term study showed no effect of budesonide on bone density and fracture rate^{175,249}, and treatment over a three-year period with 500 mcg bid fluticasone propionate alone or in combination with salmeterol was not associated with decreased bone mineral density in a population of COPD patients with high prevalence of osteoporosis²⁵⁰.

Combination Inhaled Corticosteroid/Bronchodilator Therapy. An inhaled corticosteroid combined with a long-acting beta₂-agonist is more effective than the individual components in improving lung function and health status and reducing exacerbations in patients with moderate (**Evidence B**) to very severe COPD^{195,240,243,244,246,251-253,521,522} (**Evidence A**). An inhaled corticosteroid/long-acting beta₂-agonist combination given once daily does not show relevant differences regarding efficacy compared to twice daily⁶¹⁹. A large prospective clinical trial failed to demonstrate a statistically significant effect of combination therapy on mortality¹⁹⁵, but a subsequent meta-analysis found that combination therapy may reduce mortality with a number needed to treat (NNT) of 36²⁵⁴ (**Evidence B**). Combination therapy is associated with an increased risk of pneumonia²⁵⁵, but no other significant side effect (**Evidence A**). The addition of a long-acting beta₂-agonist/inhaled corticosteroid combination to tiotropium improves lung function and quality of life^{256,257} and may further reduce exacerbations (**Evidence B**) but more studies of triple therapy are needed²⁵⁸.

Oral Corticosteroids. Oral corticosteroids have numerous side effects⁵⁹². An important side effect of long-term treatment of COPD with systemic corticosteroids is steroid myopathy²⁵⁹⁻²⁶¹, which contributes to muscle weakness, decreased functionality, and respiratory failure in subjects with very severe COPD. In

view of the well-known toxicity of long-term treatment with oral corticosteroids, prospective studies on the long-term effects of these drugs in COPD are limited^{262,263}. However, systemic corticosteroids for treating acute exacerbations have been shown to improve symptoms, lung function, reduce rate of treatment failure, and shorten length of hospital stay. The effect of preventing a subsequent exacerbation has been shown in a pooled data analysis and it was demonstrated that systemic corticosteroids when being used to treat acute exacerbations can reduce 30-day readmission rates due to recurrent exacerbations⁶²⁰.

Phosphodiesterase-4 Inhibitors

The principal action of phosphodiesterase-4 inhibitors is to reduce inflammation by inhibiting of the breakdown of intracellular cyclic AMP^{264,265}. It is a once daily oral medication with no direct bronchodilator activity, although it has been shown to improve FEV₁ in patients treated with salmeterol or tiotropium²⁶⁴. Roflumilast reduces moderate and severe exacerbations treated with corticosteroids by 15-20% in patients with chronic bronchitis, severe to very severe COPD and a history of exacerbations²⁶⁶ (**Evidence A**). The effects on lung function are also seen when roflumilast is added to long-acting bronchodilators^{266,621} (**Evidence A**). There are no direct comparison or add-on studies of roflumilast and inhaled corticosteroids. Phosphodiesterase-4 inhibitors should always be used in combination with at least one long-acting bronchodilator.

Adverse effects. Phosphodiesterase-4 inhibitors have more adverse effects than inhaled medications for COPD²⁶⁴⁻²⁶⁶. The most frequent adverse effects are nausea, reduced appetite, abdominal pain, diarrhea, sleep disturbances, and headache. Adverse effects led to increased withdrawal in clinical trials from the group receiving roflumilast. Adverse effects seem to occur early during treatment, are reversible, and diminish over time with continued treatment. In controlled studies an average unexplained weight loss of 2 kg has been seen and weight monitoring during treatment is advised as well as avoiding treatment with roflumilast in underweight patients. Roflumilast should also be used with caution in patients with depression. Roflumilast and theophylline should not be given together.

Other Pharmacologic Treatments

Vaccines. Influenza vaccination can reduce serious illness (such as lower respiratory tract infections requiring hospitalization²⁶⁷) and death in COPD patients²⁶⁸⁻²⁷⁰ (**Evidence A**). Vaccines containing killed or live, inactivated viruses are recommended²⁷¹ as they are more effective in elderly patients with COPD²⁷². The strains are adjusted each year for appropriate effectiveness and should be given once each year²⁷³. Pneumococcal polysaccharide vaccine is recommended for COPD patients 65 years and older, and also in younger patients with significant comorbid conditions such as cardiac

disease²⁷⁴⁻²⁷⁶. In addition, this vaccine has been shown to reduce the incidence of community-acquired pneumonia in COPD patients younger than age 65 with an FEV₁ < 40% predicted²⁷⁷ (**Evidence B**).

Alpha-1 Antitrypsin Augmentation Therapy. Young patients with severe hereditary alpha-1 antitrypsin deficiency and established emphysema may be candidates for alpha-1 antitrypsin augmentation therapy (**Evidence C**). However, this therapy is very expensive, is not available in most countries, and is not recommended for patients with COPD that is unrelated to alpha-1 antitrypsin deficiency.

Antibiotics. In older studies prophylactic, continuous use of antibiotics was shown to have no effect on the frequency of exacerbations in COPD²⁷⁸⁻²⁸⁰, and a study that examined the efficacy of chemoprophylaxis undertaken in winter months over a period of 5 years concluded that there was no benefit²⁸¹. Although studies have shown some effects of antibiotics on exacerbation rate^{282,283}, the role of this treatment is unclear. A trial of daily azithromycin showed efficacy on exacerbation end-points with little evidence of treatment effect among current smokers⁵⁹³; however, treatment is not recommended because of an unfavorable balance between benefits and side effects²⁸⁴. Thus, the use of antibiotics, other than for treating infectious exacerbations of COPD and other bacterial infections, is currently not indicated^{285,286} (**Evidence B**).

Mucolytic (mucokinetic, mucoregulator) and Antioxidant Agents (ambroxol, erdosteine, carbocysteine, iodinated glycerol, N-acetylcysteine). The regular use of mucolytics in COPD has been evaluated in a number of long-term studies with controversial results²⁸⁷⁻²⁸⁹. Although a few patients with viscous sputum may benefit from mucolytics^{290,291}, the overall benefits seem to be very small; the widespread use of these agents cannot be recommended at present (**Evidence D**). Drugs like N-acetylcysteine may have antioxidant effects, leading to speculation that these medications could have a role in the treatment of patients with recurrent exacerbations^{292-295,562} (**Evidence B**). In patients treated with and without inhaled corticosteroids, high doses of N-acetylcysteine significantly reduced exacerbation rates, but only in GOLD stage 2 patients⁵⁹⁴. There is some evidence that in COPD patients not receiving inhaled corticosteroids, treatment with mucolytics such as carbocysteine and N-acetylcysteine may reduce exacerbations^{296,297,562} (**Evidence B**) although a Cochrane review showed little or no effect on the overall quality of life⁵²³.

Immunoregulators (immunostimulators, immunomodulators). Studies using an immunoregulator in COPD report a decrease in the severity and frequency of exacerbations^{298,299}. However, additional studies to examine the long-term effects of this therapy are required; at present, its regular use cannot be recommended³⁰⁰.

Antitussives. Cough, although sometimes a troublesome symptom in COPD, has a significant protective role³⁰¹. The regular use of antitussives is not recommended in stable COPD (**Evidence D**).

Vasodilators. The belief that pulmonary hypertension in COPD is associated with a poorer prognosis has provoked many attempts to reduce right ventricular afterload, increase cardiac output, and improve oxygen delivery and tissue oxygenation. Many agents have been evaluated, including inhaled nitric oxide, but the results have been uniformly disappointing. In patients with COPD, in whom hypoxemia is caused primarily by ventilation-perfusion mismatching rather than by increased intrapulmonary shunt (as in noncardiogenic pulmonary edema), inhaled nitric oxide can worsen gas exchange because of altered hypoxic regulation of ventilation-perfusion balance^{302,303}. Therefore, based on the available evidence, nitric oxide is contraindicated in stable COPD. Likewise, guidelines on the treatment of pulmonary hypertension do not recommend the use of endothelium-modulating agents for the treatment of pulmonary hypertension associated with COPD until data on their safety and efficacy in this condition are available³⁰⁴.

Narcotics (morphine). Oral and parenteral opioids are effective for treating dyspnea in COPD patients with very severe disease. There is insufficient data to conclude whether nebulized opioids are effective³⁰⁵. However, some clinical studies suggest that morphine used to control dyspnea may have serious adverse effects and its benefits may be limited to a few sensitive subjects³⁰⁶⁻³¹⁰.

Others. Nedocromil and leukotriene modifiers have not been adequately tested in COPD patients and cannot be recommended. There was no evidence of benefit—and some some evidence of harm (malignancy and pneumonia)—from an anti-TNF-alpha antibody (infliximab) tested in moderate to severe COPD³¹¹. There is no evidence for the effectiveness of herbal medicines in treating COPD³¹² and other alternative healing methods (e.g., acupuncture and homeopathy) have not been adequately tested. There is evidence that sildenafil does not improve the results of rehabilitation in patients with COPD and moderately increased pulmonary artery pressure⁵⁹⁵. In unselected patients there is no evidence that supplementation of vitamin D has a positive impact on exacerbations⁵⁹⁶.

NON-PHARMACOLOGIC THERAPIES

Rehabilitation

The principal goals of pulmonary rehabilitation are to reduce symptoms, improve quality of life, and increase physical and emotional participation in everyday activities^{313,314}. To accomplish these goals, pulmonary rehabilitation covers a range of non-pulmonary problems that may not be

Table 3.5. Benefits of Pulmonary Rehabilitation in COPD

- Improves exercise capacity (**Evidence A**).
- Reduces the perceived intensity of breathlessness (**Evidence A**).
- Improves health-related quality of life (**Evidence A**).
- Reduces the number of hospitalizations and days in the hospital (**Evidence A**).
- Reduces anxiety and depression associated with COPD (**Evidence A**).
- Strength and endurance training of the upper limbs improves arm function (**Evidence B**).
- Benefits extend well beyond the immediate period of training (**Evidence B**).
- Improves survival (**Evidence B**).
- Respiratory muscle training can be beneficial, especially when combined with general exercise training (**Evidence C**).
- Improves recovery after hospitalization for an exacerbation⁵²⁴ (**Evidence A**).
- Enhances the effect of long-acting bronchodilators (**Evidence B**).

adequately addressed by medical therapy for COPD, including exercise de-conditioning, relative social isolation, altered mood states (especially depression), muscle wasting, and weight loss. Pulmonary rehabilitation has been carefully evaluated in a large number of clinical trials and shown to increase peak workload, peak oxygen consumption, and endurance time³¹⁵. Benefits have been reported from rehabilitation programs conducted in inpatient, outpatient, and home settings^{315,316}; considerations of cost and availability most often determine the choice of setting. The various benefits of pulmonary rehabilitation are summarized in **Table 3.5**^{313,314,317-320}. However the increased exercise capacity may not necessarily translate into increased daily physical activity⁵⁶³.

The minimum length of an effective rehabilitation program is 6 weeks; the longer the program continues, the more effective the results³²¹⁻³²³. However, as yet, no effective program has been developed to maintain the effects over time³²⁴. Many physicians advise patients unable to participate in a structured program to exercise on their own (e.g., walking 20 minutes daily). The benefits of this general advice have not been tested, but because observational studies have indicated significant benefits of physical activity^{325,326}, and because physical activity is good for so many other reasons, it is highly reasonable to offer such advice to patients if a formal program is not available.

Components of Pulmonary Rehabilitation Programs

The components of pulmonary rehabilitation vary widely but a comprehensive programs includes exercise training, smoking cessation, nutrition counseling, and education.

Exercise training. Exercise tolerance can be assessed by either bicycle ergometry or treadmill exercise with the

measurement of a number of physiological variables, including maximum oxygen consumption, maximum heart rate, and maximum work performed. A less complex approach is to use a self-paced, timed walking test (e.g., 6-minute walking distance). These tests require at least one practice session before data can be interpreted. Shuttle walking tests offer a compromise: they provide more complete information than an entirely self-paced test, but are simpler to perform than a treadmill test¹⁵³.

Exercise training ranges in frequency from daily to weekly, in duration from 10 minutes to 45 minutes per session, and in intensity from 50% peak oxygen consumption (VO_2 max) to maximum tolerated²⁵³. The optimum length for an exercise program has not been investigated in randomized controlled trials but most studies involving fewer than 28 exercise sessions show inferior results compared to those with longer treatment periods³²⁷. In practice, the length depends on the resources available and usually ranges from 4 to 10 weeks, with longer programs resulting in larger effects than shorter programs³¹⁸.

In many programs, especially those using simple corridor exercise training, the patient is encouraged to walk to a symptom-limited maximum, rest, and then continue walking until 20 minutes of exercise have been completed. Where possible, endurance exercise training to 60-80% of the symptom-limited maximum is preferred. Endurance training can be accomplished through continuous or interval exercise programs. The latter involve the patient doing the same total work but divided into briefer periods of high-intensity exercise, which is useful when performance is limited by other comorbidities^{328,329}. Use of a simple wheeled walking aid seems to improve walking distance and reduces breathlessness in severely disabled COPD patients³³⁰⁻³³². Other approaches to improving outcomes such as use of oxygen during exercise³³³, exercising while breathing heliox gas mixtures³³⁴, or unloading the ventilator muscles while exercising remain experimental at present.

Some programs also include upper limb exercises, usually involving an upper limb ergometer or resistive training with weights. There are no randomized clinical trial data to support the routine inclusion of these exercises, but they may be helpful in patients with comorbidities that restrict other forms of exercise and those with evidence of respiratory muscle weakness³³⁵. In contrast, inspiratory muscle training appears to provide additional benefits when included in a comprehensive pulmonary rehabilitation program³³⁶⁻³³⁸. The addition of upper limb exercises or other strength training to aerobic training is effective in improving strength, but does not improve quality of life or exercise tolerance³³⁹.

The following points summarize current knowledge of

considerations important in choosing patients for pulmonary rehabilitation:

Functional status: Benefits have been seen in patients with a wide range of disability, although those who are chair-bound appear less likely to respond even to home visiting programs³⁴⁰ (**Evidence B**).

Severity of dyspnea: Stratification by breathlessness intensity using the mMRC questionnaire may be helpful in selecting patients most likely to benefit from rehabilitation. Those with mMRC grade 4 dyspnea may not benefit³⁴⁰ (**Evidence B**).

Motivation: Selecting highly motivated participants is especially important in the case of outpatient programs³⁴¹. Depressed mood is a risk factor for non-completion of rehabilitation programs⁶²².

Smoking status: There is no evidence that smokers will benefit less than nonsmokers, although some suggest that continuing smokers are less likely to complete pulmonary rehabilitation programs than nonsmokers³⁴¹ (**Evidence B**).

Education. Most pulmonary rehabilitation programs include an educational component. The topics that seem most appropriate for an education program include: smoking cessation; basic information about COPD; general approach to therapy and specific aspects of medical treatment; self-management skills; strategies to help minimize dyspnea; advice about when to seek help; decision-making during exacerbations; and advance directives and end-of-life issues.

The intensity and content of these educational messages should vary depending on the severity of the patient's disease, although the specific contributions of education to the improvements seen after pulmonary rehabilitation remain unclear. Studies indicate that patient education alone does not improve exercise performance or lung function³⁴²⁻³⁴⁵, but it can play a role in improving skills, ability to cope with illness, and health status³⁴⁶. These outcomes are not traditionally measured in clinical trials, but they may be most important in COPD where even pharmacologic interventions generally confer only a small benefit in terms of lung function.

Patients with severe COPD often express the desire to discuss end-of-life care with clinicians, but these conversations rarely occur in clinical practice. Simple, structured approaches to facilitate these conversations may help to improve the occurrence and quality of communication from the patients' perspective⁵²⁵. In particular, patients with a chronic life-limiting illness like COPD should be informed that, should they become critically ill, they or their family members may be in a position where they would need to decide whether a) a course of intensive care is likely to achieve their

personal goals of care, and b) they are willing to accept the burdens of such treatment. Communication about end-of-life care and advance care planning gives patients the opportunity to make informed decisions about the kind of care they want and ensure that their family and clinicians understand their values, goals, and perspectives⁵²⁶. Clinicians should develop and implement methods to help patients and their families to make informed choices that are consistent with patients' values. Such methods have the potential to improve the quality of care and simultaneously may contribute to efforts to reduce health care costs by ensuring patients receive care consistent with their goals and values^{527,528}.

Assessment and Follow-up. Baseline and outcome assessments of each participant in a pulmonary rehabilitation program should be made to quantify individual gains and target areas for improvement. Assessments should include:

- Detailed history and physical examination
- Measurement of post-bronchodilator spirometry
- Assessment of exercise capacity
- Measurement of health status and impact of breathlessness (e.g., CAT and mMRC scales)
- Assessment of inspiratory and expiratory muscle strength and lower limb strength (e.g., quadriceps) in patients who suffer from muscle wasting

The first two assessments are important for establishing entry suitability and baseline status but are not used in outcome assessment. The last three assessments are baseline and outcome measures. Several detailed questionnaires for assessing health status are available, including some that are specifically designed for patients with respiratory disease (e.g., *Chronic Respiratory Disease Questionnaire*²³⁶, *St. George Respiratory Questionnaire*³⁴⁷, *Chronic Obstructive Pulmonary Disease Assessment Test*¹²⁴), and there is increasing evidence that these questionnaires may be useful in a clinical setting. Health status can also be assessed by generic questionnaires, such as the *Medical Outcomes Study Short Form (SF36)*³⁴⁸, to enable comparison of quality of life in different diseases. The *Hospital Anxiety and Depression Scale (HADS)*³⁴⁹ and the *Primary Care Evaluation of Mental Disorders (PRIME-MD) Patient Questionnaire*³⁵⁰ have been used to improve identification and treatment of anxious and depressed patients.

Nutritional support. Low-to-moderate quality evidence suggests that nutritional support promotes significant gain in weight and fat-free mass among patients with COPD, especially if malnourished. In addition, significantly greater changes from baseline have been observed in supplemented patients for six-minute walk test, respiratory muscle strength and (only in malnourished patients) overall

HRQoL as measured by SGRQ. Positive effects have been observed when nutritional supplementation is proposed alone or as an adjunct to exercise training. The optimal amount and duration of supplementation are not clearly established⁵⁶⁴.

OTHER TREATMENTS

Oxygen Therapy

The long-term administration of oxygen (> 15 hours per day) to patients with chronic respiratory failure has been shown to increase survival in patients with severe resting hypoxemia³⁵⁹ (**Evidence B**). Long-term oxygen therapy is indicated for patients who have:

- PaO₂ at or below 7.3 kPa (55 mmHg) or SaO₂ at or below 88%, with or without hypercapnia confirmed twice over a three week period (**Evidence B**); or
- PaO₂ between 7.3 kPa (55 mmHg) and 8.0 kPa (60 mmHg), or SaO₂ of 88%, if there is evidence of pulmonary hypertension, peripheral edema suggesting congestive cardiac failure, or polycythemia (hematocrit > 55%) (**Evidence D**).

A decision about the use of long-term oxygen should be based on the resting PaO₂ or saturation values repeated twice over three weeks in the stable patient. Current data do not support the use of ambulatory oxygen in patient populations that do not meet the above criteria³⁶⁰.

Although air travel is safe for most patients with chronic respiratory failure who are on long-term oxygen therapy⁵²⁹, patients should ideally be able to maintain an in-flight PaO₂ of at least 6.7 kPa (50 mmHg). Studies indicate that this can be achieved in those with moderate to severe hypoxemia at sea level by supplementary oxygen at 3 L/min by nasal cannulae or 31% by Venturi facemask³⁶¹. Those with a resting PaO₂ at sea level > 9.3 kPa (70 mmHg) are likely to be safe to fly without supplementary oxygen^{362,363}, although it is important to emphasize that a resting PaO₂ > 9.3 kPa (70 mmHg) at sea level does not exclude the development of severe hypoxemia when travelling by air (**Evidence C**). Careful consideration should be given to any comorbidity that may impair oxygen delivery to tissues (e.g., cardiac impairment, anemia). Also, walking along the aisle may profoundly aggravate hypoxemia³⁶⁴.

Ventilatory Support

Non-invasive ventilation (NIV) is increasingly used in patients with stable very severe COPD. Randomized controlled trials provide contradictory results regarding the clinical benefits of long-term NIV in patients with COPD and chronic hypercapnia, especially in terms of health status

and survival⁵⁹⁷⁻⁵⁹⁹. Thus, there is insufficient evidence to formulate recommendations. The combination of NIV with long-term oxygen therapy may be of some use in a selected subset of patients, particularly in those with pronounced daytime hypercapnia³⁶⁵. It may improve survival but does not improve quality of life³⁶⁵. However, in patients with both COPD and obstructive sleep apnea there are clear benefits from continuous positive airway pressure (CPAP) in both survival and risk of hospital admission³⁶⁶.

Surgical Treatments

Lung Volume Reduction Surgery (LVRS). LVRS is a surgical procedure in which parts of the lung are resected to reduce hyperinflation³⁶⁷, making respiratory muscles more effective pressure generators by improving their mechanical efficiency (as measured by length/tension relationship, curvature of the diaphragm, and area of apposition)^{368,369}. In addition, LVRS increases the elastic recoil pressure of the lung and thus improves expiratory flow rates and reduces exacerbations³⁷⁰. The advantage of surgery over medical therapy is more significant among patients with predominantly upper-lobe emphysema and low exercise capacity prior to treatment. A prospective economic analysis indicated that LVRS is costly relative to health-care programs not including surgery³⁷¹. In contrast to medical treatment, LVRS has been demonstrated to result in improved survival (54% vs. 39.7%) in severe emphysema patients with upper-lobe emphysema and low post-rehabilitation exercise capacity³⁷² (**Evidence A**). In similar patients with high post-pulmonary rehabilitation exercise capacity no difference in survival was noted after LVRS, although health-related quality of life and exercise capacity improved. LVRS has been demonstrated to result in higher mortality than medical management in severe emphysema patients with an FEV₁ ≤ 20% predicted and either homogeneous emphysema on high resolution computed tomography or a DL_{CO} ≤ 20% predicted³⁷³.

Bronchoscopic Lung Volume Reduction (BLVR). In a post-hoc analysis, BLVR in COPD patients with severe airflow limitation (FEV₁ 15-45% predicted), heterogeneous emphysema on CT scan, and hyperinflation (TLC > 100% and RV > 150% predicted) has been demonstrated to result in modest improvements in lung function, exercise tolerance, and symptoms at the cost of more frequent exacerbations of COPD, pneumonia, and hemoptysis after implantation³⁷⁴. Additional data are required to define the optimal technique and patient population. Several non surgical bronchoscopic lung volume reduction techniques (e.g., valves, glues, coils) are being studied. However, available evidence is insufficient to determine their benefit-risk ratios, cost-effectiveness and possible roles in the strategy of care for patients with predominant emphysema. These techniques should not be used outside clinical trials until more data are available.

Lung Transplantation. In appropriately selected patients with very severe COPD, lung transplantation has been shown to improve quality of life and functional capacity^{375,376}. The common complications seen in COPD patients after lung transplantation, apart from post-operative mortality, are acute rejection, bronchiolitis obliterans, opportunistic infections such as CMV, fungal (*Candida*, *Aspergillus*, *Cryptococcus*, *Pneumocystis*) or bacterial (*Pseudomonas*, *Staphylococcus species*) infections, and lymphoproliferative disease³⁷⁷. Lung transplantation is limited by the shortage of donor organs and costs. Criteria for referral for lung transplantation include COPD with a BODE index exceeding 5. Recommended criteria for listing include a BODE index of 7-10 and at least one of the following: history of exacerbation associated with acute hypercapnia [$\text{PaCO}_2 > 6.7$ kPa (50 mmHg)]; pulmonary hypertension, cor pulmonale, or both despite oxygen therapy; and $\text{FEV}_1 < 20\%$ predicted with either $\text{DL}_{\text{CO}} < 20\%$ predicted or homogenous distribution of emphysema³⁷⁸ (**Evidence C**).

Integrated Care Programs. COPD is a complex disease that requires the input of multiple care providers who need to work together closely. In principle, use of a formal structured program that determines how each component is delivered should make care more efficient and effective, but the evidence for this is divided. A meta-analysis of small trials concluded that an integrated care program improved a number of clinical outcomes, although not mortality⁶⁰⁰. In contrast, a large multi-center study within an existing well-organized system of care did not confirm this⁶⁰¹. The pragmatic conclusion is that well organized care is important, but there may be no advantage in structuring it tightly into a formalized program.

Bullectomy. Bullectomy is an older surgical procedure for bullous emphysema. Removal of a large bulla that does not contribute to gas exchange decompresses the adjacent lung parenchyma. Pulmonary hypertension, hypercapnia, and severe emphysema are not absolute contraindications for bullectomy.

Palliative Care, End-of-life Care, and Hospice Care.

The disease trajectory in COPD is usually marked by a gradual decline in health status and increasing symptoms, punctuated by acute exacerbations that are associated with an increased risk of dying⁵³⁰. Although mortality following hospitalization for an acute exacerbation of COPD is falling⁵³¹, it still varies between 23%⁵³² and 80%⁵³³. Progressive respiratory failure, cardiovascular diseases, malignancies and other diseases are the primary cause of death in patients with COPD hospitalized for an exacerbation⁵³³. For all these reasons, palliative care, end-of-life care, and hospice care are important components of the care of patients with advanced COPD.

Palliative care is the broadest term and incorporates (but is not limited to) both end-of-life care (care for those who are actively dying) as well as hospice care (a model for delivery of end-of-life care for patients who are terminally ill and predicted to have less than 6 months to live). The goal of palliative care is to prevent and relieve suffering, and to support the best possible quality of life for patients and their families, regardless of the stage of disease or the need for other therapies⁵³⁴. Therefore, palliative care is an important component in the management of all patients with advanced COPD and should begin at the time of the diagnosis of a chronic life-limiting illness such as COPD; yet patients with COPD are less likely to receive such services than patients with lung cancer^{535,536}. Palliative care expands traditional disease-model medical treatment to increase the focus on the goals of enhancing quality of life, optimizing function, helping with decision making about end-of-life care, providing emotional and spiritual support to patients and their families⁵³⁴. Increasingly, palliative care teams are available for consultation for hospitalized patients and such teams are rapidly increasing in numbers and capacity⁵³⁷. Availability for outpatient palliative care consultation is less common, but has been shown to improve quality of life, reduce symptoms and even prolong survival for some patients, such as those with advanced lung cancer⁵³⁶. Clinicians caring for patients with COPD should help identify patients who could benefit from palliative care services and identify available palliative care resources within their community for these patients.

For patients with the most advanced and terminal illness, hospice services may provide additional benefit. Hospice services often focus on patients with severe disability or symptom burden and may provide these services within the patient's home or in hospice beds in dedicated hospice units or other institutions such as hospitals or nursing homes. The National Hospice and Palliative Care Organization (<http://www.nhpco.org>) provides guidance for selecting patients with non-cancer diseases like COPD for access to hospice services (for example, disabling dyspnea at rest that is poorly responsive to bronchodilators and progression of advanced disease demonstrated by increasing hospitalizations or emergency department visits)^{535,536}. These guidelines discuss the difficulties in accurately predicting the prognosis of patients with advanced COPD, but recognize the appropriateness of providing hospice services for some of these patients⁵³⁴.

CHAPTER

4

MANAGEMENT OF STABLE COPD

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

CHAPTER 4: MANAGEMENT OF STABLE COPD

KEY POINTS:

- Identification and reduction of exposure to risk factors are important steps in the prevention and treatment of COPD. All individuals who smoke should be encouraged to quit.
- The level of FEV₁ is an inadequate descriptor of the impact of the disease on patients and for this reason individualized assessment of symptoms and future risk of exacerbation should also be incorporated into the management strategy for stable COPD.
- Pharmacologic therapy is used to reduce symptoms, reduce frequency and severity of exacerbations, and improve health status and exercise tolerance. Existing medications for COPD have not been conclusively shown to modify the long-term decline in lung function that is the hallmark of this disease.
- For both beta₂-agonists and anticholinergics, long-acting formulations are preferred over short-acting formulations. Based on efficacy and side effects, inhaled bronchodilators are preferred over oral bronchodilators.
- Long-term treatment with inhaled corticosteroids added to long-acting bronchodilators is recommended for patients at high risk of exacerbations.
- Long-term monotherapy with oral or inhaled corticosteroids is not recommended in COPD.
- The phosphodiesterase-4 inhibitor roflumilast may be useful to reduce exacerbations for patients with FEV₁ < 50% predicted, chronic bronchitis, and frequent exacerbations.
- Influenza vaccines can reduce the risk of serious illness (such as hospitalization due to lower respiratory tract infections) and death in COPD patients.
- Currently, the use of antibiotics is not indicated in COPD, other than for treating infectious exacerbations of COPD and other bacterial infections.
- All COPD patients with breathlessness when walking at their own pace on level ground appear to benefit from rehabilitation and maintenance of physical activity, improving their exercise tolerance and quality of life, and reducing symptoms of dyspnea and fatigue.

INTRODUCTION

Once COPD has been diagnosed, effective management should be based on an individualized assessment of disease in order to reduce both current symptoms and future risks (Table 4.1). These goals should be reached with minimal side effects from treatment, a particular challenge in COPD patients because they commonly have comorbidities that also need to be carefully identified and treated.

Table 4.1. Goals for Treatment of Stable COPD

<ul style="list-style-type: none">• Relieve symptoms• Improve exercise tolerance• Improve health status	→	REDUCE SYMPTOMS
<i>and</i>		
<ul style="list-style-type: none">• Prevent disease progression• Prevent and treat exacerbations• Reduce mortality	→	REDUCE RISK

It is crucial for patients with COPD to understand the nature of their disease, the risk factors for its progression, and their role and that of their health care workers in achieving optimal management and health outcomes. The type of health care workers seen, and the frequency of visits, will depend on the health care system. Ongoing monitoring should ensure that the goals of treatment are being met and should include continuous evaluation of exposure to risk factors and monitoring of disease progression, the effect of treatment and possible adverse effects, exacerbation history, and comorbidities. In addition, patients should receive general advice on healthy living, including diet and the fact that physical exercise is safe and encouraged for people with COPD.

Identification and reduction of exposure to risk factors are important in the treatment and prevention of COPD. Since cigarette smoking is the most commonly encountered and easily identifiable risk factor, smoking cessation should be encouraged for all individuals who smoke. Reduction of total personal exposure to occupational dusts, fumes, and gases and to indoor and outdoor air pollutants may be more difficult but should be attempted.

IDENTIFY AND REDUCE EXPOSURE TO RISK FACTORS

Tobacco Smoke

Smoking cessation is the key intervention for all COPD patients who continue to smoke (**Evidence A**). Health care providers are important to the delivery of smoking cessation messages and interventions and should encourage all patients who smoke to quit, even when patients visit a health care provider for reasons unrelated to COPD or breathing problems.

Occupational Exposures


Although studies as yet have not been done to demonstrate whether interventions to reduce occupational exposures also reduce the burden of COPD, it seems common sense to advise patients to avoid continued exposures to potential aggravants, if possible (**Evidence D**).

Indoor and Outdoor Air Pollution

Reducing the risk from indoor and outdoor air pollution is feasible and requires a combination of public policy, local and national resources, cultural changes, and protective steps taken by individual patients. Reduction of exposure to smoke from biomass fuel, particularly among women and children, is a crucial goal to reduce the prevalence of COPD worldwide. Efficient ventilation, non-polluting cooking stoves, use of flues, and similar interventions are feasible and should be recommended^{379,380} (**Evidence B**).

Table 4.2. Model of Symptom/Risk of Evaluation of COPD

When assessing risk, choose the highest risk according to GOLD grade or exacerbation history. (One or more hospitalizations for COPD exacerbations should be considered high risk.)



Patient Category	Characteristics	Spirometric Classification	Exacerbations per year	CAT	mMRC
A	Low Risk, Less Symptoms	GOLD 1-2	≤ 1	< 10	0-1
B	Low Risk, More Symptoms	GOLD 1-2	≤ 1	≥ 10	≥ 2
C	High Risk, Less Symptoms	GOLD 3-4	≥ 2	< 10	0-1
D	High Risk, More Symptoms	GOLD 3-4	≥ 2	≥ 10	≥ 2

TREATMENT OF STABLE COPD

In previous versions of the GOLD report, COPD treatment recommendations were based on spirometry only. This is in keeping with the fact that most of the clinical trial evidence about treatment efficacy in COPD is oriented around baseline FEV₁. However, FEV₁ alone is a poor descriptor of disease status and for this reason the treatment strategy for stable COPD should also consider an individual patient's symptoms and future risk of exacerbations. This individualized assessment is summarized in **Table 4.2**.

Moving from Clinical Trials to Recommendations for Routine Practice – Considerations

The guidance for clinical practice presented below is based on evidence from clinical trials, as detailed in the discussion of Evidence Levels at the beginning of this document. However, it is important to recognize that all clinical trials recruit restricted groups of patients; this limits their generalizability. In COPD the key inclusion criteria are: baseline FEV₁, acute bronchodilator reversibility, smoking history, symptoms and a prior history of exacerbations. A few general considerations related to these inclusion criteria are discussed below.

Baseline FEV₁. The evidence for pharmacological treatment of COPD is mostly based on the severity of airflow limitation (FEV₁, % predicted), and GOLD spirometry classification has often been used as an entry criterion for clinical trials. There is almost no evidence on efficacy of COPD treatments in patients with FEV₁ > 70% predicted (GOLD 1), and no evidence at all concerning anti-inflammatory treatment in patients with FEV₁ > 60% predicted. Many studies of combination medications (inhaled corticosteroids plus long-acting beta₂-agonists) have been limited to GOLD 3-4 (Severe-Very Severe airflow limitation) patients. As no trials have been carried out purely in GOLD 2 patients, evidence of the efficacy of combination treatment in this group has to be drawn from studies that included such patients as a subset of participants. Large studies such as TORCH¹⁹⁵ and UPLIFT²¹⁴ each contained over 2,000 GOLD 2 patients, albeit in the lower stratum of GOLD 2 (FEV₁ < 60% predicted). In general, it is important to draw a distinction between absence of evidence that a treatment works and presence of evidence that a treatment does not work.

Acute Bronchodilator Reversibility. Many COPD trials have used low reversibility of airflow limitation as an entry criterion. Acute reversibility is not a reliable measurement³⁸¹ and, in general, acute reversibility in response to bronchodilator is a poor predictor of a treatment's benefit for FEV₁ after one year³⁸². Thus, this common clinical trial entry criterion has limited impact on the reliability of therapeutic recommendations.

Symptoms. Almost all studies have included patients with respiratory symptoms; there are no data on asymptomatic patients. No studies have reported results based upon stratified symptom levels.

Exacerbation Prevention. Studies in which exacerbations are a major outcome often “enrich” the patient population by requiring a history of frequent exacerbations in the preceding year, as it is often easier to demonstrate an effect of treatment preventing exacerbations if the exacerbations actually occur. However, large trials that have not used this entry criterion have also shown reductions in exacerbations, even in patients with less severe airflow limitation^{195,214}. The patient’s own history of exacerbations appears to be the most powerful predictor of future exacerbations¹³², so the GOLD panel assumed that it is safe to extrapolate evidence of efficacy from clinical trials to appropriate patients in routine practice, regardless of the trial’s entry criteria concerning previous exacerbation history.

Sub-Group Analysis. Results of clinical trials potentially apply to every member of the intention-to-treat population, whether they lie in the center of the distribution of severity or at the extremes. Sub-group analysis, whether pre-specified or not, must be used with caution. For example, if a treatment has no effect in the intention-to-treat population, but appears to have an effect that is confined to one sub-group, there is a strong likelihood that one of the other groups would be worse on the treatment. In contrast, subgroup analysis is useful if it shows that a treatment effect is consistent in size and direction across the range of patients recruited to the study. In summary, sub-group analysis does not provide robust evidence that a treatment works in a specific subgroup, but it can provide confidence that the results from the intention-to-treat population apply to patients who met the study entry criteria. Subgroup analysis can also generate hypotheses to be tested in subsequent trials.

NON-PHARMACOLOGIC TREATMENT

Non-pharmacologic management of COPD according to the individualized assessment of symptoms and exacerbation risk is shown in **Table 4.3**.

Smoking Cessation

Smoking cessation should be considered the most important intervention for all COPD patients who smoke regardless of the level of disease severity.

Physical Activity

Physical activity is recommended for all patients with COPD. There is very little COPD-specific evidence to support recommendations for physical activity other than studies of pulmonary rehabilitation (the physical exercise component is believed to provide the most benefit). However, given the overall population benefits of physical exercise and its role in primary and secondary prevention of cardiovascular disease, it seems intuitively correct to recommend daily physical activity.

Rehabilitation

Although more information is needed on criteria for patient selection for pulmonary rehabilitation programs, all COPD patients appear to benefit from rehabilitation and maintenance of physical activity, improving their exercise tolerance and experiencing decreased dyspnea and fatigue³⁸³ (**Evidence A**). Several studies have documented an effect of pulmonary rehabilitation in patients with breathlessness, usually mMRC > 1, and following acute exacerbations. Data suggest that these benefits can be sustained even after a single pulmonary rehabilitation program^{341,384,385}. Benefit does wane after a rehabilitation program ends, but if exercise training is maintained at home the patient’s health status remains above pre-rehabilitation levels (**Evidence B**).

Vaccination

Decisions about vaccination in COPD patients depend on local policies, availability, and affordability.

Table 4.3. Non-Pharmacologic Management of COPD

Patient Group	Essential	Recommended	Depending on Local Guidelines
A	Smoking cessation (can include pharmacologic treatment)	Physical activity	Flu vaccination Pneumococcal vaccination
B-D	Smoking cessation (can include pharmacologic treatment) Pulmonary rehabilitation	Physical activity	Flu vaccination Pneumococcal vaccination

PHARMACOLOGIC TREATMENT

Pharmacologic therapy in COPD is used to reduce symptoms, reduce the frequency and severity of exacerbations, and improve health status and exercise tolerance. Existing medications for COPD have not been conclusively shown to modify the long-term decline in lung function that is the hallmark of this disease^{126,159,175,176}.

The classes of medications commonly used in treating COPD are shown in **Table 3.3** and a detailed description of the effects of these medications is given in Chapter 3. The choice within each class depends on the availability of medication and the patient's response. A proposed model for initial pharmacological management of COPD according to the individualized assessment of symptoms and exacerbation risk is shown in **Table 4.4**.

Group A patients have few symptoms and a low risk of exacerbations. Specific evidence for the effectiveness of pharmacologic treatments is not available for patients with FEV₁ > 80% predicted (GOLD 1). However, for all Group A patients, a short-acting bronchodilator used as needed is recommended as first choice based on its effect on lung function and breathlessness⁵⁶⁵. An alternative choice is a combination of short-acting bronchodilators or the introduction of a long-acting bronchodilator. The evidence for this step-up is weak; few studies of the combination exist^{191,386}, and most trials of therapy with long-acting bronchodilators have been performed in patients with more severe airflow limitation^{212,387}.

Group B patients have more significant symptoms but still a low risk of exacerbations. Long-acting bronchodilators are superior to short-acting bronchodilators (taken as needed, or prn) and are therefore recommended^{212,387}. There is no evidence to recommend one class of long-acting bronchodilators over another for initial treatment. In the individual patient, the choice should depend on the patient's perception of symptom relief. For patients with severe breathlessness, the alternative choice is a combination of long-acting bronchodilators^{237,238}. Other possible treatments include short-acting bronchodilators and theophylline, the latter of which can be used if inhaled bronchodilators are unavailable or unaffordable.

Group C patients have few symptoms but a high risk of exacerbations. As first choice a fixed combination of inhaled corticosteroid/long-acting beta₂-agonist or a long-acting anticholinergic is recommended^{195,212,214,240,244,251,388}. Unfortunately, there is only one study directly comparing these treatments, which makes differentiation difficult³⁸⁹. As an alternative choice a combination of two long-acting bronchodilators or the combination of inhaled corticosteroid/

long-acting anticholinergic can be used. Both long-acting anticholinergic and long-acting beta₂-agonist reduce the risk of exacerbations^{212,387}, and although good long-term studies are lacking, this principle of combination treatment seems sound (although in many countries expensive). The recommendation for a combination of inhaled corticosteroid/long-acting anticholinergic is not evidence-based, but this lack of evidence seems to be the result of lack of interest from the pharmaceutical industry rather than doubts about the rationale. A phosphodiesterase-4 inhibitor used in combination with at least one long-acting bronchodilator could be considered if the patient has chronic bronchitis^{264,266}. Other possible treatments include short-acting bronchodilators and theophylline if long-acting inhaled bronchodilators are unavailable or unaffordable.

Group D patients have many symptoms and a high risk of exacerbations. The first choice of therapy is inhaled corticosteroid plus long-acting beta₂-agonist or long-acting anticholinergic, although there are conflicting findings concerning this treatment²⁵⁷; support for it mainly comes from short-term studies^{257,538,539} (**Evidence B**). As second choice a combination of all three classes of drugs (inhaled corticosteroids/long-acting beta₂-agonist/long-acting anticholinergic) is recommended²⁵⁶. It is also possible to add a phosphodiesterase-4 inhibitor to the treatment chosen as first choice, provided the patient has chronic bronchitis²⁶⁶. A phosphodiesterase-4 inhibitor is effective when added to a long-acting bronchodilator²⁶⁴, whereas evidence of its benefit when added to inhaled corticosteroid comes from less valid secondary analyses. Other possible treatments include short-acting bronchodilators, and theophylline or carbocysteine²⁹⁶ can be used if long-acting inhaled bronchodilators are unavailable or unaffordable.

Bronchodilators – Recommendations

- For both beta₂-agonists and anticholinergics, long-acting formulations are preferred over short-acting formulations (**Evidence A**).
- The combined use of short- or long-acting beta₂-agonists and anticholinergics may be considered if symptoms are not improved with single agents⁶⁰² (**Evidence B**).
- Based on efficacy and side effects inhaled bronchodilators are preferred over oral bronchodilators (**Evidence A**).
- Based on evidence of relatively low efficacy and more side effects, treatment with theophylline is not recommended unless other long-term treatment bronchodilators are unavailable or unaffordable (**Evidence B**).

Table 4.4. Initial Pharmacologic Management of COPD*

Patient Group	Recommended First Choice	Alternative Choice	Other Possible Treatments**
A	Short-acting anticholinergic prn or Short-acting beta ₂ -agonist prn	Long-acting anticholinergic or Long-acting beta ₂ -agonist or Short-acting beta ₂ -agonist and short-acting anticholinergic	Theophylline
B	Long-acting anticholinergic or Long-acting beta ₂ -agonist	Long-acting anticholinergic and long-acting beta ₂ -agonist	Short-acting beta ₂ -agonist and/or Short-acting anticholinergic Theophylline
C	Inhaled corticosteroid + long-acting beta ₂ -agonist or Long-acting anticholinergic	Long-acting anticholinergic and long-acting beta ₂ -agonist or Long-acting anticholinergic and phosphodiesterase-4 inhibitor or Long-acting beta ₂ -agonist and phosphodiesterase-4 inhibitor	Short-acting beta ₂ -agonist and/or Short-acting anticholinergic Theophylline
D	Inhaled corticosteroid + long-acting beta ₂ -agonist and/or Long-acting anticholinergic	Inhaled corticosteroid + long-acting beta ₂ -agonist and long-acting anticholinergic or Inhaled corticosteroid + long-acting beta ₂ -agonist and phosphodiesterase-4 inhibitor or Long-acting anticholinergic and long-acting beta ₂ -agonist or Long-acting anticholinergic and phosphodiesterase-4 inhibitor	Carbocysteine N-acetylcysteine Short-acting beta ₂ -agonist and/or Short-acting anticholinergic Theophylline

*Medications in each box are mentioned in alphabetical order, and therefore not necessarily in order of preference.

**Medications in this column can be used alone or in combination with other options in the Recommended First Choice and Alternative Choice columns.

Corticosteroids and Phosphodiesterase-4 Inhibitors — Recommendations

- There is no evidence to recommend a short-term therapeutic trial with oral corticosteroids in patients with COPD to identify those who will respond to inhaled corticosteroids or other medications.
- Long-term treatment with inhaled corticosteroids is recommended for patients with severe and very severe COPD and frequent exacerbations that are not adequately controlled by long-acting bronchodilators (**Evidence A**).
- Long-term monotherapy with oral corticosteroids is not recommended in COPD (**Evidence A**).
- Long-term monotherapy with inhaled corticosteroids is not recommended in COPD because it is less effective than the combination of inhaled corticosteroids with long-acting beta₂-agonists (**Evidence A**).
- Long-term treatment containing inhaled corticosteroids should not be prescribed outside their indications, due to the risk of pneumonia and the possibility of an increased risk of fractures following long-term exposure⁵⁴⁰.
- The phosphodiesterase-4 inhibitor, roflumilast, may also be used to reduce exacerbations for patients with chronic bronchitis, severe and very severe COPD, and frequent exacerbations that are not adequately controlled by long-acting bronchodilators (**Evidence B**).

MONITORING AND FOLLOW-UP

Routine follow-up is essential in COPD. Lung function can be expected to worsen over time, even with the best available care. Symptoms and objective measures of airflow limitation should be monitored to determine when to modify therapy and to identify any complications that may develop. As at the initial assessment, follow-up visits should include a discussion of symptoms, particularly any new or worsening symptoms, and a physical examination. Comprehensive self-management or routine monitoring does not appear to show long term benefits in terms of quality of life or self efficacy over usual care alone in COPD patients in general practice⁵⁶⁶.

Monitor Disease Progression and Development of Complications

Measurements. Decline in lung function is best tracked by spirometry performed at least once a year to identify patients whose lung function is declining quickly. Questionnaires such as the COPD Assessment Test (CAT)¹²⁴ can be performed every two to three months; trends and changes are more valuable than single measurements.

Symptoms. At each visit, inquire about changes in symptoms since the last visit, including cough and sputum,

breathlessness, fatigue, activity limitation, and sleep disturbances.

Smoking Status. At each visit, determine current smoking status and smoke exposure; strongly encourage participation in programs to reduce and eliminate wherever possible exposure to COPD risk factors.

Monitor Pharmacotherapy and Other Medical Treatment

In order to adjust therapy appropriately as the disease progresses, each follow-up visit should include a discussion of the current therapeutic regimen. Dosages of various medications, adherence to the regimen, inhaler technique, effectiveness of the current regime at controlling symptoms, and side effects of treatment should be monitored. Treatment modifications should be recommended as appropriate with a focus on avoiding unnecessary polypharmacy.

At the individual patient level, measurements such as FEV₁ and questionnaires such as the CAT are useful but are not completely reliable, because the size of a clinically important response is smaller than between-assessment variability. For this reason, the following questions might be useful when deciding whether a patient has had a symptomatic response to treatment:

- Have you noticed a difference since starting this treatment?
- If you are better:
 - Are you less breathless?
 - Can you do more?
 - Can you sleep better?
 - Describe what difference it has made to you.
- Is that change worthwhile to you?

Monitor Exacerbation History

Evaluate the frequency, severity, and likely causes of any exacerbations³⁹¹. Increased sputum volume, acutely worsening dyspnea, and the presence of purulent sputum should be noted. Specific inquiry into unscheduled visits to providers, telephone calls for assistance, and use of urgent or emergency care facilities is important. Severity of exacerbations can be estimated by the increased need for bronchodilator medication or corticosteroids and by the need for antibiotic treatment. Hospitalizations should be documented, including the facility, duration of stay, and any use of critical care or mechanical ventilatory support.

Monitor Comorbidities

Comorbidities are common in COPD, amplify the disability associated with COPD, and can potentially complicate its management. Until more integrated guidance about disease management for specific comorbid problems

becomes available, the focus should be on identification and management of these individual problems in line with local treatment guidance (See also Chapter 6).

Surgery in the COPD Patient

Postoperative pulmonary complications are as important and common as postoperative cardiac complications and, consequently, are a key component of the increased risk posed by surgery in COPD patients³⁹². The principal potential factors contributing to the risk include smoking, poor general health status, age, obesity, and COPD severity. A comprehensive definition of postoperative pulmonary complications should include only major pulmonary respiratory complications, namely lung infections, atelectasis and/or increased airflow limitation, which all potentially result in acute respiratory failure and aggravation of underlying COPD^{291,393-395}.

Increased risk of postoperative pulmonary complications in COPD patients may vary with the severity of COPD, although the surgical site is the most important predictor; risk increases as the incision approaches the diaphragm³⁹⁴. Most reports conclude that epidural or spinal anesthesia have a lower risk than general anesthesia, although the results are not totally uniform.

For lung resection, the individual patient's risk factors should be identified by careful history, physical examination, chest radiography, and pulmonary function tests. Although the value of pulmonary function tests remains contentious, there is consensus that all COPD candidates for lung resection should undergo a complete battery of tests, including spirometry with bronchodilator response, static lung volumes, diffusing capacity, and arterial blood gases at rest^{396,397}. COPD patients at high risk for surgical complications due to poor lung function should undergo further lung function assessment, for example, tests of regional distribution of perfusion and exercise capacity^{396,397}.

The risk of postoperative complications from lung resection appears to be increased in patients with decreased predicted postoperative pulmonary function (FEV_1 or $DL_{CO} < 30-40\%$ predicted) or exercise capacity (peak $VO_2 < 10$ ml/kg/min or 35% predicted). The final decision to pursue surgery should be made after discussion with the surgeon, pulmonary specialist, primary clinician, and the patient. To prevent postoperative pulmonary complications, stable COPD patients clinically symptomatic and/or with limited exercise capacity should be treated intensively before surgery, with all the measures already well established for stable COPD patients who are not about to have surgery. Surgery should be postponed if an exacerbation is present.

CHAPTER

5

***MANAGEMENT
OF
EXACERBATIONS***

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

CHAPTER 5: MANAGEMENT OF EXACERBATIONS

KEY POINTS:

- An exacerbation of COPD is an acute event characterized by a worsening of the patient's respiratory symptoms that is beyond normal day-to-day variations and leads to a change in medication.
- Exacerbations of COPD can be precipitated by several factors. The most common causes appear to be viral upper respiratory tract infections and infection of the tracheobronchial tree.
- The diagnosis of an exacerbation relies exclusively on the clinical presentation of the patient complaining of an acute change of symptoms (baseline dyspnea, cough, and/or sputum production) that is beyond normal day-to-day variation.
- The goal of treatment in COPD exacerbations is to minimize the impact of the current exacerbation and to prevent the development of subsequent exacerbations.
- Short-acting inhaled beta₂-agonists with or without short-acting anticholinergics are usually the preferred bronchodilators for treatment of an exacerbation.
- Systemic corticosteroids and antibiotics can shorten recovery time, improve lung function (FEV₁) and arterial hypoxemia (PaO₂), and reduce the risk of early relapse, treatment failure, and length of hospital stay.
- COPD exacerbations can often be prevented. Smoking cessation, influenza and pneumococcal vaccination, knowledge of current therapy including inhaler technique, and treatment with long-acting inhaled bronchodilators, with or without inhaled corticosteroids, and treatment with a phosphodiesterase-4 inhibitor are all interventions that reduce the number of exacerbations and hospitalizations.

DEFINITION

*An exacerbation of COPD is an acute event characterized by a worsening of the patient's respiratory symptoms that is beyond normal day-to-day variations and leads to a change in medication*¹²⁸⁻¹³⁰.

Exacerbations of COPD are important events in the course of the disease because they:

- Negatively affect a patient's quality of life^{144,391}
- Have effects on symptoms and lung function that take several weeks to recover from³⁹⁸
- Accelerate the rate of decline of lung function^{399,400}
- Are associated with significant mortality, particularly in those requiring hospitalization
- Have high socioeconomic costs⁴⁰¹

40 MANAGEMENT OF EXACERBATIONS

In-hospital mortality of patients admitted for a hypercapnic exacerbation with acidosis is approximately 10%⁴⁰². Mortality reaches 40% at 1 year after discharge in those needing mechanical support, and all-cause mortality 3 years after hospitalization is as high as 49%⁴⁰¹⁻⁴⁰⁵. Prevention, early detection, and prompt treatment of exacerbations are vital to reduce the burden of COPD⁴⁰⁶.

Exacerbations of COPD can be precipitated by several factors. The most common causes appear to be respiratory tract infections (viral or bacterial). Bronchoscopic studies have shown that at least 50% of patients have bacteria in their lower airways during exacerbations of COPD⁴⁰⁷⁻⁴⁰⁹, but a significant proportion of these patients also have bacteria colonizing their lower airways in the stable phase of the disease. On the other hand, there is some indication that the bacterial burden increases during some exacerbations of COPD⁴¹⁰⁻⁴¹², and that acquisition of bacterial strains that are new to the patient is associated with exacerbations of COPD⁴¹³. Peaks of air pollution can also precipitate exacerbations of COPD⁴¹⁴⁻⁴¹⁶ and increase hospitalizations and mortality⁶⁰³. However, the cause of about one-third of severe exacerbations of COPD cannot be identified. Some patients appear particularly prone to suffer exacerbations of COPD whereas others do not. Those reporting two or more exacerbations of COPD per year are often defined as "frequent exacerbators"¹³², a phenotype that appears stable over time.

In addition to infections and exposure to pollutants, exacerbations of respiratory symptoms (especially dyspnea) in patients with COPD may be due to different mechanisms that may overlap in the same patients. Conditions that may mimic and/or aggravate exacerbations, including pneumonia, pulmonary embolism, congestive heart failure, cardiac arrhythmia⁶²³, pneumothorax, and pleural effusion, need to be considered in the differential diagnosis and treated if present^{128,291,398,417}. Interruption of maintenance therapy has also been shown to lead to exacerbations.

DIAGNOSIS

Currently, the diagnosis of an exacerbation relies exclusively on the clinical presentation of the patient complaining of an acute change of symptoms (baseline dyspnea, cough, and/or sputum production) that is beyond normal day-to-day variation. In the future, a biomarker or panel of biomarkers that allows a more precise etiologic diagnosis would be desirable. However, clinically useful biomarkers for COPD patients in stable condition have yet to be identified.

ASSESSMENT

The assessment of an exacerbation is based on the patient's medical history and clinical signs of severity (Tables 5.1 and 5.2) and some laboratory tests, if available.

Table 5.1. Assessment of COPD Exacerbations: Medical History

- Severity of COPD based on degree of airflow limitation
- Duration of worsening or new symptoms
- Number of previous episodes (total/hospitalizations)
- Comorbidities
- Present treatment regimen
- Previous use of mechanical ventilation

Table 5.2. Assessment of COPD Exacerbations: Signs of Severity

- Use of accessory respiratory muscles
- Paradoxical chest wall movements
- Worsening or new onset central cyanosis
- Development of peripheral edema
- Hemodynamic instability
- Deteriorated mental status

The following tests may be considered to assess the severity of an exacerbation:

- *Pulse oximetry* is useful for tracking and/or adjusting supplemental oxygen therapy. The measurement of arterial blood gases is vital if the coexistence of acute or acute-on-chronic respiratory failure is suspected ($\text{PaO}_2 < 8.0 \text{ kPa}$ (60 mmHg) with or without $\text{PaCO}_2 > 6.7 \text{ kPa}$ (50 mmHg) breathing ambient air). Assessment of the acid-base status is necessary before initiating mechanical ventilation^{291,418}.
- *Chest radiographs* are useful in excluding alternative diagnoses.
- An *ECG* may aid in the diagnosis of coexisting cardiac problems.
- *Whole blood count* may identify polycythemia (hematocrit > 55%), anemia, or leukocytosis.
- The presence of *purulent sputum* during an exacerbation can be sufficient indication for starting empirical antibiotic treatment¹¹⁴. *Hemophilus influenzae*, *Streptococcus pneumoniae*, and *Moraxella catarrhalis* are the most common bacterial pathogens involved in an exacerbation⁴⁰⁹; in GOLD 3 and GOLD 4 patients *Pseudomonas aeruginosa* becomes important. If an infectious exacerbation does not respond to the initial antibiotic treatment, a sputum culture and an antibiotic sensitivity test should be performed²⁹¹.
- *Biochemical test abnormalities* including electrolyte disturbances and hyperglycemia can be associated with exacerbations. However, these abnormalities can also be due to associated comorbidities.

Spirometry is not recommended during an exacerbation because it can be difficult to perform and measurements are not accurate enough.

TREATMENT OPTIONS

Treatment Setting

The goals of treatment for COPD exacerbations are to minimize the impact of the current exacerbation and prevent the development of subsequent exacerbations⁴¹⁹. Depending on the severity of an exacerbation and/or the severity of the underlying disease, an exacerbation can be managed in an outpatient or inpatient setting. More than 80% of exacerbations can be managed on an outpatient basis^{132,143,214} with pharmacologic therapies including bronchodilators, corticosteroids, and antibiotics.

Table 5.3 shows the indications for hospital assessment and potential admission of a patient with a COPD exacerbation. When a patient comes to the emergency department the first actions are to provide supplemental oxygen therapy and to determine whether the exacerbation is life-threatening (**Table 5.4**). If so, the patient should be admitted to the ICU immediately. Otherwise, the patient may be managed in the emergency department or hospital as detailed in **Table 5.5**. In addition to pharmacologic therapy, hospital management of exacerbations includes respiratory support (oxygen therapy, ventilation) as detailed in **Table 5.5**.

Table 5.3. Potential Indications for Hospital Assessment or Admission*

- Marked increase in intensity of symptoms, such as sudden development of resting dyspnea
- Severe underlying COPD
- Onset of new physical signs (e.g., cyanosis, peripheral edema)
- Failure of an exacerbation to respond to initial medical management
- Presence of serious comorbidities (e.g., heart failure or newly occurring arrhythmias)
- Frequent exacerbations
- Older age
- Insufficient home support

*Local resources need to be considered.

Long-term prognosis following hospitalization for COPD exacerbation is poor, with a five-year mortality rate of about 50%⁶⁰⁴. Factors independently associated with poor outcome include older age, lower body mass index, comorbidities (e.g., cardiovascular disease or lung cancer), previous admissions for COPD exacerbations, clinical severity of the index exacerbation and need for long-term oxygen therapy at discharge^{605,606}. Patients characterized by a higher prevalence and severity of respiratory symptoms,

poorer quality of life, worse lung function, lower exercise capacity, lower lung density and thickened bronchial walls on CT-scan are also at increased risk of shorter long-term survival following an acute COPD exacerbation⁶⁰⁷.

Table 5.4. Management of Severe but Not Life-Threatening Exacerbations*
<ul style="list-style-type: none"> • Assess severity of symptoms, blood gases, chest radiograph • Administer supplemental oxygen therapy and obtain serial arterial blood gas measurement • Bronchodilators: <ul style="list-style-type: none"> – Increase doses and/or frequency of short-acting bronchodilators – Combine short-acting beta₂-agonists and anticholinergics – Use spacers or air-driven nebulizers • Add oral or intravenous corticosteroids • Consider antibiotics (oral or occasionally intravenous) when signs of bacterial infection • Consider noninvasive mechanical ventilation • At all times: <ul style="list-style-type: none"> – Monitor fluid balance and nutrition – Consider subcutaneous heparin or low molecular weight heparin – Identify and treat associated conditions (e.g., heart failure, arrhythmias) – Closely monitor condition of the patient

*Local resources need to be considered.

Table 5.5. Therapeutic Components of Hospital Management
<p>RESPIRATORY SUPPORT</p> <ul style="list-style-type: none"> Oxygen therapy Ventilatory support <ul style="list-style-type: none"> Noninvasive ventilation Invasive ventilation <p>PHARMACOLOGIC TREATMENT</p> <ul style="list-style-type: none"> Bronchodilators Corticosteroids Antibiotics Adjunct therapies

Pharmacologic Treatment

Long-term prognosis following hospitalization for COPD exacerbation is poor, with a five-year mortality rate of about 50%⁶⁰⁴. Factors independently associated with poor outcome include older age, lower body mass index, comorbidities (e.g., cardiovascular disease or lung cancer), previous admissions for COPD exacerbations, clinical severity of the index exacerbation and need for long-term oxygen therapy at discharge^{605,606}. Patients characterized by a higher prevalence and severity of respiratory symptoms, poorer quality of life, worse lung function, lower exercise capacity, lower lung density and thickened bronchial walls on CT-scan are also at increased risk of shorter long-term survival following an acute COPD exacerbation⁶⁰⁷.

The three classes of medications most commonly used for exacerbations of COPD are bronchodilators, corticosteroids, and antibiotics.

Short-acting Bronchodilators. Although there are no controlled trials, short-acting inhaled beta₂-agonists with or without short-acting anticholinergics are usually the preferred bronchodilators for treatment of an exacerbation^{290,291} (**Evidence C**). There are no clinical studies that have evaluated the use of inhaled long-acting bronchodilators (either beta₂-agonists or anticholinergics) with or without inhaled corticosteroids during an exacerbation. A systematic review of the route of delivery of short-acting bronchodilators found no significant differences in FEV₁ between metered-dose inhalers (with or without a spacer device) and nebulizers⁴²⁰, although the latter can be more convenient for sicker patients. Intravenous methylxanthines (theophylline or aminophylline) are considered second-line therapy, only to be used in selected cases when there is insufficient response to short-acting bronchodilators⁴²¹⁻⁴²⁵ (**Evidence B**). Side effects of methylxanthines are significant and their beneficial effects in terms of lung function and clinical endpoints are modest and inconsistent^{426,427}.

Corticosteroids. Data from studies in secondary health care indicate that systemic corticosteroids in COPD exacerbations shorten recovery time, improve lung function (FEV₁) and arterial hypoxemia (PaO₂)⁴²⁸⁻⁴³¹ (**Evidence A**), and reduce the risk of early relapse, treatment failure⁵⁴¹, and length of hospital stay^{428,430,432}. A dose of 40 mg prednisone per day for 5 days is recommended^{567,624} (**Evidence B**), although there are insufficient data to provide firm conclusions concerning the optimal duration of corticosteroid therapy of acute exacerbations of COPD⁵⁴². Therapy with oral prednisolone is preferable⁴³³. Nebulised budesonide alone may be an alternative (although more expensive) to oral corticosteroids in the treatment of exacerbations^{429,434,435}. Nebulised magnesium as an adjuvant to salbutamol treatment in the setting of acute exacerbations of COPD has no effect on FEV₁⁵⁶⁸.

Antibiotics. Although the infectious agents in COPD exacerbations can be viral or bacterial^{273,436}, the use of antibiotics in exacerbations remains controversial⁵⁶⁹. The uncertainties originate from studies that did not differentiate between bronchitis (acute or chronic) and COPD exacerbations, studies without placebo-control, and/or studies without chest X-rays in which it was unclear if patients had signs of pneumonia. There is evidence supporting the use of antibiotics in exacerbations when patients have clinical signs of a bacterial infection, e.g., increase in sputum purulence¹¹⁴. A systematic review of the very few available placebo-controlled studies has shown that antibiotics reduce the risk of short-term mortality by 77%, treatment failure by 53% and sputum purulence by

44%. This review supports antibiotics for only moderately or severely ill patients with COPD exacerbations with increased cough and sputum purulence^{437,438}. In outpatients, sputum cultures are not feasible as they take too long (at least 2 days) and frequently do not give reliable results for technical reasons, i.e., more than 4 hours elapse between expectoration of sputum and analysis in the microbiology lab. Procalcitonin III, a marker that is specific for bacterial infections, may be of value in the decision to use antibiotics⁴³⁹, but this test is expensive and thus not widely established. A study in COPD patients with exacerbations requiring mechanical ventilation (invasive or noninvasive) indicated that not giving antibiotics was associated with increased mortality and a greater incidence of secondary nosocomial pneumonia⁴⁴⁰.

In summary, antibiotics should be given to patients with exacerbations of COPD who have three cardinal symptoms – increase in dyspnea, sputum volume, and sputum purulence (**Evidence B**); have two of the cardinal symptoms, if increased purulence of sputum is one of the two symptoms (**Evidence C**); or require mechanical ventilation (invasive or noninvasive) (**Evidence B**)^{273,351}. The recommended length of antibiotic therapy is usually 5-10 days (**Evidence D**).

The choice of the antibiotic should be based on the local bacterial resistance pattern. Usually initial empirical treatment is an aminopenicillin with or without clavulanic acid, macrolide, or tetracycline. In patients with frequent exacerbations, severe airflow limitation^{417,441}, and/or exacerbations requiring mechanical ventilation⁴¹², cultures from sputum or other materials from the lung should be performed, as gram-negative bacteria (e.g., *Pseudomonas species*) or resistant pathogens that are not sensitive to the above-mentioned antibiotics may be present. The route of administration (oral or intravenous) depends on the ability of the patient to eat and the pharmacokinetics of the antibiotic, although preferably antibiotics are given orally. Improvements in dyspnea and sputum purulence suggest clinical success.

Adjunct Therapies: Depending on the clinical condition of the patient, an appropriate fluid balance with special attention to the administration of diuretics, anticoagulants, treatment of comorbidities and nutritional aspects should be considered. At all times, health care providers should strongly enforce stringent measures against active cigarette smoking. Given that patients hospitalized because of exacerbations of COPD are at increased risk of deep vein thrombosis and pulmonary embolism^{570,571}, thromboprophylactic measures should be enhanced⁵⁷²⁻⁵⁷⁴.

Respiratory Support

Oxygen therapy. This is a key component of hospital treatment of an exacerbation. Supplemental oxygen should be titrated to improve the patient's hypoxemia with a target saturation of 88-92%⁴⁴². Once oxygen is started, arterial blood gases should be checked 30-60 minutes later to ensure satisfactory oxygenation without carbon dioxide retention or acidosis. Venturi masks (high-flow devices) offer more accurate and controlled delivery of oxygen than do nasal prongs but are less likely to be tolerated by the patient²⁹¹.

Ventilatory Support. Some patients need immediate admission to an intensive care unit (ICU) (**Table 5.6**). Admission of patients with severe exacerbations to intermediate or special respiratory care units may be appropriate if personnel, skills, and equipment exist to identify and manage acute respiratory failure successfully.

Ventilatory support in an exacerbation can be provided by either noninvasive (by nasal or facial mask) or invasive ventilation (by oral tracheal tube or tracheostomy). Respiratory stimulants are not recommended for acute respiratory failure²⁹⁰.

Table 5.6. Indications for ICU Admission *

<ul style="list-style-type: none"> • Severe dyspnea that responds inadequately to initial emergency therapy • Changes in mental status (confusion, lethargy, coma) • Persistent or worsening hypoxemia ($\text{PaO}_2 < 5.3 \text{ kPa}$, 40 mmHg) and/or severe/worsening respiratory acidosis ($\text{pH} < 7.25$) despite supplemental oxygen and noninvasive ventilation • Need for invasive mechanical ventilation • Hemodynamic instability—need for vasopressors
--

*Local resources need to be considered.

Noninvasive mechanical ventilation. The use of noninvasive mechanical ventilation (NIV) has increased significantly over time among patients hospitalized for acute exacerbations of COPD. NIV has been studied in randomized controlled trials showing a success rate of 80-85%^{443-446,543}. NIV has been shown to improve acute respiratory acidosis (increases pH and decreases PaCO_2), decrease respiratory rate, work of breathing, severity of breathlessness, complications such as ventilator associated pneumonia, and length of hospital stay (**Evidence A**). More importantly, mortality and intubation rates are reduced by this intervention^{444,447-449} (**Evidence A**). **Table 5.7** summarizes the indications for NIV⁴⁴³.

Table 5.7. Indications for Noninvasive Mechanical Ventilation^{291,445,451,452}

At least one of the following:

- Respiratory acidosis (arterial pH \leq 7.35 and/or PaCO₂ \geq 6.0 kPa, 45 mm Hg)
- Severe dyspnea with clinical signs suggestive of respiratory muscle fatigue, increased work of breathing, or both, such as use of respiratory accessory muscles, paradoxical motion of the abdomen, or retraction of the intercostal spaces

Table 5.8. Indications for Invasive Mechanical Ventilation

- Unable to tolerate NIV or NIV failure
- Respiratory or cardiac arrest
- Respiratory pauses with loss of consciousness or gasping for air
- Diminished consciousness, psychomotor agitation inadequately controlled by sedation
- Massive aspiration
- Persistent inability to remove respiratory secretions
- Heart rate $<$ 50 min⁻¹ with loss of alertness
- Severe hemodynamic instability without response to fluids and vasoactive drugs
- Severe ventricular arrhythmias
- Life-threatening hypoxemia in patients unable to tolerate NIV

Invasive mechanical ventilation. The indications for initiating invasive mechanical ventilation during an exacerbation are shown in **Table 5.8**, and include failure of an initial trial of NIV⁴⁵⁰. As experience is being gained with the generalized clinical use of NIV in COPD, several indications for invasive mechanical ventilation are being successfully treated with NIV, and in all but a few situations there is nothing to be lost by a trial of noninvasive ventilation⁴⁵⁰. The use of invasive ventilation in very severe COPD patients is influenced by the likely reversibility of the precipitating event, patient's wishes, and availability of intensive care facilities. When possible, a clear statement of the patient's own treatment wishes—an advance directive or “living will”—makes these difficult decisions much easier to resolve. Major hazards include the risk of ventilator-acquired pneumonia (especially when multi-resistant organisms are prevalent), barotrauma, and failure to wean to spontaneous ventilation.

Contrary to some opinions, acute mortality among COPD patients with respiratory failure is lower than mortality among patients ventilated for non-COPD causes⁴⁵³. Despite this, there is evidence that patients who might otherwise survive may be denied admission to intensive care for intubation because of unwarranted prognostic pessimism⁴⁵⁴. A study of a large number of COPD patients with acute respiratory failure reported in-hospital mortality of 17-49%⁴⁰³. Further deaths were reported over the next 12 months, particularly among those patients who had poor lung function before invasive ventilation (FEV₁ $<$ 30% predicted), had a non-respiratory comorbidity, or

Table 5.9. Discharge Criteria

- Able to use long acting bronchodilators, either beta₂-agonists and/or anticholinergics with or without inhaled corticosteroids
- Inhaled short-acting beta₂-agonist therapy is required no more frequently than every 4 hrs
- Patient, if previously ambulatory, is able to walk across room
- Patient is able to eat and sleep without frequent awakening by dyspnea
- Patient has been clinically stable for 12-24 hrs
- Arterial blood gases have been stable for 12-24 hrs
- Patient (or home caregiver) fully understands correct use of medications
- Follow-up and home care arrangements have been completed (e.g., visiting nurse, oxygen delivery, meal provisions)
- Patient, family, and physician are confident that the patient can manage successfully at home

Table 5.10. Checklist of Items to Assess at Time of Discharge from Hospital

- Assurance of effective home maintenance pharmacotherapy regimen
- Reassessment of inhaler technique
- Education regarding role of maintenance regimen
- Instruction regarding completion of steroid therapy and antibiotics, if prescribed
- Assess need for long-term oxygen therapy
- Assure follow-up visit in 4-6 weeks
- Provide a management plan for comorbidities and their follow-up

Table 5.11. Items to Assess at Follow-Up Visit 4-6 Weeks After Discharge from Hospital

- Ability to cope in usual environment
- Measurement of FEV₁
- Reassessment of inhaler technique
- Understanding of recommended treatment regimen
- Reassess need for long-term oxygen therapy and/or home nebulizer
- Capacity to do physical activity and activities of daily living
- CAT or mMRC
- Status of comorbidities

were housebound. Patients who did not have a previously diagnosed comorbidity, had respiratory failure due to a potentially reversible cause (such as an infection), or were relatively mobile and not using long-term oxygen did surprisingly well after ventilatory support.

Weaning or discontinuation from mechanical ventilation can be particularly difficult and hazardous in patients with COPD. The most influential determinant of mechanical ventilatory dependency in these patients is the balance between the respiratory load and the capacity of the respiratory muscles to cope with this load⁴⁵⁵. By contrast, pulmonary gas exchange by itself is not a major difficulty in patients with COPD⁴⁵⁶⁻⁴⁵⁸. Weaning patients from the ventilator can be a very difficult and prolonged process and the best method (pressure support or a T-piece trial) remains a matter of debate⁴⁵⁹⁻⁴⁶¹. In COPD patients that fail extubation, NIV facilitates weaning, prevents reintubation, and reduces mortality^{451,462}. Early NIV after extubation reduces the risk of respiratory failure and lowers 90-day mortality in patients with hypercapnia during a spontaneous breathing trial^{457,462}.

HOSPITAL DISCHARGE AND FOLLOW-UP

Insufficient clinical data exist to establish the optimal duration of hospitalization in individual patients with an exacerbation of COPD⁴⁶³⁻⁴⁶⁵, although units with more respiratory consultants and better organized care have lower mortality and reduced length of hospital stay following admission for an exacerbation⁴⁶⁶. In the hospital prior to discharge, patients should start long-acting bronchodilators, either beta₂-agonists and/or anticholinergics with or without inhaled corticosteroids. Consensus and limited data support the discharge criteria listed in **Table 5.9**. **Table 5.10** provides a checklist of items to assess at time of discharge and **Table 5.11** shows items to assess at follow-up 4 to 6 weeks after discharge from the hospital. Thereafter, follow-up is the same as for stable COPD: supervise smoking cessation, monitor the effectiveness of each medication, and monitor changes in spirometric parameters⁴⁶⁷. Prior hospital admission, oral corticosteroids, use of long-term oxygen therapy, poor health-related quality of life, and lack of routine physical activity have been found to be predictive of readmission⁴⁶⁸.

Home visits by a community nurse may permit earlier discharge of patients hospitalized with an exacerbation without increasing readmission rates^{291,469-472}. Use of a written action plan increases appropriate therapeutic interventions for an exacerbation, an effect that does not decrease health-care resource utilization⁴⁷³ (**Evidence B**) but may shorten recovery time⁴⁷⁴.

For patients who are hypoxemic during an exacerbation, arterial blood gases and/or pulse oximetry should be evaluated prior to hospital discharge and in the following 3 months. If the patient remains hypoxemic, long-term supplemental oxygen therapy may be required.

HOME MANAGEMENT OF EXACERBATIONS

The risk of dying from an exacerbation of COPD is closely related to the development of respiratory acidosis, the presence of significant comorbidities, and the need for ventilatory support⁴⁰². Patients lacking these features are not at high risk of dying. Four randomized clinical trials have shown that nurse-administered home care (also known as “hospital-at-home” care) represents an effective and practical alternative to hospitalization in selected patients with exacerbations of COPD without acidotic respiratory failure^{467-470,544} (**Evidence A**). However, the exact criteria for this approach as opposed to hospital treatment remain uncertain and will vary by health care setting^{469,470}. Treatment recommendations are the same for hospitalized patients. Supported self-management had no effect on time to first readmission or death with COPD⁵⁴⁵. Accumulating data from a variety of studies indicate that telehealth in any of its current forms has not shown benefits for patients with COPD; thus, telehealth is not recommended for use with COPD patients⁵⁷⁵⁻⁵⁷⁷.

PREVENTION OF COPD EXACERBATIONS

COPD exacerbations can often be prevented. Smoking cessation, influenza and pneumococcal vaccines, knowledge of current therapy including inhaler technique, and treatment with long-acting inhaled bronchodilators, with or without inhaled corticosteroids, and possibly phosphodiesterase-4 inhibitors, are all therapies that reduce the number of exacerbations and hospitalizations^{133,134,195,214,264,266}. A large multicenter study indicated that simvastatin has no impact on exacerbation rates^{608,625}. Early outpatient pulmonary rehabilitation after hospitalization for an exacerbation is safe and results in clinically significant improvements in exercise capacity and health status at 3 months⁴⁷⁵. Patients should be encouraged to maintain physical activity, and anxiety, depression and social problems should be discussed. Principal caregivers should be identified if the patient has a significant persisting disability.

CHAPTER

6

***COPD
AND
COMORBIDITIES***

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

CHAPTER 6: COPD AND COMORBIDITIES

KEY POINTS:

- COPD often coexists with other diseases (comorbidities) that may have a significant impact on prognosis.
- In general, the presence of comorbidities should not alter COPD treatment and comorbidities should be treated as if the patient did not have COPD.
- Cardiovascular disease is a major comorbidity in COPD and probably both the most frequent and most important disease coexisting with COPD.
- Osteoporosis and depression are also major comorbidities in COPD, are often under-diagnosed, and are associated with poor health status and prognosis.
- Lung cancer is frequently seen in patients with COPD and has been found to be the most frequent cause of death in patients with mild COPD.
- Gastroesophageal reflux (GERD) is associated with an increased risk of exacerbations and poorer health status.

INTRODUCTION

COPD often coexists with other diseases (comorbidities) that may have a significant impact on prognosis^{100,135,142,476,578,609}. Some of these arise independently of COPD whereas others may be causally related, either with shared risk factors or by one disease actually increasing the risk of another. It is possible that features of COPD, such as systemic inflammation, are shared with other diseases and as such this mechanism represents a link between COPD and some of its comorbidities⁴⁷⁷. This risk of comorbid disease can be increased by the sequelae of COPD; e.g., reduced physical activity. Whether or not COPD and comorbid diseases are related, management of the COPD patient must include identification and treatment of its comorbidities. Importantly, comorbidities with symptoms also associated with COPD may be overlooked; e.g., heart failure and lung cancer (breathlessness) or depression (fatigue and reduced physical activity). Frequent and treatable comorbidities should be prioritized.

Comorbidities are common at any severity of COPD¹³¹ and the differential diagnosis can often be difficult. For example, in a patient with both COPD and heart failure an exacerbation of COPD may be accompanied by worsening

of heart failure.

Below is a brief guide to management of COPD and some comorbidities in stable disease. The recommendations may be insufficient for the management of all patients and cannot substitute for the use of guidelines for the management of each comorbidity.

Cardiovascular Disease (CVD)

CVD is a major comorbidity in COPD and probably both the most frequent and most important disease coexisting with COPD^{135,477}. Four separate entities within CVD will be considered: ischemic heart disease, heart failure, atrial fibrillation and hypertension.

Ischemic Heart Disease (IHD): IHD is increased in COPD, to some extent because of an unfavourable IHD risk profile in COPD patients^{478,479}. There is evidence that concomitant COPD increases morbidity and mortality among patients with IHD⁶¹⁰ and that myocardial injury is overlooked and IHD is therefore under-diagnosed in COPD patients⁴⁸⁰.

Treatment of IHD in patients with COPD: IHD should be treated according to usual IHD guidelines, as there is no evidence that IHD should be treated differently in the presence of COPD. In a significant proportion of patients with IHD a beta-blocker will be indicated, either to treat angina or after a myocardial infarction. Treatment with selective beta₁-blockers is considered safe^{481,546,579,580} but this is based on relatively few short-term studies. The benefits of selective beta₁-blockers when indicated in IHD are, however, considerably larger than the potential risks associated with treatment, even in patients with severe COPD.

Treatment of COPD in patients with IHD: COPD should be treated as usual as there is no evidence that COPD should be treated differently in the presence of IHD. This statement is based on findings from large long-term studies in COPD alone^{195,214,482}, but no large long-term studies exist in patients with both COPD and IHD. Although no studies on COPD medications in patients with unstable angina exist, it seems reasonable to avoid especially high doses of beta-agonists.

Heart Failure (HF): Heart failure is a common comorbidity in COPD. Roughly 30% of patients with stable COPD will have some degree of HF⁴⁸³, and worsening of HF is a significant differential diagnosis to an exacerbation of

COPD. Approximately 30% of patients in a HF clinic have COPD⁴⁸⁴, and comorbid COPD is often the cause of admission for acute HF⁴⁸⁵ – with significant implications for prognosis as FEV₁ is a strong predictor of mortality in HF⁴⁸⁶. HF, COPD and asthma may be confused because of the common cardinal symptom of breathlessness, and caution is warranted for diagnosis and management of these comorbidities.

Treatment of HF in patients with COPD: HF should be treated according to usual HF guidelines as there is no evidence that HF should be treated differently in the presence of COPD. Treatment with selective beta₁-blockers has a significant impact on survival in HF and the presence of COPD is the most significant reason for patients not receiving sufficient therapy⁴⁸⁷. However, as in IHD, treatment with selective beta₁-blockers is considered safe for heart failure patients who also have COPD^{481,579,580}. Studies have shown that treatment with bisoprolol in HF with concomitant COPD decreased FEV₁ but without deleterious effects on symptoms and quality of life⁴⁸⁸ and that a selective beta₁-blocker is indeed preferable to a non-selective beta-blocker in HF with COPD⁴⁸⁹. In a study of patients with moderate-severe airflow limitation and heart failure (NYHA II), treatment with bisoprolol and carvedilol was well tolerated and beneficial effects on lung function were seen. Bisoprolol was superior to carvedilol on respiratory parameters⁵⁴⁷. The benefits of selective beta₁-blocker treatment in HF clearly outweigh any potential risk associated with treatment even in patients with severe COPD.

Treatment of COPD in patients with HF: COPD should be treated as usual as there is no direct evidence that COPD should be treated differently in the presence of HF. As for IHD this statement is based on findings from large long-term studies in patients with HF and comorbid COPD^{195,214,482}. An observational study found an increased risk of death and hospital admission among patients with HF treated with inhaled beta-agonists⁴⁹⁰, possibly indicating a need for close follow-up of patients with severe HF who are on this treatment for COPD.

Atrial Fibrillation (AF): Atrial fibrillation is the most frequent cardiac arrhythmia and COPD patients have an increased incidence of AF⁴⁹¹. COPD with AF presents a challenge to clinicians because of the breathlessness and disability resulting from their coexistence.

Treatment of AF in patients with COPD: AF should be treated according to usual AF guidelines, as there is no evidence that patients with COPD should be treated differently from all other patients. If beta-blockers are used, beta₁-selective drugs are preferred (see considerations

under IHD and HF above).

Treatment of COPD in patients with AF: COPD should be treated as usual; however, there are no good data on the use of COPD medication in patients with AF and these patients have often been excluded from clinical trials. It is a clinical impression that care should be taken when using high doses of beta₂-agonists as this can make appropriate heart rate control difficult.

Hypertension: Hypertension is likely to be the most frequently occurring comorbidity in COPD and has implications for prognosis⁴⁷⁷.

Treatment of hypertension in patients with COPD: Hypertension should be treated according to usual hypertension guidelines, as there is no evidence that hypertension should be treated differently in the presence of COPD. The role of treatment with selective beta-blockers is less prominent in recent hypertension guidelines; if these are used in patients with COPD, a selective beta₁-blocker should be chosen.

Treatment of COPD in patients with hypertension: COPD should be treated as usual as there is no direct evidence that COPD should be treated differently in the presence of hypertension.

Osteoporosis

Osteoporosis is a major comorbidity in COPD^{135,477}, is often under-diagnosed⁴⁹² and is associated with poor health status and prognosis. Osteoporosis may be more closely associated with emphysema than other subgroups of COPD⁴⁹³. Osteoporosis is more often associated with decreased body mass index⁴⁹⁴ and low fat-free mass⁴⁹⁵.

Treatment of osteoporosis in patients with COPD: Osteoporosis should be treated according to usual osteoporosis guidelines. There is no evidence that osteoporosis should be treated differently in the presence of COPD.

Treatment of COPD in patients with osteoporosis: COPD should be treated as usual, as there is no evidence that stable COPD should be treated differently in the presence of osteoporosis. Inhaled triamcinolone was associated with increased loss of bone mass in the Lung Health Study II⁴⁹⁶, whereas this was not the case for inhaled budesonide in the EUROSCOP trial¹⁷⁵ or for inhaled fluticasone propionate in the TORCH trial²⁵⁰. An association between inhaled corticosteroids and fractures has been found in pharmaco-epidemiological studies; however, these studies have not fully taken severity of COPD or exacerbations and their treatment into account.

Systemic corticosteroids significantly increase the risk of osteoporosis and recurrent courses of systemic corticosteroids for COPD exacerbations should be avoided if possible.

Anxiety and Depression

Anxiety and depression are major comorbidities in COPD^{117,497-499} and both are associated with a poor prognosis^{498,500}. Both are often associated with younger age, female gender, smoking, lower FEV₁, cough, higher SGRQ score, and a history of cardiovascular disease^{117,499}.

Treatment of anxiety and depression in patients with COPD: Both disorders should be treated according to usual guidelines, as there is no evidence that anxiety and depression should be treated differently in the presence of COPD. Given the large number of patients who have both depression and COPD, more research on management of depression in COPD patients is needed⁵⁰¹.

Treatment of COPD in patients with anxiety and depression: COPD should be treated as usual as there is no evidence that stable COPD should be treated differently in the presence of anxiety and depression. The potential impact of pulmonary rehabilitation should be stressed as studies have found that physical exercise has a beneficial effect on depression in general⁵⁰².

Lung Cancer

Lung cancer is frequently seen in patients with COPD and has been found to be the most frequent cause of death in patients with mild COPD²¹⁷.

Treatment of lung cancer in patients with COPD: Lung cancer should be treated according to usual lung cancer guidelines, as there is no evidence that lung cancer should be treated differently in the presence of COPD. However, often the reduced lung function of COPD patients will be a factor limiting surgical intervention for lung cancer.

Treatment of COPD in patients with lung cancer: COPD should be treated as usual as there is no evidence that stable COPD should be treated differently in the presence of lung cancer.

Infections

Serious infections, especially respiratory infections, are frequently seen in patients with COPD⁵⁰³.

Treatment of infections in patients with COPD: Macrolide antibiotics increase the serum concentration of theophylline. Apart from this, there is no evidence that

infections should be treated differently in the presence of COPD. However, repeat courses of antibiotics for exacerbations may increase the risk for the presence of antibiotic resistant bacterial strains and more extensive cultures of serious infections may be warranted. Treatment of COPD in patients with infections: COPD should be treated as usual as there is no evidence that stable COPD should be treated differently in the presence of infections. In patients who develop repeated pneumonias while on inhaled corticosteroids, this medication may be stopped in order to observe whether this medication could be the cause of repeated infections.

Metabolic Syndrome and Diabetes

Studies have shown that the presence of metabolic syndrome and manifest diabetes are more frequent in COPD and the latter is likely to impact on prognosis¹⁴².

Treatment of diabetes in patients with COPD: Diabetes should be treated according to usual guidelines for diabetes, as there is no evidence that diabetes should be treated differently in the presence of COPD. However, for patients with severe COPD, it is not advised to aim for a body mass index (BMI) less than 21 kg/m².

Treatment of COPD in patients with diabetes: COPD should be treated as usual as there is no evidence that stable COPD should be treated differently in the presence of diabetes.

Gastroesophageal reflux (GERD) is an independent risk factor for exacerbations and is associated with worse health status. It is, thus a systemic comorbidity that may have an impact on the lungs. The mechanisms responsible for increased risk of exacerbations are not yet fully established and may be more than simply acid reflux. Proton pump inhibitors are often used for treatment of GERD, but the most effective treatment for this condition in COPD has yet to be established⁶¹¹.

Bronchiectasis

Persistent airflow obstruction is a recognized feature of some patients with a primary diagnosis of bronchiectasis. However with increasing use of computed tomography in the assessment of patients with COPD, the presence of previously unrecognized radiographic bronchiectasis is being identified⁵⁸¹. This ranges from mild tubular bronchiectasis to more severe varicose change, although cystic bronchiectasis is uncommon. Whether this radiological change has the same impact as patients with a primary diagnosis of bronchiectasis remains unknown at present, although it is associated with longer exacerbations⁵⁸² and increased mortality⁵⁸³.

Treatment of bronchiectasis in patients with COPD:

Treatment should be along conventional lines for bronchiectasis with the addition of usual COPD strategies where indicated. Whether prevention of exacerbations requires more long-term use of oral or inhaled antibiotics rather than bronchodilator or inhaled corticosteroid therapy remains unknown.

Treatment of COPD in patients with bronchiectasis: COPD should be treated as usual, although some patients may need more aggressive and prolonged antibiotic therapy.

Impaired Cognitive Function

Impaired cognitive function is a feature of COPD⁶¹², and COPD significantly increases the risk of developing mild cognitive impairment⁶¹³. Currently there is no evidence for treatment benefit in such patients, but they should be referred for assessment and treatment in the same way as patients with primary dementia.

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

REFERENCES

1. *World Health Report*. Geneva: World Health Organization. Available from URL: <http://www.who.int/whr/2000/en/statistics.htm>; 2000.
2. Lopez AD, Shibuya K, Rao C, et al. Chronic obstructive pulmonary disease: current burden and future projections. *Eur Respir J* 2006;27:397-412.
3. Pauwels RA, Buist AS, Calverley PM, Jenkins CR, Hurd SS. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop summary. *Am J Respir Crit Care Med* 2001;163:1256-76.
4. Lawrence RS, Mickalide AD, Kamerow DB, Woolf SH. Report of the US Preventive Services Task Force. *JAMA* 1990;263:436-7.
5. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006;3:e442.
6. Salvi SS, Barnes PJ. Chronic obstructive pulmonary disease in non-smokers. *Lancet* 2009;374:733-43.
7. Maciewicz RA, Warburton D, Rennard SI. Can increased understanding of the role of lung development and aging drive new advances in chronic obstructive pulmonary disease? *Proc Am Thorac Soc* 2009;6:614-7.
8. Halbert RJ, Natoli JL, Gano A, Badamgarav E, Buist AS, Mannino DM. Global burden of COPD: systematic review and meta-analysis. *Eur Respir J* 2006;28:523-32.
9. van den Boom G, van Schayck CP, van Mollen MP, et al. Active detection of chronic obstructive pulmonary disease and asthma in the general population. Results and economic consequences of the DIMCA program. *Am J Respir Crit Care Med* 1998;158:1730-8.
10. Fukuchi Y, Nishimura M, Ichinose M, et al. COPD in Japan: the Nippon COPD Epidemiology study. *Respirology* 2004;9:458-65.
11. Menezes AM, Perez-Padilla R, Jardim JR, et al. Chronic obstructive pulmonary disease in five Latin American cities (the PLATINO study): a prevalence study. *Lancet* 2005;366:1875-81.
12. Schirnhofner L, Lamprecht B, Vollmer WM, et al. Results from the Burden of Obstructive Lung Disease (BOLD) Study. *Chest* 2007;131:29-36.
13. Buist AS, McBurnie MA, Vollmer WM, et al. International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet* 2007; 370: 741-50..
14. Pena VS, Miravittles M, Gabriel R, et al. Geographic variations in prevalence and underdiagnosis of COPD: results of the IBERPOC multicentre epidemiological study. *Chest* 2000;118:981-9.
15. Talamo C, de Oca MM, Halbert R, et al. Diagnostic labeling of COPD in five Latin American cities. *Chest* 2007;131:60-7.
16. Jensen HH, Godtfredsen N, Lange P, Vestbo J. Potential misclassification of causes of death from COPD in a Danish population study. *Eur Respir J* 2006;28:781-5.
17. European Respiratory Society. European Lung White Book: Huddersfield, *European Respiratory Society Journals, Ltd*; 2003.
18. National Heart, Lung, and Blood Institute. Morbidity and mortality chartbook on cardiovascular, lung and blood diseases. Bethesda, Maryland: US Department of Health and Human Services, Public Health Service, National Institutes of Health. Accessed at: <http://www.nhlbi.nih.gov/resources/docs/cht-book.htm>; 2009.
19. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet* 1997;349:1498-504.
20. Murray CJL, Lopez AD, editors. In: The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2020. Cambridge, MA: *Harvard University Press*; 1996.
21. Behrendt CE. Mild and moderate-to-severe COPD in non-smokers. Distinct demographic profiles. *Chest* 2005;128:1239-44.
22. Celli BR, Halbert RJ, Nordyke RJ, Schan B. Airway obstruction in never smokers: results from the Third National Health and Nutrition Examination Survey. *Am J Med* 2005;118:1364-72.

23. Eisner MD, Anthonisen N, Coultas D, et al. An official American Thoracic Society public policy statement: Novel risk factors and the global burden of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2010;182:693-718.
24. Lamprecht B, McBurnie MA, Vollmer WM, et al. COPD in never smokers: results from the population-based burden of obstructive lung disease study. *Chest* 2011;139:752-63.
25. Anthonisen NR, Connett JE, Murray RP. Smoking and lung function of Lung Health Study participants after 11 years. *Am J Respir Crit Care Med* 2002;166:675-9.
26. Stoller JK, Aboussouan LS. Alpha1-antitrypsin deficiency. *Lancet* 2005;365:2225-36.
27. McCloskey SC, Patel BD, Hinchliffe SJ, Reid ED, Wareham NJ, Lomas DA. Siblings of patients with severe chronic obstructive pulmonary disease have a significant risk of airflow obstruction. *Am J Respir Crit Care Med* 2001;164:1419-24.
28. Hunninghake GM, Cho MH, Tesfaigzi Y, et al. MMP12, lung function, and COPD in high-risk populations. *N Engl J Med* 2009;361:2599-608.
29. Castaldi PJ, Cho MH, Cohn M, Langerman F, Moran S, Tarragona N, et al. The COPD genetic association compendium: a comprehensive online database of COPD genetic associations. *Hum Mol Genet* 2010;19:526-34.
30. Cho MH, Boutaoui N, Klanderman BJ, Sylvia JS, Ziniti JP, Hersh CP, et al. Variants in FAM13A are associated with chronic obstructive pulmonary disease. *Nat Genet* 2010;42:200-2.
31. Pillai SG, Ge D, Zhu G, Kong X, Shianna KV, Need AC, et al. A genome-wide association study in chronic obstructive pulmonary disease (COPD): identification of two major susceptibility loci. *PLoS Genet* 2009;5:e1000421.
32. Repapi E, Sayers I, Wain LV, et al. Genome-wide association study identifies five loci associated with lung function. *Nat Genet* 2010;42:36-44.
33. Wilk JB, Chen TH, Gottlieb DJ, Walter RE, Nagle MW, Brandler BJ, et al. A genome-wide association study of pulmonary function measures in the Framingham Heart Study. *PLoS Genet* 2009;5:e1000429.
34. Mannino DM, Homa DM, Akinbami LJ, Ford ES, Redd SC. Chronic obstructive pulmonary disease surveillance-United States, 1971-2000. *MMWR Surveill Summ* 2002;51:1-16.
35. Foreman MG, Zhang L, Murphy J, et al. Early-onset chronic obstructive pulmonary disease is associated with female sex, maternal factors, and African American race in the COPD Gene Study. *Am J Respir Crit Care Med* 2011;184:414-20.
36. Lopez Varela MV, Montes de Oca M, Halbert RJ, et al. Sex-related differences in COPD in five Latin American cities: the PLATINO study. *The European respiratory journal : official journal of the European Society for Clinical Respiratory Physiology* 2010;36:1034-41.
37. Silverman EK, Weiss ST, Drazen JM, et al. Gender-related differences in severe, early-onset chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000;162:2152-8.
38. Sorheim IC, Johannessen A, Gulsvik A, Bakke PS, Silverman EK, DeMeo DL. Gender differences in COPD: are women more susceptible to smoking effects than men? *Thorax* 2010;65:480-5.
39. Barker DJ, Godfrey KM, Fall C, Osmond C, Winter PD, Shaheen SO. Relation of birth weight and childhood respiratory infection to adult lung function and death from chronic obstructive airways disease. *BMJ* 1991;303:671-5.
40. Todisco T, de Benedictis FM, Iannacci L, et al. Mild prematurity and respiratory functions. *Eur J Pediatr* 1993;152:55-8.
41. Stern DA, Morgan WJ, Wright AL, Guerra S, Martinez FD. Poor airway function in early infancy and lung function by age 22 years: a non-selective longitudinal cohort study. *Lancet* 2007;370:758-64.
42. Lawlor DA, Ebrahim S, Davey Smith G. Association of birth weight with adult lung function: findings from the British Women's Heart and Health Study and a meta-analysis. *Thorax* 2005;60:851-8.
43. Svanes C, Sunyer J, Plana E. Early life origins of chronic obstructive pulmonary disease. *Thorax* 2010;65:14-20.

44. Kohansal R, Martinez-Cambor P, Agusti A, Buist AS, Mannino DM, Soriano JB. The natural history of chronic airflow obstruction revisited: an analysis of the Framingham offspring cohort. *Am J Respir Crit Care Med* 2009;180:3-10.
45. Raad D, Gaddam S, Schunemann HJ, et al. Effects of water-pipe smoking on lung function: a systematic review and meta-analysis. *Chest* 2011;139:764-74.
46. Tan WC, Lo C, Jong A, et al. Marijuana and chronic obstructive lung disease: a population-based study. *CMAJ* 2009;180:814-20.
47. WHO Study Group on Tobacco Product Regulation. Water Pipe smoking: health effects, research needs, and recommended actions by regulators. *World Health Organization Publication*. ISBN 92 4 159385. 2005.
48. Tetrault JM, Crothers K, Moore BA, Mehra R, Concato J, Fiellin DA. Effects of marijuana smoking on pulmonary function and respiratory complications: a systematic review. *Arch Intern Med* 2007;167:221-8.
49. The Health Consequences of Involuntary Exposure to Tobacco Smoke: *A Report of the Surgeon General, Department of Health and Human Services*. Washington, DC, US; 2006.
50. Eisner MD, Balmes J, Katz BP, Trupin L, Yelin E, Blanc P. Lifetime environmental tobacco smoke exposure and the risk of chronic obstructive pulmonary disease. *Environ Health Perspect* 2005;4:7-15.
51. Dayal HH, Khuder S, Sharrar R, Trief N. Passive smoking in obstructive respiratory disease in an industrialized urban population. *Environ Res* 1994;65:161-71.
52. Leuenberger P, Schwartz J, Ackermann-Liebrich U, et al. Passive smoking exposure in adults and chronic respiratory symptoms (SAPALDIA Study). Swiss Study on Air Pollution and Lung Diseases in Adults, SAPALDIA Team. *Am J Respir Crit Care Med* 1994;150:1222-8.
53. Holt PG. Immune and inflammatory function in cigarette smokers. *Thorax* 1987;42:241-9.
54. Tager IB, Ngo L, Hanrahan JP. Maternal smoking during pregnancy. Effects on lung function during the first 18 months of life. *Am J Respir Crit Care Med* 1995;152:977-83.
55. Trupin L, Earnest G, San Pedro M, et al. The occupational burden of chronic obstructive pulmonary disease. *Eur Respir J* 2003;22:462-9.
56. Matheson MC, Benke G, Raven J, et al. Biological dust exposure in the workplace is a risk factor for chronic obstructive pulmonary disease. *Thorax* 2005;60:645-51.
57. Hnizdo E, Sullivan PA, Bang KM, Wagner G. Airflow obstruction attributable to work in industry and occupation among U.S. race/ethnic groups: a study of NHANES III data. *Am J Ind Med* 2004;46:126-35.
58. Hnizdo E, Sullivan PA, Bang KM, Wagner G. Association between chronic obstructive pulmonary disease and employment by industry and occupation in the US population: a study of data from the Third National Health and Nutrition Examination Survey. *Am J Epidemiol* 2002;156:738-46.
59. Balmes J, Becklake M, Blanc P, et al. American Thoracic Society Statement: Occupational contribution to the burden of airway disease. *Am J Respir Crit Care Med* 2003;167:787-97.
60. Boman C, Forsberg B, Sandstrom T. Shedding new light on wood smoke: a risk factor for respiratory health. *Eur Respir J* 2006;27:446-7.
61. Ezzati M. Indoor air pollution and health in developing countries. *Lancet* 2005;366:104-6.
62. Mishra V, Dai X, Smith KR, Mika L. Maternal exposure to biomass smoke and reduced birth weight in Zimbabwe. *Ann Epidemiol* 2004;14:740-7.
63. Orozco-Levi M, Garcia -Aymerich J, Villar J, Ramirez-Sarmiento A, Anto JM, Gea J. Wood smoke exposure and risk of chronic obstructive pulmonary disease. *Eur Respir J* 2006;27:542-6.
64. Sezer H, Akkurt I, Guler N, Marakoglu K, Berk S. A case-control study on the effect of exposure to different substances on the development of COPD. *Ann Epidemiol* 2006;16:59-62.
65. Smith KR, Mehta S, Maeusezahl-Feuz M. Indoor air-pollution from household solid fuel use. In: Ezzati, M., Lopez, A. D., Rodgers, M., Murray, C. J., eds. Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors. Geneva: *World Health Organization*; 2004.

66. Warwick H, Doig A. Smoke the killer in the kitchen: Indoor air pollution in developing countries. ITDG Publishing, 103-105 Southampton Row, London WC1B HLD, UK 2004:URL: <http://www.itdgpublishing.org.uk>.
67. Torres-Duque C, Maldonado D, Perez-Padilla R, Ezzati M, Viegi G. Biomass fuels and respiratory diseases: a review of the evidence. *Proc Am Thorac Soc* 2008;5:577-90.
68. Abbey DE, Burchette RJ, Knutsen SF, McDonnell WF, Lebowitz MD, Enright PL. Long-term particulate and other air pollutants and lung function in nonsmokers. *Am J Respir Crit Care Med* 1998;158:289-98.
69. Prescott E, Lange P, Vestbo J. Socioeconomic status, lung function and admission to hospital for COPD: results from the Copenhagen City Heart Study. *Eur Respir J* 1999;13:1109-14.
70. Silva GE, Sherrill DL, Guerra S, Barbee RA. Asthma as a risk factor for COPD in a longitudinal study. *Chest* 2004;126:59-65.
71. Vonk JM, Jongepier H, Panhuysen CI, Schouten JP, Bleecker ER, Postma DS. Risk factors associated with the presence of irreversible airflow limitation and reduced transfer coefficient in patients with asthma after 26 years of follow up. *Thorax* 2003;58:322-7.
72. Lange P, Parner J, Vestbo J, Schnohr P, Jensen G. A 15-year follow-up study of ventilatory function in adults with asthma. *N Engl J Med* 1998;339:1194-200.
73. de Marco R, Accordini S, Marcon A, et al. Risk factors for chronic obstructive pulmonary disease in a European cohort of young adults. *Am J Respir Crit Care Med* 2011;183:891-7.
74. Fabbri LM, Romagnoli M, Corbetta L, et al. Differences in airway inflammation in patients with fixed airflow obstruction due to asthma or chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003;167:418-24.
75. Rijcken B, Schouten JP, Weiss ST, Speizer FE, van der Lende R. The relationship of nonspecific bronchial responsiveness to respiratory symptoms in a random population sample. *Am Rev Respir Dis* 1987;136:62-8.
76. Tashkin DP, Altose MD, Connett JE, Kanner RE, Lee WW, Wise RA. Methacholine reactivity predicts changes in lung function over time in smokers with early chronic obstructive pulmonary disease. The Lung Health Study Research Group. *Am J Respir Crit Care Med* 1996;153:1802-11.
77. Fletcher C, Peto R. The natural history of chronic airflow obstruction. *BMJ* 1977;1:1645-8.
78. Vestbo J, Prescott E, Lange P, Group at CCHS. Association between chronic mucus hypersecretion with FEV₁ decline and COPD morbidity. *Am J Respir Crit Care Med* 1996;153:1530-5.
79. Guerra S, Sherrill DL, Venker C, Ceccato CM, Halonen M, F.D. M. Chronic bronchitis before age 50 years predicts incident airflow limitation and mortality risk. *Thorax* 2009;64:894-900.
80. de Marco R, Accordini S, Cerveri I, et al. Incidence of chronic obstructive pulmonary disease in a cohort of young adults according to the presence of chronic cough and phlegm. *Am J Respir Crit Care Med* 2007;175:32-9.
81. Crothers K HL, Goulet JL, Goetz MB, Brown ST. HIV infection and risk for incident pulmonary diseases in the combination antiretroviral therapy era. *Am J Respir Crit Care Med* 2011;183:388-95.
82. Lam KB, Jiang CQ, Jordan RE, et al. Prior TB, smoking, and airflow obstruction: a cross-sectional analysis of the Guangzhou Biobank Cohort Study. *Chest*;137:593-600.
83. Menezes AM, Hallal PC, Perez-Padilla R, et al. Tuberculosis and airflow obstruction: evidence from the PLATINO study in Latin America. *Eur Respir J* 2007;30:1180-5.
84. Jordan TS, Spencer EM, Davies P. Tuberculosis, bronchiectasis and chronic airflow obstruction. *Respirology* 2010;15:623-8.
85. Barnes PJ, Shapiro SD, Pauwels RA. Chronic obstructive pulmonary disease: molecular and cellular mechanisms. *Eur Respir J* 2003;22:672-88.
86. Hogg JC. Pathophysiology of airflow limitation in chronic obstructive pulmonary disease. *Lancet* 2004;364:709-21.
87. Cosio MG, Saetta M, Agusti A. Immunologic aspects of chronic obstructive pulmonary disease. *N Engl J Med* 2009;360:2445-54.

88. Rahman I. Oxidative stress in pathogenesis of chronic obstructive pulmonary disease: cellular and molecular mechanisms. *Cell Biochem Biophys* 2005;43:167-88.
89. Malhotra D, Thimmulappa R, Navas-Acien A, et al. Decline in NRF2-regulated antioxidants in chronic obstructive pulmonary disease lungs due to loss of its positive regulator, DJ-1. *Am J Respir Crit Care Med* 2008;178:592-604.
90. Hogg JC, Chu F, Utokaparch S, et al. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 2004;350:2645-53.
91. Bestall JC, Paul EA, Garrod R, Garnham R, Jones PW, Wedzicha JA. Usefulness of the Medical Research Council (MRC) dyspnoea scale as a measure of disability in patients with chronic obstructive pulmonary disease. *Thorax* 1999;54:581-6.
92. Nishimura K, Izumi T, Tsukino M, Oga T. Dyspnea is a better predictor of 5-year survival than airway obstruction in patients with COPD. *Chest* 2002;121:1434-40.
93. O'Donnell DE, Laveneziana P. Dyspnea and activity limitation in COPD: mechanical factors. *COPD* 2007;4:225-36.
94. O'Donnell DE, Laveneziana P, Ora J, Webb KA, Lam VM, Ofir D. Evaluation of acute bronchodilator reversibility in patients with symptoms of GOLD stage I COPD. *Thorax* 2009;64:216-23.
95. Rodriguez-Roisin R, Drakulovic M, Rodriguez DA, Roca J, Barbera JA, Wagner PD. Ventilation-perfusion imbalance and chronic obstructive pulmonary disease staging severity. *J Appl Physiol* 2009;106:1902-8.
96. Burgel PR, Nadel JA. Roles of epidermal growth factor receptor activation in epithelial cell repair and mucin production in airway epithelium. *Thorax* 2004;59:992-6.
97. Peinado VI, Pizarro S, Barbera JA. Pulmonary vascular involvement in COPD. *Chest* 2008;134:808-14.
98. Parker CM, Voduc N, Aaron SD, Webb KA, O'Donnell DE. Physiological changes during symptom recovery from moderate exacerbations of COPD. *Eur Respir J* 2005;26:420-8.
99. Barbera JA, Roca J, Ferrer A, et al. Mechanisms of worsening gas exchange during acute exacerbations of chronic obstructive pulmonary disease. *Eur Respir J* 1997;10:1285-91.
100. Barnes PJ, Celli BR. Systemic manifestations and comorbidities of COPD. *Eur Respir J* 2009;33:1165-85.
101. Barr RG, Bluemke DA, Ahmed FS, et al. Percent emphysema, airflow obstruction, and impaired left ventricular filling. *N Engl J Med* 2010;362:217-27.
102. Hole DJ, Watt GC, Davey-Smith G, Hart CL, Gillis CR, Hawthorne VM. Impaired lung function and mortality risk in men and women: findings from the Renfrew and Paisley prospective population study. *BMJ* 1996;313:711-5; discussion 5-6.
103. Young RP, Hopkins RJ, Christmas T, Black PN, Metcalf P, Gamble GD. COPD prevalence is increased in lung cancer, independent of age, sex and smoking history. *Eur Respir J* 2009;34:380-6.
104. Screening for chronic obstructive pulmonary disease using spirometry: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2008;148:529-34.
105. Hardie JA, Buist AS, Vollmer WM, Ellingsen I, Bakke PS, Morkve O. Risk of over-diagnosis of COPD in asymptomatic elderly never-smokers. *Eur Respir J* 2002;20:1117-22.
106. Cerveri I, Corsico AG, Accordini S, et al. Underestimation of airflow obstruction among young adults using FEV1/FVC <70% as a fixed cut-off: a longitudinal evaluation of clinical and functional outcomes. *Thorax* 2008;63:1040-5.
107. Mohamed Hoesein FA, Zanen P, Lammers JW. Lower limit of normal or FEV(1)/FVC <0.70 in diagnosing COPD: An evidence-based review. *Respir Med* 2011;105:907-15.
108. Simon PM, Schwartzstein RM, Weiss JW, Fencel V, Teghtsoonian M, Weinberger SE. Distinguishable types of dyspnea in patients with shortness of breath. *Am Rev Respir Dis* 1990;142:1009-14.
109. Elliott MW, Adams L, Cockcroft A, MacRae KD, Murphy K, Guz A. The language of breathlessness. Use of verbal descriptors by patients with cardiopulmonary disease. *Am Rev Respir Dis* 1991;144:826-32.

110. Georgopoulos D, Anthonisen NR. Symptoms and signs of COPD. In: Cherniack NS, ed. Chronic obstructive pulmonary disease. Toronto: WB Saunders Co; 1991:357-63.
111. Burrows B, Niden AH, Barclay WR, Kasik JE. Chronic obstructive lung disease II. Relationships of clinical and physiological findings to the severity of airways obstruction. *Am Rev Respir Dis* 1965;91:665-78.
112. Definition and classification of chronic bronchitis for clinical and epidemiological purposes. A report to the Medical Research Council by their Committee on the Aetiology of Chronic Bronchitis. *Lancet* 1965;1:775-9.
113. Hill AT, Bayley D, Stockley RA. The interrelationship of sputum inflammatory markers in patients with chronic bronchitis. *Am J Respir Crit Care Med* 1999;160:893-8.
114. Stockley RA, O'Brien C, Pye A, Hill SL. Relationship of sputum color to nature and outpatient management of acute exacerbations of COPD. *Chest* 2000;117:1638-45.
115. Schols AM, Soeters PB, Dingemans AM, Mostert R, Frantzen PJ, Wouters EF. Prevalence and characteristics of nutritional depletion in patients with stable COPD eligible for pulmonary rehabilitation. *Am Rev Respir Dis* 1993;147:1151-6.
116. Schols AM, Slangen J, Volovics L, Wouters EF. Weight loss is a reversible factor in the prognosis of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;157:1791-7.
117. Hanania NA, Mullerova H, Locantore NW, et al. Determinants of depression in the ECLIPSE chronic obstructive pulmonary disease cohort. *Am J Respir Crit Care Med* 2011;183:604-11.
118. Holguin F, Folch E, Redd SC, Mannino DM. Comorbidity and mortality in COPD-related hospitalizations in the United States, 1979 to 2001. *Chest* 2005;128:2005-11.
119. Kesten S, Chapman KR. Physician perceptions and management of COPD. *Chest* 1993;104:254-8.
120. Loveridge B, West P, Kryger MH, Anthonisen NR. Alteration in breathing pattern with progression of chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1986;134:930-4.
121. Jackson H, Hubbard R. Detecting chronic obstructive pulmonary disease using peak flow rate: cross sectional survey. *BMJ* 2003;327:653-4.
122. Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. *Eur Respir J* 2005;26:948-68.
123. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J* 2005;26:319-38.
124. Jones PW, Harding G, Berry P, Wiklund I, Chen WH, Kline Leidy N. Development and first validation of the COPD Assessment Test. *Eur Respir J* 2009;34:648-54.
125. Dodd JW, Hogg L, Nolan J, et al. The COPD assessment test (CAT): response to pulmonary rehabilitation. A multicentre, prospective study. *Thorax* 2011;66:425-9.
126. Burge PS, Calverley PM, Jones PW, Spencer S, Anderson JA, Maslen TK. Randomised, double blind, placebo controlled study of fluticasone propionate in patients with moderate to severe chronic obstructive pulmonary disease: the ISOLDE trial. *BMJ* 2000;320:1297-303.
127. Jones PW. Health status and the spiral of decline. *COPD* 2009;6:59-63.
128. Rodriguez-Roisin R. Toward a consensus definition for COPD exacerbations. *Chest* 2000;117:398S-401S.
129. Burge S, Wedzicha JA. COPD exacerbations: definitions and classifications. *Eur Respir J Suppl* 2003;41:46s-53s.
130. Celli BR, Barnes PJ. Exacerbations of chronic obstructive pulmonary disease. *Eur Respir J* 2007;29:1224-38.
131. Agusti A, Calverley PM, Celli B, et al. Characterisation of COPD heterogeneity in the ECLIPSE cohort. *Respir Res* 2010;11:122.
132. Hurst JR, Vestbo J, Anzueto A, et al. Susceptibility to exacerbation in chronic obstructive pulmonary disease. *N Engl J Med* 2010;363:1128-38.
133. Decramer M, Celli B, Kesten S, Lystig T, Mehra S, Tashkin DP. Effect of tiotropium on outcomes in patients with moderate chronic obstructive pulmonary disease (UPLIFT): a prespecified subgroup analysis of a randomised controlled trial. *Lancet* 2009;374:1171-8.

134. Jenkins CR, Jones PW, Calverley PM, et al. Efficacy of salmeterol/fluticasone propionate by GOLD stage of chronic obstructive pulmonary disease: analysis from the randomised, placebo-controlled TORCH study. *Respir Res* 2009;10:59.
135. Soriano JB, Visick GT, Muellerova H, Payvandi N, Hansell AL. Patterns of comorbidities in newly diagnosed COPD and asthma in primary care. *Chest* 2005;128:2099-107.
136. Wagner PD. Possible mechanisms underlying the development of cachexia in COPD. *Eur Respir J* 2008;31:492-501.
137. American Thoracic Society and European Respiratory Society. Skeletal muscle dysfunction in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:S1-40.
138. Lange P, Nyboe J, Appleyard M, Jensen G, Schnohr P. Ventilatory function and chronic mucus hypersecretion as predictors of death from lung cancer. *Am Rev Respir Dis* 1990;141:613-7.
139. Skillrud DM, Offord KP, Miller RD. Higher risk of lung cancer in chronic obstructive pulmonary disease. A prospective, matched, controlled study. *Ann Intern Med* 1986;105:503-7.
140. Stavem K, Aaser E, Sandvik L, et al. Lung function, smoking and mortality in a 26-year follow-up of healthy middle-aged males. *Eur Respir J* 2005;25:618-25.
141. Tockman MS, Anthonisen NR, Wright EC, Donithan MG. Airways obstruction and the risk for lung cancer. *Ann Intern Med* 1987;106:512-8.
142. Mannino DM, Thorn D, Swensen A, Holguin F. Prevalence and outcomes of diabetes, hypertension and cardiovascular disease in COPD. *Eur Respir J* 2008;32:962-9.
143. Celli BR, Thomas NE, Anderson JA, et al. Effect of pharmacotherapy on rate of decline of lung function in chronic obstructive pulmonary disease: results from the TORCH study. *Am J Respir Crit Care Med* 2008;178:332-8.
144. Spencer S, Calverley PM, Burge PS, Jones PW. Impact of preventing exacerbations on deterioration of health status in COPD. *Eur Respir J* 2004;23:698-702.
145. Jones P, Tabberer M, Chen W-H. Creating scenarios of the impact of copd and their relationship to copd assessment test (CATTM) scores. *BMC Pulmonary Medicine* 2011;11:42.
146. Fishman A, Martinez F, Naunheim K, et al. A randomized trial comparing lung-volume-reduction surgery with medical therapy for severe emphysema. *N Engl J Med* 2003;348:2059-73.
147. Kelly AM, McAlpine R, Kyle E. How accurate are pulse oximeters in patients with acute exacerbations of chronic obstructive airways disease? *Respiratory medicine* 2001;95:336-40.
148. Alpha-1 Antitrypsin Deficiency: Memorandum from a WHO Meeting. *Bulletin of the World Health Organization* 1997;75:397-415.
149. Pinto-Plata VM, Cote C, Cabral H, Taylor J, Celli BR. The 6-min walk distance: change over time and value as a predictor of survival in severe COPD. *Eur Respir J* 2004;23:28-33.
150. Oga T, Nishimura K, Tsukino M, Sato S, Hajiro T. Analysis of the factors related to mortality in chronic obstructive pulmonary disease: role of exercise capacity and health status. *Am J Respir Crit Care Med* 2003;167:544-9.
151. Jones PW. Health status measurement in chronic obstructive pulmonary disease. *Thorax* 2001;56:880-7.
152. Revall SM, Morgan MD, Singh SJ, Williams J, Hardman AE. The endurance shuttle walk: a new field test for the assessment of endurance capacity in chronic obstructive pulmonary disease. *Thorax* 1999;54:213-22.
153. Singh SJ, Morgan MD, Scott S, Walters D, Hardman AE. Development of a shuttle walking test of disability in patients with chronic airways obstruction. *Thorax* 1992;47:1019-24.
154. Casanova C, Cote CG, Marin JM, et al. The 6-min walking distance: long-term follow up in patients with COPD. *Eur Respir J* 2007;29:535-40.
155. Waschki B KA, Holz O, Muller KC, Meyer T, Watz H, Magnussen H. Physical activity is the strongest predictor of all-cause mortality in patients with chronic obstructive pulmonary disease: a prospective cohort study. *Chest* 2011; Aug;140(2):331-42.

156. Celli BR, Cote CG, Marin JM, et al. The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Engl J Med* 2004;350:1005-12.
157. Jones RC, Donaldson GC, Chavannes NH, et al. Derivation and validation of a composite index of severity in chronic obstructive pulmonary disease: the DOSE Index. *Am J Respir Crit Care Med* 2009;180:1189-95.
158. Puhan MA, Garcia-Aymerich J, Frey M, et al. Expansion of the prognostic assessment of patients with chronic obstructive pulmonary disease: the updated BODE index and the ADO index. *Lancet* 2009;374:704-11.
159. Anthonisen NR, Connett JE, Kiley JP, et al. Effects of smoking intervention and the use of an inhaled anticholinergic bronchodilator on the rate of decline of FEV1. The Lung Health Study. *JAMA* 1994;272:1497-505.
160. The tobacco use and dependence clinical practice guideline panel, staff, and consortium representatives. A clinical practice guideline for treating tobacco use and dependence. *JAMA* 2000;28:3244-54.
161. Lancaster T, Stead L, Silagy C, Sowden A. Effectiveness of interventions to help people stop smoking: findings from the Cochrane Library. *BMJ* 2000;321:355-8.
162. Tonnesen P, Mikkelsen K, Bremann L. Nurse-conducted smoking cessation in patients with COPD using nicotine sublingual tablets and behavioral support. *Chest* 2006;130:334-42.
163. Fiore MC, Bailey WC, Cohen SJ. Smoking cessation: information for specialists. Rockville, MD: *US Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research and Centers for Disease Control and Prevention*; 1996.
164. Tashkin DP, Rennard S, Hays JT, Ma W, Lawrence D, Lee TC. Effects of varenicline on smoking cessation in patients with mild to moderate COPD: a randomized controlled trial. *Chest* 2011;139:591-9.
165. Tashkin D, Kanner R, Bailey W, et al. Smoking cessation in patients with chronic obstructive pulmonary disease: a double-blind, placebo-controlled, randomised trial. *Lancet* 2001;357:1571-5.
166. Jorenby DE, Leischow SJ, Nides MA, et al. A controlled trial of sustained-release bupropion, a nicotine patch, or both for smoking cessation. *N Engl J Med* 1999;340:685-91.
167. American Medical Association. Guidelines for the diagnosis and treatment of nicotine dependence: how to help patients stop smoking. Washington DC: *American Medical Association*; 1994.
168. Glynn TJ, Manley MW. How to help your patients stop smoking. A National Cancer Institute manual for physicians. Bethesda, MD: *US Department of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute*; 1990.
169. Glynn TJ, Manley MW, Pechacek TF. Physician-initiated smoking cessation program: the National Cancer Institute trials. *Prog Clin Biol Res* 1990;339:11-25.
170. Baillie AJ, Mattick RP, Hall W, Webster P. Meta-analytic review of the efficacy of smoking cessation interventions. *Drug and Alcohol Review* 1994;13:157-70.
171. Wilson DH, Wakefield MA, Steven ID, Rohrsheim RA, Esterman AJ, Graham NM. "Sick of Smoking": evaluation of a targeted minimal smoking cessation intervention in general practice. *Med J Aust* 1990;152:518-21.
172. Kottke TE, Battista RN, DeFries GH, Brekke ML. Attributes of successful smoking cessation interventions in medical practice. A meta-analysis of 39 controlled trials. *JAMA* 1988;259:2883-9.
173. Ockene JK, Kristeller J, Goldberg R, et al. Increasing the efficacy of physician-delivered smoking interventions: a randomized clinical trial. *J Gen Intern Med* 1991;6:1-8.
174. Katz DA, Muehlenbruch DR, Brown RL, Fiore MC, Baker TB. Effectiveness of implementing the agency for healthcare research and quality smoking cessation clinical practice guideline: a randomized, controlled trial. *J Natl Cancer Inst* 2004;96:594-603.
175. Pauwels RA, Lofdahl CG, Laitinen LA, et al. Long-term treatment with inhaled budesonide in persons with mild chronic obstructive pulmonary disease who continue smoking. European Respiratory Society Study on Chronic Obstructive Pulmonary Disease. *N Engl J Med* 1999;340:1948-53.

176. Vestbo J, Sorensen T, Lange P, Brix A, Torre P, Viskum K. Long-term effect of inhaled budesonide in mild and moderate chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet* 1999;353:1819-23.
177. Al-Showair RA, Tarsin WY, Assi KH, Pearson SB, Chrystyn H. Can all patients with COPD use the correct inhalation flow with all inhalers and does training help? *Respir Med* 2007;101:2395-401.
178. Ericsson CH, Svartengren K, Svartengren M, et al. Repeatability of airway deposition and tracheobronchial clearance rate over three days in chronic bronchitis. *Eur Respir J* 1995;8:1886-93.
179. Kim CS, Kang TC. Comparative measurement of lung deposition of inhaled fine particles in normal subjects and patients with obstructive airway disease. *Am J Respir Crit Care Med* 1997;155:899-905.
180. Calverley PMA. Symptomatic bronchodilator treatment. In: Calverley PMA, Pride NB, eds. Chronic obstructive pulmonary disease. London: *Chapman and Hall*; 1995:419-45.
181. O'Donnell DE, Fluge T, Gerken F, et al. Effects of tiotropium on lung hyperinflation, dyspnoea and exercise tolerance in COPD. *Eur Respir J* 2004;23:832-40.
182. O'Donnell DE, Sciruba F, Celli B, et al. Effect of fluticasone propionate/salmeterol on lung hyperinflation and exercise endurance in COPD. *Chest* 2006;130:647-56.
183. Berger R, Smith D. Effect of inhaled metaproterenol on exercise performance in patients with stable "fixed" airway obstruction. *Am Rev Respir Dis* 1988;138:624-9.
184. Hay JG, Stone P, Carter J, et al. Bronchodilator reversibility, exercise performance and breathlessness in stable chronic obstructive pulmonary disease. *Eur Respir J* 1992;5:659-64.
185. Chrystyn H, Mulley BA, Peake MD. Dose response relation to oral theophylline in severe chronic obstructive airways disease. *BMJ* 1988;297:1506-10.
186. Gross NJ, Petty TL, Friedman M, Skorodin MS, Silvers GW, Donohue JF. Dose response to ipratropium as a nebulized solution in patients with chronic obstructive pulmonary disease. A three-center study. *Am Rev Respir Dis* 1989;139:1188-91.
187. Higgins BG, Powell RM, Cooper S, Tattersfield AE. Effect of salbutamol and ipratropium bromide on airway calibre and bronchial reactivity in asthma and chronic bronchitis. *Eur Respir J* 1991;4:415-20.
188. Vathenen AS, Britton JR, Ebden P, Cookson JB, Wharrad HJ, Tattersfield AE. High-dose inhaled albuterol in severe chronic airflow limitation. *Am Rev Respir Dis* 1988;138:850-5.
189. O'Driscoll BR, Kay EA, Taylor RJ, Weatherby H, Chetty MC, Bernstein A. A long-term prospective assessment of home nebulizer treatment. *Respir Med* 1992;86:317-25.
190. Jenkins SC, Heaton RW, Fulton TJ, Moxham J. Comparison of domiciliary nebulized salbutamol and salbutamol from a metered-dose inhaler in stable chronic airflow limitation. *Chest* 1987;91:804-7.
191. COMBIVENT Inhalation Aerosol Study Group. In chronic obstructive pulmonary disease, a combination of ipratropium and albuterol is more effective than either agent alone. An 85-day multicenter trial. *Chest* 1994;105:1411-9.
192. van Schayck CP, Folgering H, Harbers H, Maas KL, van Weel C. Effects of allergy and age on responses to salbutamol and ipratropium bromide in moderate asthma and chronic bronchitis. *Thorax* 1991;46:355-9.
193. Sestini P, Cappiello V, Aliani M, et al. Prescription bias and factors associated with improper use of inhalers. *J Aerosol Med* 2006;19:127-36.
194. Datta D, Vitale A, Lahiri B, ZuWallack R. An evaluation of nebulized levalbuterol in stable COPD. *Chest* 2003;124:844-9.
195. Calverley PM, Anderson JA, Celli B, et al. Salmeterol and fluticasone propionate and survival in chronic obstructive pulmonary disease. *N Engl J Med* 2007;356:775-89.
196. Boyd G, Morice AH, Pounsford JC, Siebert M, Pelsis N, Crawford C. An evaluation of salmeterol in the treatment of chronic obstructive pulmonary disease (COPD) [published erratum appears in *Eur Respir J* 1997 Jul;10(7):1696]. *Eur Respir J* 1997;10:815-21.
197. Cazzola M, Matera MG, Santangelo G, Vinciguerra A, Rossi F, D'Amato G. Salmeterol and formoterol in partially reversible severe chronic obstructive pulmonary disease: a dose-response study. *Respir Med* 1995;89:357-62.

198. Rossi A, Kristufek P, Levine BE, et al. Comparison of the efficacy, tolerability, and safety of formoterol dry powder and oral, slow-release theophylline in the treatment of COPD. *Chest* 2002;121:1058-69.
199. Ulrik CS. Efficacy of inhaled salmeterol in the management of smokers with chronic obstructive pulmonary disease: a single centre randomised, double blind, placebo controlled, crossover study. *Thorax* 1995;50:750-4.
200. Tashkin DP, Fabbri LM. Long-acting beta-agonists in the management of chronic obstructive pulmonary disease: current and future agents. *Respir Res* 2010;11:149.
201. Donohue JF, Fogarty C, Lotvall J, et al. Once-daily bronchodilators for chronic obstructive pulmonary disease: indacaterol versus tiotropium. *Am J Respir Crit Care Med* 2010;182:155-62.
202. Kornmann O, Dahl R, Centanni S, et al. Once-daily indacaterol versus twice-daily salmeterol for COPD: a placebo-controlled comparison. *Eur Respir J* 2011;37:273-9.
203. Lipworth BJ, McDevitt DG, Struthers AD. Hypokalemic and ECG sequelae of combined beta-agonist/diuretic therapy. Protection by conventional doses of spironolactone but not triamterene. *Chest* 1990;98:811-5.
204. Uren NG, Davies SW, Jordan SL, Lipkin DP. Inhaled bronchodilators increase maximum oxygen consumption in chronic left ventricular failure. *Eur Heart J* 1993;14:744-50.
205. Khoukaz G, Gross NJ. Effects of salmeterol on arterial blood gases in patients with stable chronic obstructive pulmonary disease. Comparison with albuterol and ipratropium. *Am J Respir Crit Care Med* 1999;160:1028-30.
206. Polverino E, Gomez FP, Manrique H, et al. Gas exchange response to short-acting beta2-agonists in chronic obstructive pulmonary disease severe exacerbations. *Am J Respir Crit Care Med* 2007;176:350-5.
207. Barnes PJ. Bronchodilators: basic pharmacology. In: Calverley PMA, Pride NB, eds. *Chronic obstructive pulmonary disease*. London: *Chapman and Hall*; 1995:391-417.
208. Disse B, Speck GA, Rominger KL, Witek TJ, Jr., Hammer R. Tiotropium (Spiriva): mechanistical considerations and clinical profile in obstructive lung disease. *Life Sci* 1999;64:457-64.
209. van Noord JA, Bantje TA, Eland ME, Korducki L, Cornelissen PJ. A randomised controlled comparison of tiotropium and ipratropium in the treatment of chronic obstructive pulmonary disease. The Dutch Tiotropium Study Group. *Thorax* 2000;55:289-94.
210. Vincken W, van Noord JA, Greefhorst AP, et al. Improved health outcomes in patients with COPD during 1 yr's treatment with tiotropium. *Eur Respir J* 2002;19:209-16.
211. Casaburi R, Mahler DA, Jones PW, et al. A long-term evaluation of once-daily inhaled tiotropium in chronic obstructive pulmonary disease. *Eur Respir J* 2002;19:217-24.
212. Barr RG, Bourbeau J, Camargo CA, Ram FS. Inhaled tiotropium for stable chronic obstructive pulmonary disease. *Cochrane database of systematic reviews* 2005:CD002876.
213. Kesten S, Casaburi R, Kukafka D, Cooper CB. Improvement in self-reported exercise participation with the combination of tiotropium and rehabilitative exercise training in COPD patients. *Int J Chron Obstruct Pulmon Dis* 2008;3:127-36.
214. Tashkin DP, Celli B, Senn S, et al. A 4-year trial of tiotropium in chronic obstructive pulmonary disease. *N Engl J Med* 2008;359:1543-54.
215. Vogelmeier C, Hederer B, Glaab T, et al. Tiotropium versus salmeterol for the prevention of exacerbations of COPD. *N Engl J Med* 2011;364:1093-103.
216. Tashkin DP. Long-acting anticholinergic use in chronic obstructive pulmonary disease: efficacy and safety. *Current opinion in pulmonary medicine* 2010;16:97-105.
217. Anthonisen NR, Connett JE, Enright PL, Manfreda J. Hospitalizations and mortality in the Lung Health Study. *Am J Respir Crit Care Med* 2002;166:333-9.
218. Michele TM, Pinheiro S, Iyasu S. The safety of tiotropium--the FDA's conclusions. *N Engl J Med* 2010;363:1097-9.
219. Singh S, Loke YK, Enright PL, Furberg CD. Mortality associated with tiotropium mist inhaler in patients with chronic obstructive pulmonary disease: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2011;342:d3215.

220. Aubier M. Pharmacotherapy of respiratory muscles. *Clin Chest Med* 1988;9:311-24.
221. McKay SE, Howie CA, Thomson AH, Whiting B, Addis GJ. Value of theophylline treatment in patients handicapped by chronic obstructive lung disease. *Thorax* 1993;48:227-32.
222. Moxham J. Aminophylline and the respiratory muscles: an alternative view. *Clin Chest Med* 1988;9:325-36.
223. Murciano D, Auclair MH, Pariente R, Aubier M. A randomized, controlled trial of theophylline in patients with severe chronic obstructive pulmonary disease. *N Engl J Med* 1989;320:1521-5.
224. Taylor DR, Buick B, Kinney C, Lowry R, McDevitt DG. The efficacy of orally administered theophylline, inhaled salbutamol, and a combination of the two as chronic therapy in the management of chronic bronchitis with reversible air-flow obstruction. *Am Rev Respir Dis* 1985;131:747-51.
225. Rossi A, Gottfried SB, Higgs BD, Zocchi L, Grassino A, Milic-Emili J. Respiratory mechanics in mechanically ventilated patients with respiratory failure. *J Appl Physiol* 1985;58:1849-58.
226. Ram FS, Jones PW, Castro AA, et al. Oral theophylline for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2002;4:CD003902.
227. Zhou Y, Wang X, Zeng X, et al. Positive benefits of theophylline in a randomized, double-blind, parallel-group, placebo-controlled study of low-dose, slow-release theophylline in the treatment of COPD for 1 year. *Respirology* 2006;11:603-10.
228. ZuWallack RL, Mahler DA, Reilly D, et al. Salmeterol plus theophylline combination therapy in the treatment of COPD. *Chest* 2001;119:1661-70.
229. Ram FS. Use of theophylline in chronic obstructive pulmonary disease: examining the evidence. *Curr Opin Pulm Med* 2006;12:132-9.
230. Vogelmeier C, Kardos P, Harari S, Gans SJ, Stenglein S, Thirlwell J. Formoterol mono- and combination therapy with tiotropium in patients with COPD: a 6-month study. *Respir Med* 2008;102:1511-20.
231. The COMBIVENT Inhalation Solution Study Group. Routine nebulized ipratropium and albuterol together are better than either alone in COPD. *Chest* 1997;112:1514-21.
232. Gross N, Tashkin D, Miller R, Oren J, Coleman W, Linberg S. Inhalation by nebulization of albuterol-ipratropium combination (Dey combination) is superior to either agent alone in the treatment of chronic obstructive pulmonary disease. Dey Combination Solution Study Group. *Respiration* 1998;65:354-62.
233. Bellia V, Foresi A, Bianco S, et al. Efficacy and safety of oxitropium bromide, theophylline and their combination in COPD patients: a double-blind, randomized, multicentre study (BREATH Trial). *Respir Med* 2002;96:881-9.
234. Rabe KF, Timmer W, Sakkrotis A, Viel K. Comparison of a combination of tiotropium plus formoterol to salmeterol plus fluticasone in moderate COPD. *Chest* 2008;134:255-62.
235. van Noord JA, de Munck DR, Bantje TA, Hop WC, Akveld ML, Bommer AM. Long-term treatment of chronic obstructive pulmonary disease with salmeterol and the additive effect of ipratropium. *Eur Respir J* 2000;15:878-85.
236. Guyatt GH, Berman LB, Townsend M, Pugsley SO, Chambers LW. A measure of quality of life for clinical trials in chronic lung disease. *Thorax* 1987;42:773-8.
237. Tashkin DP, Pearle J, Iezzoni D, Varghese ST. Formoterol and tiotropium compared with tiotropium alone for treatment of COPD. *COPD* 2009;6:17-25.
238. van Noord JA, Aumann JL, Janssens E, et al. Comparison of tiotropium once daily, formoterol twice daily and both combined once daily in patients with COPD. *Eur Respir J* 2005;26:214-22.
239. Global Strategy for Asthma Management and Prevention, Revised 2014. Available from <http://www.ginasthma.org>.
240. Calverley P, Pauwels R, Vestbo J, et al. Combined salmeterol and fluticasone in the treatment of chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet* 2003;361:449-56.
241. Calverley PM, Spencer S, Willits L, Burge PS, Jones PW. Withdrawal from treatment as an outcome in the ISOLDE study of COPD. *Chest* 2003;124:1350-6.

242. Jones PW, Willits LR, Burge PS, Calverley PM. Disease severity and the effect of fluticasone propionate on chronic obstructive pulmonary disease exacerbations. *Eur Respir J* 2003;21:68-73.
243. Mahler DA, Wire P, Horstman D, et al. Effectiveness of fluticasone propionate and salmeterol combination delivered via the Diskus device in the treatment of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2002;166(8):1084-91.
244. Szafranski W, Cukier A, Ramirez A, et al. Efficacy and safety of budesonide/formoterol in the management of chronic obstructive pulmonary disease. *Eur Respir J* 2003;21:74-81.
245. van der Valk P, Monninkhof E, van der Palen J, Zielhuis G, van Herwaarden C. Effect of discontinuation of inhaled corticosteroids in patients with chronic obstructive pulmonary disease: the COPE study. *Am J Respir Crit Care Med* 2002;166:1358-63.
246. Drummond MB, Dasenbrook EC, Pitz MW, Murphy DJ, Fan E. Inhaled corticosteroids in patients with stable chronic obstructive pulmonary disease: a systematic review and meta-analysis. *JAMA* 2008;300:2407-16.
247. Singh S, Amin AV, Loke YK. Long-term use of inhaled corticosteroids and the risk of pneumonia in chronic obstructive pulmonary disease: a meta-analysis. *Arch Intern Med* 2009;169:219-29.
248. Calverley PM, Stockley RA, Seemungal TA, et al. Reported pneumonia in patients with COPD: findings from the INSPIRE study. *Chest* 2011;139:506-12.
249. Johnell O, Pauwels R, Lofdahl CG, et al. Bone mineral density in patients with chronic obstructive pulmonary disease treated with budesonide Turbuhaler. *Eur Respir J* 2002;19:1058-63.
250. Ferguson GT, Calverley PM, Anderson JA, et al. Prevalence and progression of osteoporosis in patients with COPD: results from the TOWARDS a Revolution in COPD Health study. *Chest* 2009;136:1456-65.
251. Calverley PM, Boonsawat W, Cseke Z, Zhong N, Peterson S, Olsson H. Maintenance therapy with budesonide and formoterol in chronic obstructive pulmonary disease. *Eur Respir J* 2003;22:912-9.
252. Hanania NA, Darken P, Horstman D, et al. The efficacy and safety of fluticasone propionate (250 microg)/salmeterol (50 microg) combined in the Diskus inhaler for the treatment of COPD. *Chest* 2003;124:834-43.
253. Mahler DA. Pulmonary rehabilitation. *Chest* 1998;113:263S-8S.
254. Nannini LJ, Cates CJ, Lasserson TJ, Poole P. Combined corticosteroid and long-acting beta-agonist in one inhaler versus placebo for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2007; Issue 4. Art. No.: CD003794.
255. Crim C, Calverley PM, Anderson JA, et al. Pneumonia risk in COPD patients receiving inhaled corticosteroids alone or in combination: TORCH study results. *Eur Respir J* 2009;34:641-7.
256. Welte T, Miravittles M, Hernandez P, et al. Efficacy and tolerability of budesonide/formoterol added to tiotropium in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2009;180:741-50.
257. Aaron SD, Vandemheen KL, Fergusson D, et al. Tiotropium in combination with placebo, salmeterol, or fluticasone-salmeterol for treatment of chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 2007;146:545-55.
258. Karner C, Cates CJ. Combination inhaled steroid and long-acting beta(2)-agonist in addition to tiotropium versus tiotropium or combination alone for chronic obstructive pulmonary disease. *Cochrane database of systematic reviews* 2011;3:CD008532.
259. Decramer M, de Bock V, Dom R. Functional and histologic picture of steroid-induced myopathy in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1996;153:1958-64.
260. Decramer M, Lacquet LM, Fagard R, Rogiers P. Corticosteroids contribute to muscle weakness in chronic airflow obstruction. *Am J Respir Crit Care Med* 1994;150:11-6.
261. Decramer M, Stas KJ. Corticosteroid-induced myopathy involving respiratory muscles in patients with chronic obstructive pulmonary disease or asthma. *Am Rev Respir Dis* 1992;146:800-2.

262. Renkema TE, Schouten JP, Koeter GH, Postma DS. Effects of long-term treatment with corticosteroids in COPD. *Chest* 1996;109:1156-62.
263. Rice KL, Rubins JB, Lebahn F, et al. Withdrawal of chronic systemic corticosteroids in patients with COPD: a randomized trial. *Am J Respir Crit Care Med* 2000;162:174-8.
264. Fabbri LM, Calverley PM, Izquierdo-Alonso JL, et al. Roflumilast in moderate-to-severe chronic obstructive pulmonary disease treated with longacting bronchodilators: two randomised clinical trials. *Lancet* 2009;374:695-703.
265. Rabe KF. Update on roflumilast, a phosphodiesterase 4 inhibitor for the treatment of chronic obstructive pulmonary disease. *Br J Pharmacol* 2011;163:53-67.
266. Calverley PM, Rabe KF, Goehring UM, Kristiansen S, Fabbri LM, Martinez FJ. Roflumilast in symptomatic chronic obstructive pulmonary disease: two randomised clinical trials. *Lancet* 2009;374:685-94.
267. Wongsurakiat P, Maranetra KN, Wasi C, Kositanont U, Dejsomritrutai W, Charoenratanakul S. Acute respiratory illness in patients with COPD and the effectiveness of influenza vaccination: a randomized controlled study. *Chest* 2004;125:2011-20.
268. Nichol KL, Margolis KL, Wuorenma J, Von Sternberg T. The efficacy and cost effectiveness of vaccination against influenza among elderly persons living in the community. *N Engl J Med* 1994;331:778-84.
269. Wongsurakiat P, Lertakyamane J, Maranetra KN, Jongriratanakul S, Sangkaew S. Economic evaluation of influenza vaccination in Thai chronic obstructive pulmonary disease patients. *J Med Assoc Thai* 2003;86:497-508.
270. Centers for Disease Control and Prevention. Prevention and control of seasonal influenza with vaccines. Recommendations of the Advisory Committee on Immunization Practices (ACIP) *MMWR Morb Mortal Wkly Rep* 2009;58 (RR08):1-52.
271. Edwards KM, Dupont WD, Westrich MK, Plummer WD, Jr., Palmer PS, Wright PF. A randomized controlled trial of cold-adapted and inactivated vaccines for the prevention of influenza A disease. *J Infect Dis* 1994;169:68-76.
272. Hak E, van Essen GA, Buskens E, Stalman W, de Melker RA. Is immunising all patients with chronic lung disease in the community against influenza cost effective? Evidence from a general practice based clinical prospective cohort study in Utrecht, The Netherlands. *J Epidemiol Community Health* 1998;52:120-5.
273. Woodhead M, Blasi F, Ewig S, et al. Guidelines for the management of adult lower respiratory tract infections. *Eur Respir J* 2005;26:1138-80.
274. Jackson LA, Neuzil KM, Yu O, et al. Effectiveness of pneumococcal polysaccharide vaccine in older adults. *N Engl J Med* 2003;348:1747-55.
275. Prevention of Pneumococcal Disease: Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 1997;46 (RR-08):1-24 <http://www.cdc.gov/mmwr/preview/mmwrhtml/00047135.htm>.
276. Centers for Disease Control and Prevention. Recommended adult immunization schedule. United States, 2010. *MMWR Morb Mortal Wkly Rep* 2011;60:1-4.
277. Alfageme I, Vazquez R, Reyes N, et al. Clinical efficacy of anti-pneumococcal vaccination in patients with COPD. *Thorax* 2006;61:189-95.
278. Francis RS, May JR, Spicer CC. Chemotherapy of bronchitis: influence of penicillin and tetracycline administered daily, or intermittently for exacerbations. *BMJ* 1961;2:979-85.
279. Francis RS, Spicer CC. Chemotherapy in chronic bronchitis: influence of daily penicillin and tetracycline on exacerbations and their cost. A report to the research committee of the British Tuberculosis Association by their Chronic Bronchitis subcommittee. *BMJ* 1960;1:297-303.
280. Fletcher CM, Ball JD, Carstairs LW, et al. Value of chemoprophylaxis and chemotherapy in early chronic bronchitis. A report to the Medical Research Council by their Working Party on trials of chemotherapy in early chronic bronchitis. In: *BMJ*; 1966:1317-22.
281. Johnston RN, McNeill RS, Smith DH, et al. Five-year winter chemoprophylaxis for chronic bronchitis. *Br Med J* 1969;4:265-9.

282. Seemungal TA, Wilkinson TM, Hurst JR, Perera WR, Sapsford RJ, Wedzicha JA. Long-term erythromycin therapy is associated with decreased chronic obstructive pulmonary disease exacerbations. *Am J Respir Crit Care Med* 2008;178:1139-47.
283. Sethi S, Jones PW, Theron MS, et al. Pulsed moxifloxacin for the prevention of exacerbations of chronic obstructive pulmonary disease: a randomized controlled trial. *Respiratory research* 2010;11:10.
284. Albert RK, Connett J, Bailey WC, et al. Azithromycin for prevention of exacerbations of COPD. *N Engl J Med* 2011;365:689-98.
285. Isada CM, Stoller JK. Chronic bronchitis: the role of antibiotics. In: Niederman MS, Sarosi GA, Glassroth J, eds. *Respiratory infections: a scientific basis for management*. London: WB Saunders; 1994:621-33.
286. Siafakas NM, Celli BR. Overall management of stable chronic obstructive Pulmonary disease. In: *Management of Chronic Obstructive Pulmonary Disease* Edited by N.M. Siafakas. *Eur Respir Mon* 2006; 38: 258-265.
287. Allegra L, Cordaro CI, Grassi C. Prevention of acute exacerbations of chronic obstructive bronchitis with carbocysteine lysine salt monohydrate: a multicenter, double-blind, placebo-controlled trial. *Respiration* 1996;63:174-80.
288. Guyatt GH, Townsend M, Kazim F, Newhouse MT. A controlled trial of ambroxol in chronic bronchitis. *Chest* 1987;92:618-20.
289. Petty TL. The National Mucolytic Study. Results of a randomized, double-blind, placebo-controlled study of iodinated glycerol in chronic obstructive bronchitis. *Chest* 1990;97:75-83.
290. National Institute for Clinical Excellence (NICE). Chronic obstructive pulmonary disease. Management of chronic obstructive pulmonary disease in adults in primary and secondary care. <http://guidancencice.org.uk/CG101/Guidance/pdf/English> 2010.
291. Celli BR, MacNee W. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J* 2004;23:932-46.
292. Hansen NC, Skriver A, Brorsen-Riis L, et al. Orally administered N-acetylcysteine may improve general well-being in patients with mild chronic bronchitis. *Respir Med* 1994;88:531-5.
293. British Thoracic Society Research Committee. Oral N-acetylcysteine and exacerbation rates in patients with chronic bronchitis and severe airways obstruction. *Thorax* 1985;40:832-5.
294. Boman G, Backer U, Larsson S, Melander B, Wahlander L. Oral acetylcysteine reduces exacerbation rate in chronic bronchitis: report of a trial organized by the Swedish Society for Pulmonary Diseases. *Eur J Respir Dis* 1983;64:405-15.
295. Rasmussen JB, Glenmow C. Reduction in days of illness after long-term treatment with N-acetylcysteine controlled-release tablets in patients with chronic bronchitis. *Eur Respir J* 1988;1:351-5.
296. Zheng JP, Kang J, Huang SG, et al. Effect of carbocysteine on acute exacerbation of chronic obstructive pulmonary disease (PEACE Study): a randomised placebo-controlled study. *Lancet* 2008;371:2013-8.
297. Decramer M, Rutten-van Molken M, Dekhuijzen PN, et al. Effects of N-acetylcysteine on outcomes in chronic obstructive pulmonary disease (Bronchitis Randomized on NAC Cost-Utility Study, BRONCUS): a randomised placebo-controlled trial. *Lancet* 2005;365:1552-60.
298. Collet JP, Shapiro P, Ernst P, Renzi T, Ducruet T, Robinson A. Effects of an immunostimulating agent on acute exacerbations and hospitalizations in patients with chronic obstructive pulmonary disease. The PARI-IS Study Steering Committee and Research Group. Prevention of Acute Respiratory Infection by an Immunostimulant. *Am J Respir Crit Care Med* 1997;156:1719-24.
299. Li J, Zheng JP, Yuan JP, Zeng GQ, Zhong NS, Lin CY. Protective effect of a bacterial extract against acute exacerbation in patients with chronic bronchitis accompanied by chronic obstructive pulmonary disease. *Chin Med J (Engl)* 2004;117:828-34.
300. Anthonisen NR. OM-8BV for COPD. *Am J Respir Crit Care Med* 1997;156:1713-4.

301. Irwin RS, Boulet LP, Cloutier MM, et al. Managing cough as a defense mechanism and as a symptom. A consensus panel report of the American College of Chest Physicians. *Chest* 1998;114:133S-81S.
302. Barbera JA, Roger N, Roca J, Rovira I, Higenbottam TW, Rodriguez-Roisin R. Worsening of pulmonary gas exchange with nitric oxide inhalation in chronic obstructive pulmonary disease. *Lancet* 1996;347:436-40.
303. Jones AT, Evans TW. NO: COPD and beyond. *Thorax* 1997;52 Suppl 3:S16-21.
304. Galie N, Hoepfer MM, Humbert M, et al. Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eur Respir J* 2009;34:1219-63.
305. Jennings AL, Davies AN, Higgins JP, Gibbs JS, Broadley KE. A systematic review of the use of opioids in the management of dyspnoea. *Thorax* 2002;57:939-44.
306. Eiser N, Denman WT, West C, Luce P. Oral diamorphine: lack of effect on dyspnoea and exercise tolerance in the "pink puffer" syndrome. *Eur Respir J* 1991;4:926-31.
307. Young IH, Daviskas E, Keena VA. Effect of low dose nebulised morphine on exercise endurance in patients with chronic lung disease. *Thorax* 1989;44:387-90.
308. Woodcock AA, Gross ER, Gellert A, Shah S, Johnson M, Geddes DM. Effects of dihydrocodeine, alcohol, and caffeine on breathlessness and exercise tolerance in patients with chronic obstructive lung disease and normal blood gases. *N Engl J Med* 1981;305:1611-6.
309. Rice KL, Kronenberg RS, Hedemark LL, Niewoehner DE. Effects of chronic administration of codeine and promethazine on breathlessness and exercise tolerance in patients with chronic airflow obstruction. *Br J Dis Chest* 1987;81:287-92.
310. Poole PJ, Veale AG, Black PN. The effect of sustained-release morphine on breathlessness and quality of life in severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;157:1877-80.
311. Rennard SI, Fogarty C, Kelsen S, et al. The safety and efficacy of infliximab in moderate to severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2007;175:926-34.
312. Guo R, Pittler MH, Ernst E. Herbal medicines for the treatment of COPD: a systematic review. *Eur Respir J* 2006;28:330-8.
313. Nici L, Donner C, Wouters E, et al. American Thoracic Society/European Respiratory Society statement on pulmonary rehabilitation. *Am J Respir Crit Care Med* 2006;173:1390-413.
314. Ries AL, Bauldoff GS, Carlin BW, et al. Pulmonary Rehabilitation: Joint ACCP/AACVPR Evidence-Based Clinical Practice Guidelines. *Chest* 2007;131:4S-42S.
315. Lacasse Y, Brosseau L, Milne S, et al. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2002:CD003793.
316. Maltais F, Bourbeau J, Shapiro S, et al. Effects of home-based pulmonary rehabilitation in patients with chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 2008;149:869-78.
317. Casaburi R, Kukafka D, Cooper CB, Witek TJ, Jr., Kesten S. Improvement in exercise tolerance with the combination of tiotropium and pulmonary rehabilitation in patients with COPD. *Chest* 2005;127:809-17.
318. Lacasse Y, Wong E, Guyatt GH, King D, Cook DJ, Goldstein RS. Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. *Lancet* 1996;348:1115-9.
319. Puhan M, Scharplatz M, Troosters T, Walters EH, Steurer J. Pulmonary rehabilitation following exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2009:CD005305.
320. Lacasse Y, Goldstein R, Lasserson TJ, Martin S. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2006:CD003793.
321. Behnke M, Taube C, Kirsten D, Lehnigk B, Jorres RA, Magnussen H. Home-based exercise is capable of preserving hospital-based improvements in severe chronic obstructive pulmonary disease. *Respir Med* 2000;94:1184-91.
322. Finnerty JP, Keeping I, Bullough I, Jones J. The effectiveness of outpatient pulmonary rehabilitation in chronic lung disease: a randomized controlled trial. *Chest* 2001;119:1705-10.

323. Green RH, Singh SJ, Williams J, Morgan MD. A randomised controlled trial of four weeks versus seven weeks of pulmonary rehabilitation in chronic obstructive pulmonary disease. *Thorax* 2001;56:143-5.
324. Ries AL, Kaplan RM, Myers R, Prewitt LM. Maintenance after pulmonary rehabilitation in chronic lung disease: a randomized trial. *Am J Respir Crit Care Med* 2003;167:880-8.
325. Esteban C, Quintana JM, Aburto M, et al. Impact of changes in physical activity on health-related quality of life among patients with COPD. *Eur Respir J* 2010;36:292-300
326. Garcia-Aymerich J, Lange P, Benet M, Schnohr P, Anto JM. Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort study. *Thorax* 2006;61:772-8.
327. Troosters T, Casaburi R, Gosselink R, Decramer M. Pulmonary rehabilitation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2005;172:19-38.
328. Puhan MA, Busching G, Schunemann HJ, VanOort E, Zaugg C, Frey M. Interval versus continuous high-intensity exercise in chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 2006;145:816-25.
329. Vogiatzis I, Nanas S, Roussos C. Interval training as an alternative modality to continuous exercise in patients with COPD. *Eur Respir J* 2002;20:12-9.
330. Honeyman P, Barr P, Stubbing DG. Effect of a walking aid on disability, oxygenation, and breathlessness in patients with chronic airflow limitation. *J Cardiopulm Rehabil* 1996;16:63-7.
331. Roomi J, Yohannes AM, Connolly MJ. The effect of walking aids on exercise capacity and oxygenation in elderly patients with chronic obstructive pulmonary disease. *Age Ageing* 1998;27:703-6.
332. Yohannes AM, Connolly MJ. Early mobilization with walking aids following hospital admission with acute exacerbation of chronic obstructive pulmonary disease. *Clin Rehabil* 2003;17:465-71.
333. Emtner M, Porszasz J, Burns M, Somfay A, Casaburi R. Benefits of supplemental oxygen in exercise training in nonhypoxemic chronic obstructive pulmonary disease patients. *Am J Respir Crit Care Med* 2003;168:1034-42.
334. Palange P, Valli G, Onorati P, et al. Effect of heliox on lung dynamic hyperinflation, dyspnea, and exercise endurance capacity in COPD patients. *J Appl Physiol* 2004;97:1637-42.
335. Belman MJ, Botnick WC, Nathan SD, Chon KH. Ventilatory load characteristics during ventilatory muscle training. *Am J Respir Crit Care Med* 1994;149:925-9.
336. Lotters F, van Tol B, Kwakkel G, Gosselink R. Effects of controlled inspiratory muscle training in patients with COPD: a meta-analysis. *Eur Respir J* 2002;20:570-6.
337. Magadle R, McConnell AK, Beckerman M, Weiner P. Inspiratory muscle training in pulmonary rehabilitation program in COPD patients. *Respir Med* 2007;101:1500-5.
338. O'Brien K, Geddes BL, Reid WD, Brooks D, Crowe J. Inspiratory muscle training compared with other rehabilitation interventions in chronic obstructive pulmonary disease: a systematic review update. *J Cardiopulm Rehabil Prev* 2008;28:128-41.
339. Bernard S, Whittom F, Leblanc P, et al. Aerobic and strength training in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:896-901.
340. Wedzicha JA, Bestall JC, Garrod R, Garnham R, Paul EA, Jones PW. Randomized controlled trial of pulmonary rehabilitation in severe chronic obstructive pulmonary disease patients, stratified with the MRC dyspnoea scale. *Eur Respir J* 1998;12:363-9.
341. Young P, Dewse M, Fergusson W, Kolbe J. Improvements in outcomes for chronic obstructive pulmonary disease (COPD) attributable to a hospital-based respiratory rehabilitation programme. *Aust N Z J Med* 1999;29:59-65.
342. Ashikaga T, Vacek PM, Lewis SO. Evaluation of a community-based education program for individuals with chronic obstructive pulmonary disease. *J Rehabil* 1980;46:23-7.
343. Janelli LM, Scherer YK, Schmieder LE. Can a pulmonary health teaching program alter patients' ability to cope with COPD? *Rehabil Nurs* 1991;16:199-202.
344. Reis AL. Response to bronchodilators. In: Clausen J, ed. *Pulmonary function testing: guidelines and controversies*. New York: Academic Press; 1982.

345. Toshima MT, Kaplan RM, Ries AL. Experimental evaluation of rehabilitation in chronic obstructive pulmonary disease: short-term effects on exercise endurance and health status. *Health Psychol* 1990;9:237-52.
346. Celli BR. Pulmonary rehabilitation in patients with COPD. *Am J Respir Crit Care Med* 1995;152:861-4.
347. Jones PW, Quirk FH, Baveystock CM, Littlejohns P. A self-complete measure for chronic airflow limitation - the St George's Respiratory Questionnaire. *Amer Rev Respir Dis* 1992;145:1321-7
348. Ware JE, Jr., Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473-83.
349. Dowson C, Laing R, Barraclough R, et al. The use of the Hospital Anxiety and Depression Scale (HADS) in patients with chronic obstructive pulmonary disease: a pilot study. *N Z Med J* 2001;114:447-9.
350. Kunik ME, Veazey C, Cully JA, et al. COPD education and cognitive behavioral therapy group treatment for clinically significant symptoms of depression and anxiety in COPD patients: a randomized controlled trial. *Psychol Med* 2008;38:385-96.
351. Anthonisen NR, Manfreda J, Warren CP, Hershfield ES, Harding GK, Nelson NA. Antibiotic therapy in exacerbations of chronic obstructive pulmonary disease. *Ann Intern Med* 1987;106:196-204.
352. Reference deleted
353. Reference deleted
354. Reference deleted
355. Reference deleted
356. Reference deleted
357. Reference deleted
358. Reference deleted
359. Stoller JK, Panos RJ, Krachman S, Doherty DE, Make B. Oxygen therapy for patients with COPD: current evidence and the long-term oxygen treatment trial. *Chest* 2010;138:179-87.
360. Moore RP, Berlowitz DJ, Denehy L, et al. A randomised trial of domiciliary, ambulatory oxygen in patients with COPD and dyspnoea but without resting hypoxaemia. *Thorax* 2011;66:32-7.
361. Berg BW, Dillard TA, Rajagopal KR, Mehm WJ. Oxygen supplementation during air travel in patients with chronic obstructive lung disease. *Chest* 1992;101:638-41.
362. Gong H, Jr. Air travel and oxygen therapy in cardiopulmonary patients. *Chest* 1992;101:1104-13.
363. Gong H, Jr., Tashkin DP, Lee EY, Simmons MS. Hypoxia-altitude simulation test. Evaluation of patients with chronic airway obstruction. *Am Rev Respir Dis* 1984;130:980-6.
364. Christensen CC, Ryg M, Refvem OK, Skjonsberg OH. Development of severe hypoxaemia in chronic obstructive pulmonary disease patients at 2,438 m (8,000 ft) altitude. *Eur Respir J* 2000;15:635-9.
365. McEvoy RD, Pierce RJ, Hillman D, et al. Nocturnal non-invasive nasal ventilation in stable hypercapnic COPD: a randomised controlled trial. *Thorax* 2009;64:561-6.
366. Marin JM, Soriano JB, Carrizo SJ, Boldova A, Celli BR. Outcomes in patients with chronic obstructive pulmonary disease and obstructive sleep apnea: the overlap syndrome. *Am J Respir Crit Care Med* 2010;182:325-31.
367. Cooper JD, Trulock EP, Triantafillou AN, et al. Bilateral pneumectomy (volume reduction) for chronic obstructive pulmonary disease. *J Thorac Cardiovasc Surg* 1995;109:106-16.
368. Criner G, Cordova FC, Leyenson V, et al. Effect of lung volume reduction surgery on diaphragm strength. *Am J Respir Crit Care Med* 1998;157:1578-85.
369. Martinez FJ, de Oca MM, Whyte RI, Stetz J, Gay SE, Celli BR. Lung-volume reduction improves dyspnea, dynamic hyperinflation, and respiratory muscle function. *Am J Respir Crit Care Med* 1997;155:1984-90.
370. Fessler HE, Permutt S. Lung volume reduction surgery and airflow limitation. *Am J Respir Crit Care Med* 1998;157:715-22.
371. Ramsey SD, Shroyer AL, Sullivan SD, Wood DE. Updated evaluation of the cost-effectiveness of lung volume reduction surgery. *Chest* 2007;131:823-32.

372. Naunheim KS, Wood DE, Mohsenifar Z, et al. Long-term follow-up of patients receiving lung-volume-reduction surgery versus medical therapy for severe emphysema by the National Emphysema Treatment Trial Research Group. *Ann Thorac Surg* 2006;82:431-43.
373. National Emphysema Treatment Trial Research Group. Patients at high risk of death after lung-volume-reduction surgery. *N Engl J Med* 2001;345:1075-83.
374. Sciruba FC, Ernst A, Herth FJ, et al. A randomized study of endobronchial valves for advanced emphysema. *N Engl J Med* 2010;363:1233-44.
375. Christie JD, Edwards LB, Kucheryavaya AY, et al. The Registry of the International Society for Heart and Lung Transplantation: twenty-seventh official adult lung and heart-lung transplant report--2010. *J Heart Lung Transplant* 2010;29:1104-18.
376. Trulock EP. Lung transplantation. *Am J Respir Crit Care Med* 1997;155:789-818.
377. Theodore J, Lewiston N. Lung transplantation comes of age. *N Engl J Med* 1990;322:772-4.
378. Orens JB, Estenne M, Arcasoy S, et al. International guidelines for the selection of lung transplant candidates. 2006 update--a consensus report from the Pulmonary Scientific Council of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant* 2006;25:745-55.
379. Romieu I, Riojas-Rodriguez H, Marron-Mares AT, Schilman A, Perez-Padilla R, Maseria O. Improved biomass stove intervention in rural Mexico: impact on the respiratory health of women. *Am J Respir Crit Care Med* 2009;180:649-56.
380. Liu S, Zhou Y, Wang X, et al. Biomass fuels are the probable risk factor for chronic obstructive pulmonary disease in rural South China. *Thorax* 2007;62:889-97.
381. Calverley PM, Burge PS, Spencer S, Anderson JA, Jones PW. Bronchodilator reversibility testing in chronic obstructive pulmonary disease. *Thorax* 2003;58:659-64.
382. Tashkin D, Kesten S. Long-term treatment benefits with tiotropium in COPD patients with and without short-term bronchodilator responses. *Chest* 2003;123:1441-9.
383. Berry MJ, Rejeski WJ, Adair NE, Zaccaro D. Exercise rehabilitation and chronic obstructive pulmonary disease stage. *Am J Respir Crit Care Med* 1999;160:1248-53.
384. Foglio K, Bianchi L, Bruletti G, Battista L, Pagani M, Ambrosino N. Long-term effectiveness of pulmonary rehabilitation in patients with chronic airway obstruction. *Eur Respir J* 1999;13:125-32.
385. Griffiths TL, Burr ML, Campbell IA, et al. Results at 1 year of outpatient multidisciplinary pulmonary rehabilitation: a randomised controlled trial [published erratum appears in *Lancet* 2000;355:1280]. *Lancet* 2000;355:362-8.
386. Friedman M, Serby CW, Menjoge SS, Wilson JD, Hilleman DE, Witek TJ, Jr. Pharmacoeconomic evaluation of a combination of ipratropium plus albuterol compared with ipratropium alone and albuterol alone in COPD. *Chest* 1999;115:635-41.
387. Appleton S, Poole P, Smith B, Veale A, Lasserson TJ, Chan MM. Long-acting beta2-agonists for poorly reversible chronic obstructive pulmonary disease. *Cochrane database of systematic reviews* 2006;3:CD001104.
388. Niewoehner DE, Rice K, Cote C, et al. Prevention of exacerbations of chronic obstructive pulmonary disease with tiotropium, a once-daily inhaled anticholinergic bronchodilator: a randomized trial. *Ann Intern Med* 2005;143:317-26.
389. Wedzicha JA, Calverley PM, Seemungal TA, Hagan G, Ansari Z, Stockley RA. The prevention of chronic obstructive pulmonary disease exacerbations by salmeterol/fluticasone propionate or tiotropium bromide. *Am J Respir Crit Care Med* 2008;177:19-26.
390. Reference deleted
391. Kessler R, Stahl E, Vogelmeier C, et al. Patient understanding, detection, and experience of COPD exacerbations: an observational, interview-based study. *Chest* 2006;130:133-42.
392. Mazzone PJ. Preoperative evaluation of the lung cancer resection candidate. *Expert Rev Respir Med* 2010;4:97-113.
393. Schuurmans MM, Diacon AH, Bolliger CT. Functional evaluation before lung resection. *Clin Chest Med* 2002;23:159-72.

394. Smetana GW. Preoperative pulmonary evaluation. *N Engl J Med* 1999;340:937-44.
395. Trayner E, Jr., Celli BR. Postoperative pulmonary complications. *Med Clin North Am* 2001;85:1129-39.
396. Brunelli A, Charloux A, Bolliger CT, et al. ERS/ESTS clinical guidelines on fitness for radical therapy in lung cancer patients (surgery and chemo-radiotherapy). *Eur Respir J* 2009;34:17-41.
397. Colice GL, Shafazand S, Griffin JP, Keenan R, Bolliger CT. Physiologic evaluation of the patient with lung cancer being considered for resectional surgery: ACCP evidenced-based clinical practice guidelines (2nd edition). *Chest* 2007;132:161S-77S.
398. Seemungal TA, Donaldson GC, Bhowmik A, Jeffries DJ, Wedzicha JA. Time course and recovery of exacerbations in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2000;161:1608-13.
399. Donaldson GC, Seemungal TA, Bhowmik A, Wedzicha JA. Relationship between exacerbation frequency and lung function decline in chronic obstructive pulmonary disease. *Thorax* 2002;57:847-52.
400. Kanner RE, Anthonisen NR, Connett JE. Lower respiratory illnesses promote FEV(1) decline in current smokers but not ex-smokers with mild chronic obstructive pulmonary disease: results from the lung health study. *Am J Respir Crit Care Med* 2001;164:358-64.
401. Wouters EF. The burden of COPD in The Netherlands: results from the Confronting COPD survey. *Respir Med* 2003;97 Suppl C:S51-9.
402. Connors AF, Jr., Dawson NV, Thomas C, et al. Outcomes following acute exacerbation of severe chronic obstructive lung disease. The SUPPORT investigators (Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments). *Am J Respir Crit Care Med* 1996;154:959-67.
403. Gunen H, Hacievliyagil SS, Kosar F, et al. Factors affecting survival of hospitalised patients with COPD. *Eur Respir J* 2005;26:234-41.
404. Kong GK, Belman MJ, Weingarten S. Reducing length of stay for patients hospitalized with exacerbation of COPD by using a practice guideline. *Chest* 1997;111:89-94.
405. Seneff MG, Wagner DP, Wagner RP, Zimmerman JE, Knaus WA. Hospital and 1-year survival of patients admitted to intensive care units with acute exacerbation of chronic obstructive pulmonary disease. *JAMA* 1995;274:1852-7.
406. Wilkinson TM, Donaldson GC, Hurst JR, Seemungal TA, Wedzicha JA. Early therapy improves outcomes of exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004;169:1298-303.
407. Monso E, Rosell A, Bonet G, et al. Risk factors for lower airway bacterial colonization in chronic bronchitis. *Eur Respir J* 1999;13:338-42.
408. Pela R, Marchesani F, Agostinelli C, et al. Airways microbial flora in COPD patients in stable clinical conditions and during exacerbations: a bronchoscopic investigation. *Monaldi Arch Chest Dis* 1998;53:262-7.
409. Sethi S, Murphy TF. Infection in the pathogenesis and course of chronic obstructive pulmonary disease. *N Engl J Med* 2008;359:2355-65.
410. Fagon JY, Chastre J, Trouillet JL, et al. Characterization of distal bronchial microflora during acute exacerbation of chronic bronchitis. Use of the protected specimen brush technique in 54 mechanically ventilated patients. *Am Rev Respir Dis* 1990;142:1004-8.
411. Monso E, Ruiz J, Rosell A, et al. Bacterial infection in chronic obstructive pulmonary disease. A study of stable and exacerbated outpatients using the protected specimen brush. *Am J Respir Crit Care Med* 1995;152:1316-20.
412. Soler N, Torres A, Ewig S, et al. Bronchial microbial patterns in severe exacerbations of chronic obstructive pulmonary disease (COPD) requiring mechanical ventilation. *Am J Respir Crit Care Med* 1998;157:1498-505.
413. Sethi S, Wrona C, Grant BJ, Murphy TF. Strain-specific immune response to Haemophilus influenzae in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004;169:448-53.
414. Ling SH, van Eeden SF. Particulate matter air pollution exposure: role in the development and exacerbation of chronic obstructive pulmonary disease. *International journal of chronic obstructive pulmonary disease* 2009;4:233-43.

415. Sint T, Donohue JF, Ghio AJ. Ambient air pollution particles and the acute exacerbation of chronic obstructive pulmonary disease. *Inhalation toxicology* 2008;20:25-9.
416. Peacock JL, Anderson HR, Bremner SA, et al. Outdoor air pollution and respiratory health in patients with COPD. *Thorax* 2011;66:591-6.
417. Adams S, J. M, Luther M. Antibiotics are associated with lower relapse rates in outpatients with acute exacerbations of chronic obstructive pulmonary disease. *Chest* 2000;117:1345-52.
418. Emerman CL, Connors AF, Lukens TW, Effron D, May ME. Relationship between arterial blood gases and spirometry in acute exacerbations of chronic obstructive pulmonary disease. *Ann Emerg Med* 1989;18:523-7.
419. Martinez FJ, Han MK, Flaherty K, Curtis J. Role of infection and antimicrobial therapy in acute exacerbations of chronic obstructive pulmonary disease. *Expert Rev Anti Infect Ther* 2006;4:101-24.
420. Turner MO, Patel A, Ginsburg S, FitzGerald JM. Bronchodilator delivery in acute airflow obstruction. A meta-analysis. *Arch Intern Med* 1997;157:1736-44.
421. Barbera JA, Reyes A, Roca J, Montserrat JM, Wagner PD, Rodriguez-Roisin R. Effect of intravenously administered aminophylline on ventilation/perfusion inequality during recovery from exacerbations of chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1992;145:1328-33.
422. Emerman CL, Connors AF, Lukens TW, May ME, Effron D. Theophylline concentrations in patients with acute exacerbation of COPD. *Am J Emerg Med* 1990;8:289-92.
423. Lloberes P, Ramis L, Montserrat JM, et al. Effect of three different bronchodilators during an exacerbation of chronic obstructive pulmonary disease. *Eur Respir J* 1988;1:536-9.
424. Mahon JL, Laupacis A, Hodder RV, et al. Theophylline for irreversible chronic airflow limitation: a randomized study comparing n of 1 trials to standard practice. *Chest* 1999;115:38-48.
425. Murciano D, Aubier M, Lecocguic Y, Pariente R. Effects of theophylline on diaphragmatic strength and fatigue in patients with chronic obstructive pulmonary disease. *N Engl J Med* 1984;311:349-53.
426. Barr RG, Rowe BH, Camargo CA, Jr. Methylxanthines for exacerbations of chronic obstructive pulmonary disease: meta-analysis of randomised trials. *BMJ* 2003;327:643.
427. Duffy N, Walker P, Diamantea F, Calverley PM, Davies L. Intravenous aminophylline in patients admitted to hospital with non-acidotic exacerbations of chronic obstructive pulmonary disease: a prospective randomised controlled trial. *Thorax* 2005;60:713-7.
428. Davies L, Angus RM, Calverley PM. Oral corticosteroids in patients admitted to hospital with exacerbations of chronic obstructive pulmonary disease: a prospective randomised controlled trial. *Lancet* 1999;354:456-60.
429. Maltais F, Ostinelli J, Bourbeau J, et al. Comparison of nebulized budesonide and oral prednisolone with placebo in the treatment of acute exacerbations of chronic obstructive pulmonary disease: a randomized controlled trial. *Am J Respir Crit Care Med* 2002;165:698-703.
430. Niewehner DE, Erbland ML, Deupree RH, et al. Effect of systemic glucocorticoids on exacerbations of chronic obstructive pulmonary disease. Department of Veterans Affairs Cooperative Study Group. *N Engl J Med* 1999;340:1941-7.
431. Thompson WH, Nielson CP, Carvalho P, Charan NB, Crowley JJ. Controlled trial of oral prednisone in outpatients with acute COPD exacerbation. *Am J Respir Crit Care Med* 1996;154:407-12.
432. Aaron SD, Vandemheen KL, Hebert P, et al. Outpatient oral prednisone after emergency treatment of chronic obstructive pulmonary disease. *N Engl J Med* 2003;348:2618-25.
433. de Jong YP, Uil SM, Grotjohan HP, Postma DS, Kerstjens HA, van den Berg JW. Oral or IV prednisolone in the treatment of COPD exacerbations: a randomized, controlled, double-blind study. *Chest* 2007;132:1741-7.
434. Gunen H, Hacievliyagil SS, Yetkin O, Gulbas G, Mutlu LC, In E. The role of nebulised budesonide in the treatment of exacerbations of COPD. *Eur Respir J* 2007;29:660-7.
435. Stallberg B, Selroos O, Vogelmeier C, Andersson E, Ekstrom T, Larsson K. Budesonide/formoterol as effective as prednisolone plus formoterol in acute exacerbations of COPD. A double-blind, randomised, non-inferiority, parallel-group, multicentre study. *Respir Res* 2009;10:11.

436. Seemungal T, Harper-Owen R, Bhowmik A, et al. Respiratory viruses, symptoms, and inflammatory markers in acute exacerbations and stable chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001;164:1618-23.
437. Ram FS, Rodriguez-Roisin R, Granados-Navarrete A, Garcia-Aymerich J, Barnes NC. Antibiotics for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2006:CD004403.
438. Quon BS, Gan WQ, Sin DD. Contemporary management of acute exacerbations of COPD: a systematic review and metaanalysis. *Chest* 2008;133:756-66.
439. Christ-Crain M, Jaccard-Stolz D, Bingisser R, et al. Effect of procalcitonin-guided treatment on antibiotic use and outcome in lower respiratory tract infections: cluster-randomised, single-blinded intervention trial. *Lancet* 2004;363:600-7.
440. Nouira S, Marghli S, Belghith M, Besbes L, Elatrous S, Abroug F. Once daily oral ofloxacin in chronic obstructive pulmonary disease exacerbation requiring mechanical ventilation: a randomised placebo- controlled trial. *Lancet* 2001;358:2020-5.
441. Miravittles M, Espinosa C, Fernandez-Laso E, Martos JA, Maldonado JA, Gallego M. Relationship between bacterial flora in sputum and functional impairment in patients with acute exacerbations of COPD. Study Group of Bacterial Infection in COPD. *Chest* 1999;116:40-6.
442. Austin MA, Wills KE, Blizzard L, Walters EH, Wood-Baker R. Effect of high flow oxygen on mortality in chronic obstructive pulmonary disease patients in prehospital setting: randomised controlled trial. *BMJ* 2010;341:c5462.
443. Consensus conference report. Clinical indications for noninvasive positive pressure ventilation in chronic respiratory failure due to restrictive lung disease, COPD, and nocturnal hypoventilation. *Chest* 1999;116:521-34.
444. Brochard L, Mancebo J, Wysocki M, et al. Noninvasive ventilation for acute exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* 1995;333:817-22.
445. Lightowler JV, Wedzicha JA, Elliott MW, Ram FS. Non-invasive positive pressure ventilation to treat respiratory failure resulting from exacerbations of chronic obstructive pulmonary disease: Cochrane systematic review and meta-analysis. *BMJ* 2003;326:185.
446. Meyer TJ, Hill NS. Noninvasive positive pressure ventilation to treat respiratory failure. *Ann Intern Med* 1994;120:760-70.
447. Bott J, Carroll MP, Conway JH, et al. Randomised controlled trial of nasal ventilation in acute ventilatory failure due to chronic obstructive airways disease. *Lancet* 1993;341:1555-7.
448. Kramer N, Meyer TJ, Meharg J, Cece RD, Hill NS. Randomized, prospective trial of noninvasive positive pressure ventilation in acute respiratory failure. *Am J Respir Crit Care Med* 1995;151:1799-806.
449. Plant PK, Owen JL, Elliott MW. Early use of non-invasive ventilation for acute exacerbations of chronic obstructive pulmonary disease on general respiratory wards: a multicentre randomised controlled trial. *Lancet* 2000;355:1931-5.
450. Conti G, Antonelli M, Navalesi P, et al. Noninvasive vs. conventional mechanical ventilation in patients with chronic obstructive pulmonary disease after failure of medical treatment in the ward: a randomized trial. *Intensive Care Med* 2002;28:1701-7.
451. International Consensus Conferences in *Intensive Care Medicine*: noninvasive positive pressure ventilation in acute respiratory failure. *Am J Respir Crit Care Med* 2001;163(1):283-91.
452. Esteban A, Anzueto A, Alia I, et al. How is mechanical ventilation employed in the intensive care unit? An international utilization review. *Am J Respir Crit Care Med* 2000;161:1450-8.
453. Esteban A, Anzueto A, Frutos F, et al. Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study. *JAMA* 2002;287:345-55.
454. Wildman MJ, Sanderson C, Groves J, et al. Implications of prognostic pessimism in patients with chronic obstructive pulmonary disease (COPD) or asthma admitted to intensive care in the UK within the COPD and asthma outcome study (CAOS): multicentre observational cohort study. *BMJ* 2007;335:1132.
455. Purro A, Appendini L, De Gaetano A, Gudjonsdottir M, Donner CF, Rossi A. Physiologic determinants of ventilator dependence in long-term mechanically ventilated patients. *Am J Respir Crit Care Med* 2000;161:1115-23.

456. Beydon L, Cinotti L, Rekik N, et al. Changes in the distribution of ventilation and perfusion associated with separation from mechanical ventilation in patients with obstructive pulmonary disease. *Anesthesiology* 1991;75:730-8.
457. Nava S, Ambrosino N, Clini E, et al. Noninvasive mechanical ventilation in the weaning of patients with respiratory failure due to chronic obstructive pulmonary disease. A randomized, controlled trial. *Ann Intern Med* 1998;128:721-8.
458. Torres A, Reyes A, Roca J, Wagner PD, Rodriguez-Roisin R. Ventilation-perfusion mismatching in chronic obstructive pulmonary disease during ventilator weaning. *Am Rev Respir Dis* 1989;140:1246-50.
459. Brochard L, Rauss A, Benito S, et al. Comparison of three methods of gradual withdrawal from ventilatory support during weaning from mechanical ventilation. *Am J Respir Crit Care Med* 1994;150:896-903.
460. Esteban A, Frutos F, Tobin MJ, et al. A comparison of four methods of weaning patients from mechanical ventilation. Spanish Lung Failure Collaborative Group. *N Engl J Med* 1995;332:345-50.
461. Hilbert G, Gruson D, Portel L, Gbikpi-Benissan G, Cardinaud JP. Noninvasive pressure support ventilation in COPD patients with postextubation hypercapnic respiratory insufficiency. *Eur Respir J* 1998;11:1349-53.
462. Ferrer M, Sellares J, Valencia M, et al. Non-invasive ventilation after extubation in hypercapnic patients with chronic respiratory disorders: randomised controlled trial. *Lancet* 2009;374:1082-8.
463. Kessler R, Faller M, Fourgaut G, Mennecier B, Weitzenblum E. Predictive factors of hospitalization for acute exacerbation in a series of 64 patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999;159:158-64.
464. Mushlin AI, Black ER, Connolly CA, Buonaccorso KM, Eberly SW. The necessary length of hospital stay for chronic pulmonary disease. *JAMA* 1991;266:80-3.
465. Regueiro CR, Hamel MB, Davis RB, Desbiens N, Connors AF, Jr., Phillips RS. A comparison of generalist and pulmonologist care for patients hospitalized with severe chronic obstructive pulmonary disease: resource intensity, hospital costs, and survival. SUPPORT Investigators. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. *Am J Med* 1998;105:366-72.
466. Price LC, Lowe D, Hosker HS, Anstey K, Pearson MG, Roberts CM. UK National COPD Audit 2003: Impact of hospital resources and organisation of care on patient outcome following admission for acute COPD exacerbation. *Thorax* 2006;61:837-42.
467. Gravit JH, Al-Rawas OA, Cotton MM, Flanigan U, Irwin A, Stevenson RD. Home treatment of exacerbations of chronic obstructive pulmonary disease by an acute respiratory assessment service. *Lancet* 1998;351:1853-5.
468. Bahadon K, FitzGerald JM. Risk factors of hospitalization and readmission of patients with COPD exacerbation-systematic review. *Int J Chron Obstruct Pulmon Dis* 2007;2:241-51.
469. Cotton MM, Bucknall CE, Dagg KD, et al. Early discharge for patients with exacerbations of chronic obstructive pulmonary disease: a randomized controlled trial. *Thorax* 2000;55:902-6.
470. Hermiz O, Comino E, Marks G, Daffurn K, Wilson S, Harris M. Randomised controlled trial of home based care of patients with chronic obstructive pulmonary disease. *BMJ* 2002;325:938.
471. Hughes SL, Weaver FM, Giobbie-Hurder A, et al. Effectiveness of team-managed home-based primary care: a randomized multicenter trial. *JAMA* 2000;284:2877-85.
472. Siafakas NM, Vermeire P, Pride NB, et al. Optimal assessment and management of chronic obstructive pulmonary disease (COPD). The European Respiratory Society Task Force. *Eur Respir J* 1995;8:1398-420.
473. Wood-Baker R, McGlone S, Venn A, Walters EH. Written action plans in chronic obstructive pulmonary disease increase appropriate treatment for acute exacerbations. *Respirology* 2006;11:619-26.
474. Bischoff EW, Hamd DH, Sedeno M, et al. Effects of written action plan adherence on COPD exacerbation recovery. *Thorax* 2011;66:26-31.

475. Man WD, Polkey MI, Donaldson N, Gray BJ, Moxham J. Community pulmonary rehabilitation after hospitalisation for acute exacerbations of chronic obstructive pulmonary disease: randomised controlled study. *BMJ* 2004;329:1209.
476. Sin DD, Anthonisen NR, Soriano JB, Agusti AG. Mortality in COPD: Role of comorbidities. *Eur Respir J* 2006;28:1245-57.
477. Fabbri LM, Luppi F, Beghe B, Rabe KF. Complex chronic comorbidities of COPD. *Eur Respir J* 2008;31:204-12.
478. Johnston AK, Mannino DM, Hagan GW, Davis KJ, Kiri VA. Relationship between lung function impairment and incidence or recurrence of cardiovascular events in a middle-aged cohort. *Thorax* 2008;63:599-605.
479. Lange P, Mogelvang R, Marott JL, Vestbo J, Jensen JS. Cardiovascular morbidity in COPD: A study of the general population. *COPD* 2010;7:5-10.
480. Brekke PH, Omland T, Smith P, Soyseth V. Underdiagnosis of myocardial infarction in COPD - Cardiac Infarction Injury Score (CIIS) in patients hospitalised for COPD exacerbation. *Respir Med* 2008;102:1243-7.
481. Salpeter S, Ormiston T, Salpeter E. Cardioselective beta-blockers for chronic obstructive pulmonary disease. *Cochrane database of systematic reviews* 2005:CD003566.
482. Calverley PM, Anderson JA, Celli B, et al. Cardiovascular events in patients with COPD: TORCH study results. *Thorax* 2010;65:719-25.
483. Rutten FH, Cramer MJ, Grobbee DE, et al. Unrecognized heart failure in elderly patients with stable chronic obstructive pulmonary disease. *Eur Heart J* 2005;26:1887-94.
484. Hawkins NM, Huang Z, Pieper KS, et al. Chronic obstructive pulmonary disease is an independent predictor of death but not atherosclerotic events in patients with myocardial infarction: analysis of the Valsartan in Acute Myocardial Infarction Trial (VALIANT). *Eur J Heart Fail* 2009;11:292-8.
485. Iversen KK, Kjaergaard J, Akkan D, et al. Chronic obstructive pulmonary disease in patients admitted with heart failure. *J Intern Med* 2008;264:361-9.
486. Iversen KK, Kjaergaard J, Akkan D, et al. The prognostic importance of lung function in patients admitted with heart failure. *Eur J Heart Fail* 2010;12:685-91.
487. Hawkins NM, Jhund PS, Simpson CR, et al. Primary care burden and treatment of patients with heart failure and chronic obstructive pulmonary disease in Scotland. *Eur J Heart Fail* 2010;12:17-24.
488. Hawkins NM, MacDonald MR, Petrie MC, et al. Bisoprolol in patients with heart failure and moderate to severe chronic obstructive pulmonary disease: a randomized controlled trial. *Eur J Heart Fail* 2009;11:684-90.
489. Jabbour A, Macdonald PS, Keogh AM, et al. Differences between beta-blockers in patients with chronic heart failure and chronic obstructive pulmonary disease: a randomized crossover trial. *J Am Coll Cardiol* 2010;55:1780-7.
490. Au DH, Udris EM, Fan VS, Curtis JR, McDonnell MB, Fihn SD. Risk of mortality and heart failure exacerbations associated with inhaled beta-adrenoceptor agonists among patients with known left ventricular systolic dysfunction. *Chest* 2003;123:1964-9.
491. Buch P, Friberg J, Scharling H, Lange P, Prescott E. Reduced lung function and risk of atrial fibrillation in the Copenhagen City Heart Study. *Eur Respir J* 2003;21:1012-6.
492. Madsen H, Brixen K, Hallas J. Screening, prevention and treatment of osteoporosis in patients with chronic obstructive pulmonary disease - a population-based database study. *Clin Respir J* 2010;4:22-9.
493. McAllister DA, Maclay JD, Mills NL, et al. Arterial stiffness is independently associated with emphysema severity in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2007;176:1208-14.
494. Bolton CE, Cannings-John R, Edwards PH, et al. What community measurements can be used to predict bone disease in patients with COPD? *Respir Med* 2008;102:651-7.
495. Bolton CE, Ionescu AA, Shiels KM, et al. Associated loss of fat-free mass and bone mineral density in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004;170:1286-93.

496. The Lung Health Study Research Group. Effect of inhaled triamcinolone on the decline in pulmonary function in chronic obstructive pulmonary disease: Lung Health Study II. *N Engl J Med* 2000;343:1902-9.
497. Kunik ME, Roundy K, Veazey C, et al. Surprisingly high prevalence of anxiety and depression in chronic breathing disorders. *Chest* 2005;127:1205-11.
498. Ng TP, Niti M, Tan WC, Cao Z, Ong KC, Eng P. Depressive symptoms and chronic obstructive pulmonary disease: effect on mortality, hospital readmission, symptom burden, functional status, and quality of life. *Arch Intern Med* 2007;167:60-7.
499. Maurer J, Rebbapragada V, Borson S, et al. Anxiety and depression in COPD: current understanding, unanswered questions, and research needs. *Chest* 2008;134:43S-56S.
500. Eisner MD, Blanc PD, Yelin EH, et al. Influence of anxiety on health outcomes in COPD. *Thorax* 2010;65:229-34.
501. National Institute of Clinical Excellence. Depression in adults with a chronic physical health problem: full guideline. Available at: <http://www.nice.org.uk/nicemedia/pdf/CG91FullGuideline.pdf>. 2009.
502. Coventry PA, Bower P, Keyworth C, Kenning C, Knopp J, Garrett C, Hind D, Malpass A, Dickens C. The effect of complex interventions on depression and anxiety in chronic obstructive pulmonary disease: systematic review and meta-analysis. *PLoS One*. 2013 Apr 5;8(4):e60532.
503. Benfield T, Lange P, Vestbo J. COPD stage and risk of hospitalization for infectious disease. *Chest* 2008;134:46-53.
504. McGarvey LP, Magder S, Burkhart D, Kesten S, Liu D, Manuel RC, Niewoehner DE. Cause-specific mortality adjudication in the UPLIFT® COPD trial: findings and recommendations. *Respir Med* 2012 Apr;106(4):515-21.
505. Zwar NA, Marks GB, Hermiz O, Middleton S, Comino EJ, Hasan I, et al. Predictors of accuracy of diagnosis of chronic obstructive pulmonary disease in general practice. *Med J Aust* 2011 Aug 15;195(4):168-71.
506. Albert P, Agusti A, Edwards L, Tal-Singer R, Yates J, Bakke P, et al. Bronchodilator responsiveness as a phenotypic characteristic of established chronic obstructive pulmonary disease. *Thorax* 2012 Aug;67(8):701-8.
507. Kessler R, Partridge MR, Miravittles M, Cazzola, M, Vogelmeier, C, Leynaud, D, Ostinelli, J. Symptom variability in patients with severe COPD: a pan-European cross-sectional study. *Eur Respir J* 2011;37:264-72.
508. Espinosa de los Monteros MJ, Pena C, Soto Hurtado EJ, Jareno J, Miravittles M. Variability of respiratory symptoms in severe COPD. *Arch Bronconeumol* 2012;48:3-7.
509. van der Molen T, Willemse BW, Schokker S, ten Hacken NH, Postma DS, Juniper EF. Development, validity and responsiveness of the Clinical COPD Questionnaire. *Health Qual Life Outcomes*. 2003 Apr 28;1:13
510. Reda AA, Kotz D, Kocks JW, Wesseling G, van Schayck CP. Reliability and validity of the clinical COPD questionnaire and chronic respiratory questionnaire. *Respir Med* 2010 Nov;104(11):1675-82.
511. Trappenburg JC, Touwen I, de Weert-van Oene GH, Bourbeau J, Monninkhof EM, Verheij TJ, et al. Detecting exacerbations using the Clinical COPD Questionnaire. *Health Qual Life Outcomes* 2010 Sep 16;8:102
512. Aaron SD, Donaldson GC, Whitmore GA, Hurst JR, Ramsay T, Wedzicha JA. Time course and pattern of COPD exacerbation onset. *Thorax* 2012 Mar;67(3):238-43.
513. Kornmann O, Dahl R, Centanni S, Dogra A, Owen R, Lassen C, Kramer B. Once-daily indacaterol vs twice-daily salmeterol for COPD: a placebo-controlled comparison. *Eur Respir J* 2011;37:273-9.
514. Dahl R, Chung KF, Buhl R, Magnussen H, Nonikov V, Jack D, et al; INVOLVE (INDacaterol: Value in COPD: Longer Term Validation of Efficacy and Safety) Study Investigators. Efficacy of a new once-daily long-acting inhaled beta2-agonist indacaterol versus twice-daily formoterol in COPD. *Thorax* 2010;65:473-9.
515. Buhl R, Dunn LJ, Disdier C, Lassen C, Amos C, Henley M, Kramer B; INTENSITY study investigators. Blinded 12-week comparison of once-daily indacaterol and tiotropium in COPD. *Eur Respir J* 2011;38:797-803.
516. Chapman KR, Rennard SI, Dogra A, Owen R, Lassen C, Kramer B; INDORSE Study Investigators. Long-term safety and efficacy of indacaterol, a long-acting β 2-agonist, in subjects with COPD: a randomized, placebo-controlled study. *Chest* 2011;140:68-75.

517. Chong J, Karner C, Poole P. Tiotropium versus long-acting beta-agonists for stable chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 Sep 12;9:CD009157.
518. Karner C, Chong J, Poole P. Tiotropium versus placebo for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 Jul 11;7:CD009285.
519. Beasley R, Singh S, Loke YK, Enright P, Furberg CD. Call for worldwide withdrawal of tiotropium Respimat mist inhaler. *BMJ* 2012 Nov 9;345:e7390.
520. Yang IA, Clarke MS, Sim EH, Fong KM. Inhaled corticosteroids for stable chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 Jul 11;7:CD002991.
521. Doherty DE, Tashkin DP, Kerwin E, Knorr BA, Shekar T, Banerjee S, Staudinger H. Effects of mometasone furoate/formoterol fumarate fixed-dose combination formulation on chronic obstructive pulmonary disease (COPD): results from a 52-week Phase III trial in subjects with moderate-to-very severe COPD. *Int J Chron Obstruct Pulmon Dis* 2012;7:57-71.
522. Nannini LJ, Lasserson TJ, Poole P. Combined corticosteroid and long-acting beta(2)-agonist in one inhaler versus long-acting beta(2)-agonists for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 Sep 12;9:CD006829.
523. Poole P, Black PN, Cates CJ. Mucolytic agents for chronic bronchitis or chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 Aug 15;8:CD001287
524. Puhan MA, Gimeno-Santos E, Scharplatz M, Troosters T, Walters EH, Steurer J. Pulmonary rehabilitation following exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2011 Oct 5;(10):CD005305.
525. Au DH, Udris EM, Engelberg RA, Diehr PH, Bryson CL, Reinke LF, Curtis JR. A randomized trial to improve communication about end-of-life care among patients with COPD. *Chest* 2012 Mar;141(3):726-35.
526. Sudore RL, Fried TR. Redefining the "planning" in advance care planning: preparing for end-of-life decision making. *Ann Intern Med* 2010 Aug 17;153(4):256-61.
527. Curtis JR, Engelberg RA, Bensink ME, Ramsey SD. End-of-Life Care in the Intensive Care Unit: Can We Simultaneously Increase Quality and Reduce Costs? *Am J Respir Crit Care Med* 2012 Oct 1;186(7):587-92.
528. Pinnock H, Kendall M, Murray SA, Worth A, Levack P, Porter M, MacNee W, Sheikh A. Living and dying with severe chronic obstructive pulmonary disease: multi-perspective longitudinal qualitative study. *BMJ* 2011 Jan 24;342:d142.
529. Ahmedzai S, Balfour-Lynn IM, Bewick T, Buchdahl R, Coker RK, Cummin AR, et al. British Thoracic Society Standards of Care Committee. Managing passengers with stable respiratory disease planning air travel: British Thoracic Society recommendations. *Thorax* 2011 Sep;66 Suppl 1:i1-30.
530. Murray SA, Kendall M, Boyd K, Sheikh A. Illness trajectories and palliative care. *BMJ*. 2005;330:1007-11.
531. Eriksen N, Vestbo J. Management and survival of patients admitted with an exacerbation of COPD: comparison of two Danish patient cohorts. *Clin Respir J* 2010 Oct;4(4):208-14.
532. Groenewegen KH, Schols AM, Wouters EF. Mortality and mortality-related factors after hospitalization for acute exacerbation of COPD. *Chest* 2003;124:459-67.
533. Gudmundsson G, Ulrik CS, Gislason T, Lindberg E, Brøndum E, Bakke P, Janson C. Long-term survival in patients hospitalized for chronic obstructive pulmonary disease: a prospective observational study in the Nordic countries. *Int J Chron Obstruct Pulmon Dis* 2012;7:571-6.
534. National Consensus Project for Quality Palliative Care: Clinical Practice Guidelines for quality palliative care, executive summary. *J Palliat Med* 2004;7(5):611-27.
535. Au DH, Udris EM, Fihn SD, McDonnell MB, Curtis JR. Differences in health care utilization at the end of life among patients with chronic obstructive pulmonary disease and patients with lung cancer. *Arch Intern Med* 2006;166(3):326-31.
536. Levy MH, Adolph MD, Back A, Block S, Codada SN, Dalai S, et al. Palliative care. *J Natl Compr Canc Netw* 2012 Oct1;10(10):1284-309.

537. Morrison RS, Maroney-Galin C, Kralovec PD, Meier DE. The growth of palliative care programs in United States hospitals. *J Palliat Med* 2005 Dec; 8(6):1127-34.
538. Hanania NA, Crater GD, Morris AN, Emmett AH, O'Dell DM, Niewoehner DE. Benefits of adding fluticasone propionate/salmeterol to tiotropium in moderate to severe COPD. *Respir Med* 2012 Jan;106(1):91-101.
539. Welte T, Miravittles M, Hernandez P, Eriksson G, Peterson S, Polanowski T, Kessler R. Efficacy and tolerability of budesonide/formoterol added to tiotropium in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2009 Oct 15;180(8):741-50.
540. Loke YK, Cavallazzi R, Singh S. Risk of fractures with inhaled corticosteroids in COPD: systematic review and meta-analysis of randomized controlled trials and observational studies. *Thorax* 2011 Aug;66(8):699-708.
541. Alia I, de la Cal MA, Esteban A, Abella A, Ferrer R, Molina FJ, et al. Efficacy of corticosteroid therapy in patients with an acute exacerbation of chronic obstructive pulmonary disease receiving ventilatory support. *Arch Intern Med* 2011 Nov 28;171(21):1939-46.
542. Walters JA, Wang W, Morley C, Soltani A, Wood-Baker R. Different durations of corticosteroid therapy for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2011 Oct 5;(10):CD006897.
543. Chandra D, Stamm JA, Taylor B, Ramos RM, Satterwhite L, Krishnan JA, et al. Outcomes of noninvasive ventilation for acute exacerbations of chronic obstructive pulmonary disease in the United States, 1998-2008. *Am J Respir Crit Care Med* 2012 Jan 15;185(2):152-9.
544. Jeppesen E, Brurberg KG, Vist GE, Wedzicha JA, Wright JJ, Greenstone M, Walters JA. Hospital at home for acute exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 May 16;5:CD003573
545. Bucknall CE, Miller G, Lloyd SM, Cleland J, McCluskey S, Cotton M, et al. Glasgow supported self-management trial (GSuST) for patients with moderate to severe COPD: randomised controlled trial. *BMJ* 2012 Mar 6;344:e1060.
546. Mainguy V, Girard D, Maltais F, Saey D, Milot J, Sénéchal M, Poirier P, Provencher S. Effect of bisoprolol on respiratory function and exercise capacity in chronic obstructive pulmonary disease. *Am J Cardiol* 2012 Jul 15;110(2):258-63.
547. Lainscak M, Podbregar M, Kovacic D, Rozman J, von Haehling S. Differences between bisoprolol and carvedilol in patients with chronic heart failure and chronic obstructive pulmonary disease: a randomized trial. *Respir Med* 2011 Oct;105 Suppl 1:S44-9.
548. Jones PW. COPD assessment test --rationale, development, validation and performance. *J COPD* 2013;10(2):269-71.
549. Miravittles M, Soriano JB, García-Río F, Muñoz L, Duran-Tauleria E, Sanchez G, et al. Prevalence of COPD in Spain: impact of undiagnosed COPD on quality of life and daily life activities. *Thorax*. 2009;64(10):863-8.
550. D'Urzo A, Ferguson G, van Noord J, Hirata K, Martin C, Horton R, et al. Efficacy and safety of once-daily NVA237 in patients with moderate-to-severe COPD: the GLOW1 trial. *Respir Res*. 2011;12(1):156.
551. Kerwin FM, D'Urzo AD, Gelb AF, Lakkis H, Gil EG, Caracta CF. Efficacy and Safety of a 12-week Treatment with Twice-daily Acclidinium Bromide in COPD Patients (ACCORD COPD I). *J COPD*. 2012;9(2):90-101.
552. Jones PW, Singh D, Bateman ED, Agusti A, Lamarca R, de Miquel G, et al. Efficacy and safety of twice-daily acclidinium bromide in COPD patients: the ATTAIN study. *Eur Respir J*. 2012;40(4):830-6.
553. Vogelmeier CF, Bateman ED, Pallante J, Alagappan VKT, D'Andrea P, Chen H, et al. Efficacy and safety of once-daily QVA149 compared with twice-daily salmeterol/fluticasone in patients with chronic obstructive pulmonary disease (ILLUMINATE): a randomised, double-blind, parallel group study. *The Lancet Respir Med*. 2013;1(1):51-60.
554. Jones P, Tabberer M, Chen W-H. Creating scenarios of the impact of COPD and their relationship to COPD assessment test (CATTM) scores. *BMC Pul Med*. 2011;11(1):42.
555. Jones PW, Adamek L, Nadeau G, Banik N. Comparisons of health status scores with MRC grades in COPD: implications for the GOLD 2011 classification. *Eur Respir J* 2013;42:647-54.
556. Soler-Cataluña JJ, Martínez-García MÁ, P. RS, Salcedo E, Navarro M, Ochando R. Severe acute exacerbations and mortality in patients with chronic obstructive pulmonary disease. *Thorax* 2005;60:925-31.

557. Polkey MI, Spruit MA, Edwards LD, Watkins ML, Pinto-Plata V, Vestbo J, et al; Evaluation of COPD Longitudinally to Identify Predictive Surrogate Endpoints (ECLIPSE) Study Investigators. Six-minute-walk test in chronic obstructive pulmonary disease: minimal clinically important difference for death or hospitalization. *Am J Respir Crit Care Med* 2013 Feb 15;187(4):382-6.
558. Kerwin E, Hébert J, Gallagher N, Martin C, Overend T, Alagappan VK, Lu Y, Banerji D. Efficacy and safety of NVA237 versus placebo and tiotropium in patients with COPD: the GLOW2 study. *Eur Respir J* 2012 Nov;40(5):1106-14.
559. Wise RA, Anzueto A, Cotton D, Dahl R, Devins T, Disse B, et al for the TIOSPIR Investigators. Tiotropium Respimat Inhaler and the Risk of Death in COPD. *N Engl J Med* 2013 Oct 17;369(16):1491-1501.
560. Bateman ED, Ferguson GT, Barnes N, Gallagher N, Green Y, Henley M, Banerji D. Dual bronchodilation with QVA149 versus single bronchodilator therapy: the SHINE study. *Eur Respir J* 2013 Dec;42(6):1484-94.
561. Wedzicha JA, Decramer M, Ficker JH, Niewoehner DE, SandstroANm T, Taylor AF, et al. Analysis of chronic obstructive pulmonary disease exacerbations with the dual bronchodilator QVA149 compared with glycopyrronium and tiotropium (SPARK): a randomised, double-blind, parallel-group study. *Lancet Respir Med* 2013;1:199-209.
562. Tse HN, Raiteri L, Wong KY, Yee KS, Ng LY, Wai KY, Loo CK, Chan MH. High-dose N-acetylcysteine in stable COPD: the 1-year, double-blind, randomized, placebo-controlled HIACE study. *Chest* 2013 Jul;144(1):106-18.
563. Egan C, Deering BM, Blake C, Fullen BM, McCormack NM, Spruit MA, Costello RW. Short term and long term effects of pulmonary rehabilitation on physical activity in COPD. *Respir Med* 2012 Dec;106(12):1671-9.
564. Ferreira IM, Brooks D, White J, Goldstein R. Nutritional supplementation for stable chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 Dec 12;12:CD000998.
565. Gagnon P, Saey D, Provencher S, Milot J, Bourbeau J, Tan WC, Martel S, Maltais F. Walking exercise response to bronchodilation in mild COPD: a randomized trial. *Respir Med* 2012 Dec;106(12):1695-705.
566. Bischoff EW, Akkermans R, Bourbeau J, van Weel C, Vercoelen JH, Schermer TR. Comprehensive self management and routine monitoring in chronic obstructive pulmonary disease patients in general practice: randomised controlled trial. *BMJ* 2012 Nov 28;345:e7642.
567. Leuppi JD, Schuetz P, Bingisser R, Bodmer M, Briel M, Drescher T, et al.. Short-term vs conventional glucocorticoid therapy in acute exacerbations of chronic obstructive pulmonary disease: the REDUCE randomized clinical trial. *JAMA* 2013 Jun 5;309(21):2223-31.
568. Edwards L, Shirtcliffe P, Wadsworth K, Healy B, Jefferies S, Weatherall M, Beasley R; Magnesium COPD Study Team. Use of nebulised magnesium sulphate as an adjuvant in the treatment of acute exacerbations of COPD in adults: a randomised double-blind placebo-controlled trial. *Thorax* 2013 Apr;68(4):338-43.
569. Vollenweider DJ, Jarrett H, Steurer-Stey CA, Garcia-Aymerich J, Puhan MA. Antibiotics for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2012 Dec 12;12:CD010257.
570. Rizkallah J, Man SF, Sin DD. Prevalence of pulmonary embolism in exacerbations of COPD: a systematic review and metaanalysis. *Chest* 2009 Mar;135(3):786-93.
571. Gunen H, Gulbas G, In E, Yetkin O, Hacievliyagil SS. Venous thromboemboli and exacerbations of COPD. *Eur Respir J* 2010;35(6):1243-8.
572. Qaseem A, Chou R, Humphrey LL, Starkey M, Shekelle P; Clinical Guidelines Committee of the American College of Physicians. Venous thromboembolism prophylaxis in hospitalized patients: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2011 Nov 1;155(9):625-32.
573. Kahn SR, Lim W, Dunn AS, Cushman M, Dentali F, Akl EA, et al; American College of Chest Physicians. Prevention of VTE in nonsurgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012 Feb;141(2 Suppl):e195S-226S.
574. Bertoletti L, Quenet S, Laporte S, Sahuquillo JC, Conget F, Pedrajas JM, Martin M, Casado I, Riera-Mestre A, Monreal M; RIETE Investigators. Pulmonary embolism and 3-month outcomes in 4036 patients with venous thromboembolism and chronic obstructive pulmonary disease: data from the RIETE registry. *Respir Res* 2013 Jul 18;14:75.

575. Cartwright M, Hirani SP, Rixon L, Beynon M, Doll H, Bower P, et al; Whole Systems Demonstrator Evaluation Team. Effect of telehealth on quality of life and psychological outcomes over 12 months (Whole Systems Demonstrator telehealth questionnaire study): nested study of patient reported outcomes in a pragmatic, cluster randomised controlled trial. *BMJ* 2013 Feb 26;346:f653.
576. Henderson C, Knapp M, Fernández JL, Beecham J, Hirani SP, Cartwright M, et al; Whole System Demonstrator evaluation team. Cost effectiveness of telehealth for patients with long term conditions (Whole Systems Demonstrator telehealth questionnaire study): nested economic evaluation in a pragmatic, cluster randomised controlled trial. *BMJ* 2013 Mar 20;346:f1035.
577. Pinnock H, Hanley J, McCloughan L, Todd A, Krishan A, Lewis S, et al. Effectiveness of telemonitoring integrated into existing clinical services on hospital admission for exacerbation of chronic obstructive pulmonary disease: researcher blind, multicentre, randomised controlled trial. *BMJ* 2013 Oct 17;347:f6070.
578. Almagro P, Cabrera FJ, Diez J, Boixeda R, Alonso Ortiz MB, Murio C, Soriano JB; Working Group on COPD, Spanish Society of Internal Medicine. Comorbidities and short-term prognosis in patients hospitalized for acute exacerbation of COPD: the EPOC en Servicios de medicina interna (ESMI) study. *Chest* 2012 Nov;142(5):1126-33.
579. Mainguy V, Girard D, Maltais F, Saey D, Milot J, Sénéchal M, Poirier P, Provencher S. Effect of bisoprolol on respiratory function and exercise capacity in chronic obstructive pulmonary disease. *Am J Cardiol* 2012 Jul 15;110(2):258-63.
580. Stefan MS, Rothberg MB, Priya A, Pekow PS, Au DH, Lindenauer PK. Association between beta-blocker therapy and outcomes in patients hospitalised with acute exacerbations of chronic obstructive lung disease with underlying ischaemic heart disease, heart failure or hypertension. *Thorax* 2012 Nov;67(11):977-84.
581. O'Brien C, Guest PJ, Hill SL, Stockley RA. Physiological and radiological characterisation of patients diagnosed with chronic obstructive pulmonary disease in primary care. *Thorax* 2000;55:635-642.
582. Patel IS, Vlahos I, Wilkinson TMA, et al. Bronchiectasis, exacerbation indices, and inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004;170:400-407.
583. Martínez-García MA, de la Rosa Carrillo D, Soler-Cataluña JJ, Donat-Sanz Y, Serra PC, Lerma MA, Ballestín J, Sánchez IV, Selma Ferrer MJ, Dalfo AR, Valdecillos MB. Prognostic value of bronchiectasis in patients with moderate-to-severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2013 Apr 15;187(8):823-31.
584. Dirven JA, Tange HJ, Muris JW, van Haaren KM, Vink G, van Schayck OC. Early detection of COPD in general practice: implementation, workload and socioeconomic status. A mixed methods observational study. *Prim Care Respir J* 2013 Sep;22(3):338-43.
585. Beekman E, Mesters I, Hendriks EJ, Klaassen MP, Gosselink R, van Schayck OC, de Bie RA. Course length of 30 metres versus 10 metres has a significant influence on six-minute walk distance in patients with COPD: an experimental crossover study. *J Physiother* 2013 Sep;59(3):169-76.
586. Kew KM, Mavergames C, Walters JA. Long-acting beta2-agonists for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2013 Oct 15;10:CD010177.
587. Cheyne L, Irvin-Sellers MJ, White J. Tiotropium versus ipratropium bromide for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2013 Sep 16;9:CD009552.
588. Beier J, Kirsten AM, Mróz R, Segarra R, Chuecos F, Caracta C, Gil EG. Efficacy and safety of acclidinium bromide compared with placebo and tiotropium in patients with moderate-to-severe chronic obstructive pulmonary disease: results from a 6-week, randomized, controlled Phase IIIb study. *COPD* 2013 Aug;10(4):511-22.
589. Donohue JF, Maleki-Yazdi MR, Kilbride S, Mehta R, Kalberg C, Church A. Efficacy and safety of once-daily umeclidinium/vilanterol 62.5/25 mcg in COPD. *Respir Med* 2013 Oct;107(10):1538-46.
590. Magnussen H, Disse B, Rodriguez-Roisin R, Kirsten A, Watz H, Tetzlaff K, Towse L, Finnigan H, Dahl R, Decramer M, Chanez P, Wouters EF, Calverley PM; WISDOM Investigators. Withdrawal of inhaled glucocorticoids and exacerbations of COPD. *N Engl J Med* 2014 Oct 2;371(14):1285-94.

591. Kew KM, Seniukovich A. Inhaled steroids and risk of pneumonia for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2014 Mar10;3:CD010115.
592. Horita N, Miyazawa N, Morita S, Kojima R, Inoue M, Ishigatsubo Y, Kaneko T. Evidence suggesting that oral corticosteroids increase mortality in stable chronic obstructive pulmonary disease. *Respir Res* 2014 Apr 3;15:37.
593. Han MK, Tayob N, Murray S, Dransfield MT, Washko G, Scanlon PD, et al. Predictors of chronic obstructive pulmonary disease exacerbation reduction in response to daily azithromycin therapy. *Am J Respir Crit Care Med* 2014 Jun 15;189(12):1503-8. Sep;42(3):636-46.
594. Zheng JP, Wen FQ, Bai CX, Wan HY, Kang J, Chen P, et al; PANTHEON study group. Twice daily N-acetylcysteine 600 mg for exacerbations of chronic obstructive pulmonary disease (PANTHEON): a randomised, double-blind placebo-controlled trial. *Lancet Respir Med* 2014 Mar;2(3):187-94.
595. Blanco I, Santos S, Gea J, Güell R, Torres F, Gimeno-Santos E, et al. Sildenafil to improve respiratory rehabilitation outcomes in COPD: a controlled trial. *Eur Respir J* 2013 Oct;42(4):982-92.
596. Lehouck A, Mathieu C, Carremans C, Baeke F, Verhaegen J, Van Eldere J, et al. High doses of vitamin D to reduce exacerbations in chronic obstructive pulmonary disease. *Ann Intern Med* 2012 Jan 17;156(2):105-14.
597. Struik FM, Sprooten RT, Kerstjens HA, Bladder G, Zijnen M, Asin J, et al. Nocturnal non-invasive ventilation in COPD patients with prolonged hypercapnia after ventilatory support for acute respiratory failure: a randomised, controlled, parallel-group study. *Thorax* 2014 Sep;69(9):826-34.
598. Struik FM, Lacasse Y, Goldstein RS, Kerstjens HA, Wijkstra PJ. Nocturnal noninvasive positive pressure ventilation in stable COPD: a systematic review and individual patient data meta-analysis. *Respir Med* 2014 Feb;108(2):329-37.
599. Köhnlein T, Windisch W, Köhler D, Drabik A, Geiseler J, Hartl S, et al. Non-invasive positive pressure ventilation for the treatment of severe stable chronic obstructive pulmonary disease: a prospective, multicentre, randomised, controlled clinical trial. *Lancet Respir Med* 2014 Sep;2(9):698-705
600. Kruis AL, Smidt N, Assendelft WJJ, Gussekloo J, Boland MRS, et al. Integrated disease management interventions for patients with chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2013, Issue 10. Art. No.: CD009437.
601. Kruis AL, Boland MRS, Assendelft WJJ, Gussekloo J, Tsiachristas A, Stijnen T, et al. Effectiveness of integrated disease management for primary care chronic obstructive pulmonary disease patients: results of cluster randomised trial. *BMJ* 2014;349:g5392
602. Vincken W, Aumann J, Chen H, Henley M, McBryan D, Goyal P. Efficacy and safety of co-administration of once-daily indacaterol and glycopyrronium versus indacaterol alone in COPD patients: the GLOW6 study. *Int J COPD* 2014 Feb 24;9:215-28.
603. Faustini A, Staroggia M, Colais P, Berti G, Bisanti L, Cadum E, et al; EpiAir Collaborative Group. Air pollution and multiple acute respiratory outcomes. *Eur Respir J* 2013 Aug;42(2):304-13.
604. Hoogendoorn M, Hoogenveen RT, Rutten-van Mölken MP, Vestbo J, Feenstra TL. Case fatality of COPD exacerbations: a meta-analysis and statistical modelling approach. *Eur. Respir. J.* 2011; 37: 508–515.
605. Piquet J, Chavaillon J-M, David P, Martin F, Blanchon F, Roche N, French College of General Hospital Respiratory Physicians (CPHG). High-risk patients following hospitalisation for an acute exacerbation of COPD. *Eur Respir J* 2013; 42: 946–955.
606. Singanayagam A, Schembri S, Chalmers JD. Predictors of Mortality in Hospitalized Adults with Acute Exacerbation of Chronic Obstructive Pulmonary Disease. A Systematic Review and Meta-analysis. *Ann. Am. Thorac. Soc.* 2013; 10: 81–89.
607. Garcia-Aymerich J, Gómez FP, Benet M, Farrero E, Basagaña X, Gayete À, et al. Identification and prospective validation of clinically relevant chronic obstructive pulmonary disease (COPD) subtypes. *Thorax* 2011; 66: 430–437.
608. Criner GJ, Connett JE, Aaron SD, Albert RK, Bailey WC, Casaburi R, et al; COPD Clinical Research Network; Canadian Institutes of Health Research. Simvastatin for the prevention of exacerbations in moderate-to-severe COPD. *N Engl J Med* 2014 Jun 5;370(23):2201-10.

609. Miller J, Edwards LD, Agustí A, Bakke P, Calverley PM, Celli B, et al. Evaluation of COPD Longitudinally to Identify Predictive Surrogate Endpoints (ECLIPSE) Investigators. Comorbidity, systemic inflammation and outcomes in the ECLIPSE cohort. *Respir Med* 2013 Sep;107(9):1376-84.
610. Campo G, Guastaroba P, Marzocchi A, Santarelli A, Varani E, Vignali L, et al. Impact of COPD on long-term outcome after ST-segment elevation myocardial infarction receiving primary percutaneous coronary intervention. *Chest* 2013 Sep;144(3):750-7.
611. Martinez CH, Okajima Y, Murray S, Washko GR, Martinez FJ, Silverman EK, et al.; COPDGene Investigators. Impact of self-reported gastroesophageal reflux disease in subjects from COPDGene cohort. *Respir Res* 2014 Jun 3;15:62.
612. Dodd JW, Getov SV, Jones PW. Cognitive function in COPD. *Eur Respir J* 2010;35(4):913-22.
613. Singh B, Mielke MM, Parsaik AK, Cha RH, Roberts RO, Scanlon PD, et al. A prospective study of chronic obstructive pulmonary disease and the risk for mild cognitive impairment. *JAMA Neurol* 2014 May;71(5):581-8.
614. She L, Yang P, Wang Y, Qin X, Fan L, Wang Y, Gao G, Luo G, et al. Chinese water-pipe smoking and the risk of capno. *Chest*. 2014 Oct;146(4):924-31.
615. Ramirez-Venegas A, Sansores RH, Quintana-Carrillo RH, Velazquez-Uncal M, Hernandez-Zenteno RJ, Sanchez-Romero C, et al. FEV1 decline in patients with chronic obstructive pulmonary disease associated with biomass exposure. *Am J Respir Crit Care Med*. 2014 Nov 1;190(9):996-1002.
616. Putchu N, Drummond MB, Connett JE, Scanlon PD, Tashkin DP, Hansel NN, Wise RA. Chronic productive cough is associated with death in smokers with early COPD. *COPD*. 2014 Aug;11(4):451-8.
617. Kon SS, Dilaver D, Mittal M, Nolan CM, Clark AL, Canavan JL, et al. The Clinical COPD Questionnaire: response to pulmonary rehabilitation and minimal clinically important difference. *Thorax*. 2014 Sep;69(9):793-8.
618. Rossi A, Guerriero M, Corrado A; OPTIMO/AIPO Study Group. Withdrawal of inhaled corticosteroids can be safe in COPD patients at low risk of exacerbation: a real-life study on the appropriateness of treatment in moderate COPD patients (OPTIMO). *Respir Res*. 2014 Jul 8;15:77.
619. Dransfield MT, Feldman G, Korenblat P, LaForce CF, Locantore N, Pistolesi M, et al. Efficacy and safety of once-daily fluticasone furoate/vilanterol (100/25 mcg) versus twice-daily fluticasone propionate/salmeterol (250/50 mcg) in COPD patients. *Respir Med*. 2014 Aug;108(8):1171-9.
620. Criner GJ, Bourbeau J, Diekemper RL, Ouellette DR, Goodridge D, Hernandez P, et al. Prevention of acute exacerbations of COPD: American College of Chest Physicians and Canadian Thoracic Society Guideline. *Chest*. 2015 Apr;147(4):894-2.
621. P Yu T, Fain K, Boyd CM, Singh S, Weiss CO, Li T, et al. Benefits and harms of roflumilast in moderate to severe COPD. *Thorax*. 2014 Jul;69(7):616-22. g 28: Busch AM, Scott-Sheldon LA, Pierce J, Chattillion EA, Cunningham K, Buckley ML, et al. Depressed mood predicts pulmonary rehabilitation completion among women, but not men. *Respir Med*. 2014 Jul;108(7):1007-13.
622. Busch AM, Scott-Sheldon LA, Pierce J, Chattillion EA, Cunningham K, Buckley ML, et al. Depressed mood predicts pulmonary rehabilitation completion among women, but not men. *Respir Med*. 2014 Jul;108(7):1007-13.
623. Konecny T, Park JY, Somers KR, Konecny D, Orban M, Soucek F, Parker KO, et al. Relation of chronic obstructive pulmonary disease to atrial and ventricular arrhythmias. *Am J Cardiol*. 2014 Jul 15;114(2):272-7.
624. Walters JA, Tan DJ, White CJ, Wood-Baker R. Different durations of corticosteroid therapy for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2014 Dec 10;12:CD006897.f
625. Ingebrigtsen TS, Marott JL, Nordestgaard BG, Lange P, Hallas J, Vestbo J. Statin use and exacerbations in individuals with chronic obstructive pulmonary disease. *Thorax*. 2015 Jan;70(1):33-40.

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

Appendix

Asthma and COPD Overlap Syndrome (ACOS)

Appendix

Diagnosis Of Diseases Of Chronic Airflow Limitation: Asthma, COPD and Asthma–COPD Overlap Syndrome

A joint project of GINA and GOLD[#]

- Distinguishing asthma from COPD can be problematic, particularly in smokers and older adults
- ACOS is identified by the features that it shares with both asthma and COPD.
- A stepwise approach to diagnosis is advised, comprising recognition of the presence of a chronic airways disease, syndromic categorization as asthma, COPD or the overlap between asthma and COPD (the Asthma COPD Overlap Syndrome (ACOS)), confirmation by spirometry and, if necessary, referral for specialized investigations.
- Although initial recognition and treatment of ACOS may be made in primary care, referral for confirmatory investigations is encouraged, as outcomes for ACOS are often worse than for asthma or COPD alone.
- Initial treatment should be selected to ensure that:
 - o Patients with features of asthma receive adequate controller therapy including inhaled corticosteroids, but not long-acting bronchodilators alone (as monotherapy), and
 - o Patients with COPD receive appropriate symptomatic treatment with bronchodilators or combination therapy, but not inhaled corticosteroids alone (as monotherapy).
- The consensus-based description of the Asthma COPD Overlap Syndrome (ACOS) is intended to stimulate further study of the character and treatments for this common clinical problem.

KEY POINTS

OBJECTIVE

This consensus-based document aims to assist clinicians to:

- Identify patients who have a disease of chronic airflow limitation
- Distinguish asthma from COPD and the Asthma-COPD Overlap Syndrome (ACOS)
- Decide on initial treatment and/or need for referral

[#] This chapter also appears in the Global Strategy for Asthma Management and Prevention 2014, available from <http://www.ginasthma.org>

DEFINITIONS

Table 1. Current definitions of asthma and COPD, and clinical description of ACOS

<p style="text-align: center;">Asthma</p> <p>Asthma is a heterogeneous disease, usually characterized by chronic airway inflammation. It is defined by the history of respiratory symptoms such as wheeze, shortness of breath, chest tightness and cough that vary over time and in intensity, together with variable expiratory airflow limitation. [GINA 2014]</p>
<p style="text-align: center;">COPD</p> <p>COPD is a common preventable and treatable disease, characterized by persistent airflow limitation that is usually progressive and associated with enhanced chronic inflammatory responses in the airways and the lungs to noxious particles or gases. Exacerbations and comorbidities contribute to the overall severity in individual patients. [GOLD 2014]²¹</p>
<p style="text-align: center;">Asthma-COPD Overlap Syndrome (ACOS) – a description for clinical use</p> <p>Asthma-COPD overlap syndrome (ACOS) is characterized by persistent airflow limitation with several features usually associated with asthma and several features usually associated with COPD. ACOS is therefore identified by the features that it shares with both asthma and COPD.</p>

A summary of the typical characteristics of asthma, COPD and ACOS is presented in Table 2a, showing the similarities and differences in history and investigations.

STEP-WISE APPROACH TO DIAGNOSIS OF PATIENTS WITH RESPIRATORY SYMPTOMS

Step 1: Does the patient have chronic airways disease?

A first step in diagnosing these conditions is to identify patients at risk of, or with significant likelihood of having chronic airways disease, and to exclude other potential causes of respiratory symptoms. This is based on a detailed medical history, physical examination, and other investigations.^{3,22-24}

Clinical history

Features that should prompt consideration of chronic airways disease include:

- History of chronic or recurrent cough, sputum production, dyspnea, or wheezing; or repeated acute lower respiratory tract infections
- Report of a previous doctor diagnosis of asthma or COPD
- History of prior treatment with inhaled medications
- History of smoking tobacco and/or other substances
- Exposure to environmental hazards, e.g. occupational or domestic exposures to airborne pollutants

Physical examination

- May be normal
- Evidence of hyperinflation and other features of chronic lung disease or respiratory insufficiency
- Abnormal auscultation (wheeze and/or crackles)

Radiology

- May be normal, particularly in early stages
- Abnormalities on chest X-ray or CT scan (performed for other reasons such as screening for lung cancer), including hyperinflation, airway wall thickening, air trapping, hyperlucency, bullae or other features of emphysema.
- May identify an alternative diagnosis, including bronchiectasis, evidence of lung infections such as tuberculosis, interstitial lung diseases or cardiac failure.

Screening questionnaires

Many screening questionnaires have been proposed to help the clinician identifying subjects at risk of chronic airways disease, based on the above risk factors and clinical features.²⁵⁻²⁷ These questionnaires are usually context-specific, so they are not necessarily relevant to all countries (where risk factors and comorbid diseases differ), to all practice settings and uses (population screening versus primary or secondary care), or to all groups of patients (case-finding versus self-presenting with respiratory symptoms versus referred consultation). Examples of these questionnaires are provided on both the GINA and GOLD websites.

STEP 2. The syndromic diagnosis of asthma, COPD and ACOS in an adult patient

Given the extent of overlap between features of asthma and COPD (Table 2a), the approach proposed focuses on the features that are most helpful in distinguishing asthma and COPD (Table 2b).

a. Assemble the features that favor a diagnosis of asthma or of COPD

From a careful history that considers age, symptoms (in particular onset and progression, variability, seasonality or periodicity and persistence), past history, social and occupational risk factors including smoking history, previous diagnoses and treatment and response to treatment, the features favoring the diagnostic profile of asthma or of COPD can be assembled. The check boxes in Table 2b can be used to identify the features that are most consistent with asthma and/or COPD. Note that not all of the features of asthma and COPD are listed, but only those that most easily distinguish between asthma and COPD.

b. Compare the number of features in favor of a diagnosis of asthma or a diagnosis of COPD

From Table-2b, count the number of checked boxes in each column. Having several (three or more) of the features listed for either asthma or for COPD, in the absence of those for the alternative diagnosis, provides a strong likelihood of a correct diagnosis.²⁷ However, the absence of any of these features has less predictive value, and does not rule out the diagnosis of either disease. For example, a history of allergies increases the probability that respiratory symptoms are due to asthma, but is not essential for the diagnosis of asthma since non-allergic asthma is a well-recognized asthma phenotype; and atopy is common in the general population including in patients who develop COPD in later years. When a patient has similar numbers of features of both asthma and COPD, the diagnosis of ACOS should be considered.

c. Consider the level of certainty around the diagnosis of asthma or COPD, or whether there are features of both suggesting Asthma-COPD Overlap Syndrome

In the absence of pathognomonic features, clinicians recognize that diagnoses are made on the weight of evidence, provided there are no features that clearly make the diagnosis untenable. Clinicians are able to provide an estimate of their level of certainty and factor it into their decision to treat. Doing so consciously may assist in the selection of treatment and, where there is significant doubt, it may direct therapy towards the safest option - namely, treatment for the condition that should not be missed and left untreated.

Table 2a. Usual features of asthma, COPD and ACOS

Table 2a. Usual features of asthma, COPD and ACOS				Table 2b. Features that favor asthma or COPD	
Feature	Asthma	COPD	ACOS	Favors Asthma	Favors COPD
Age of onset	Usually childhood onset but can commence at any age.	Usually > 40 years of age	Usually age ≥40 years, but may have had symptoms in childhood or early adulthood	Onset before age 20 years	Onset after age 40 years
Pattern of respiratory symptoms	Symptoms may vary over time (day to day, or over longer periods), often limiting activity. Often triggered by exercise, emotions including laughter, dust or exposure to allergens	Chronic usually continuous symptoms, particularly during exercise, with 'better' and 'worse' days	Respiratory symptoms including exertional dyspnea are persistent but variability may be prominent	<ul style="list-style-type: none"> <input type="checkbox"/> Variation in symptoms over minutes, hours or days <input type="checkbox"/> Symptoms worse during the night or early morning <input type="checkbox"/> Symptoms triggered by exercise, emotions including laughter, dust or exposure to allergens 	<ul style="list-style-type: none"> <input type="checkbox"/> Persistence of symptoms despite treatment <input type="checkbox"/> Good and bad days but always daily symptoms and exertional dyspnea <input type="checkbox"/> Chronic cough and sputum preceded onset of dyspnea, unrelated to triggers
Lung function	Current and/or historical variable airflow limitation, e.g. BD reversibility, AHR	FEV ₁ may be improved by therapy, but post-BD FEV ₁ /FVC < 0.7 persists	Airflow limitation not fully reversible, but often with current or historical variability	<ul style="list-style-type: none"> <input type="checkbox"/> Record of variable airflow limitation (spirometry, peak flow) 	<ul style="list-style-type: none"> <input type="checkbox"/> Record of persistent airflow limitation (post-bronchodilator FEV₁/FVC < 0.7)
Lung function between symptoms	May be normal between symptoms	Persistent airflow limitation	Persistent airflow limitation	<ul style="list-style-type: none"> <input type="checkbox"/> Lung function normal between symptoms 	<ul style="list-style-type: none"> <input type="checkbox"/> Lung function abnormal between symptoms
Past history or family history	Many patients have allergies and a personal history of asthma in childhood, and/or family history of asthma	History of exposure to noxious particles and gases (mainly tobacco smoking and biomass fuels)	Frequently a history of doctor-diagnosed asthma (current or previous), allergies and a family history of asthma, and/or a history of noxious exposures	<ul style="list-style-type: none"> <input type="checkbox"/> Previous doctor diagnosis of asthma <input type="checkbox"/> Family history of asthma, and other allergic condition 	<ul style="list-style-type: none"> <input type="checkbox"/> Previous doctor diagnosis of COPD, chronic bronchitis or emphysema <input type="checkbox"/> Heavy exposure to a risk factor: tobacco smoke, biomass fuels
Time course	Often improves spontaneously or with treatment, but may result in fixed airflow limitation	Generally, slowly progressive over years despite treatment	Symptoms are partly but significantly reduced by treatment. Progression is usual and treatment needs are high	<ul style="list-style-type: none"> <input type="checkbox"/> No worsening of symptoms over time. Symptoms vary either seasonally, or from year to year <input type="checkbox"/> May improve spontaneously or have an immediate response to BD or to ICS over weeks 	<ul style="list-style-type: none"> <input type="checkbox"/> Symptoms slowly worsening over time (progressive course over years) <input type="checkbox"/> Rapid-acting bronchodilator treatment provides only limited relief.
Chest X-ray	Usually normal	Severe hyperinflation & other changes of COPD	Similar to COPD	<ul style="list-style-type: none"> <input type="checkbox"/> Normal 	<ul style="list-style-type: none"> <input type="checkbox"/> Severe hyperinflation
Exacerbations	Exacerbations occur, but the risk of exacerbations can be considerably reduced by treatment	Exacerbations can be reduced by treatment. If present, comorbidities contribute to impairment	Exacerbations may be more common than in COPD but are reduced by treatment. Comorbidities can contribute to impairment	<p>*Syndromic diagnosis of airways disease: how to use Table 2b Shaded columns list features that, when present, best distinguish between asthma and COPD. For a patient, count the number of checked boxes in each column. If three or more boxes are checked for either asthma or COPD, that diagnosis is suggested. If there are similar numbers of checked boxes in each column, the diagnosis of ACOS should be considered. See Step 2 for more details.</p>	
Typical airway inflammation	Eosinophils and/or neutrophils	Neutrophils in sputum, lymphocytes in airways, may have systemic inflammation	Eosinophils and/or neutrophils in sputum.		

STEP 3: Spirometry

Spirometry is essential for the assessment of patients with suspected chronic disease of the airways. It must be performed at either the initial or a subsequent visit, if possible before and after a trial of treatment. Early confirmation or exclusion of the diagnosis may avoid needless trials of therapy, or delays in initiating other investigations. Spirometry confirms chronic airflow limitation but is of more limited value in distinguishing between asthma with fixed airflow obstruction, COPD and ACOS (Table 3).

Measurement of peak expiratory flow (PEF), although not an alternative to spirometry, if performed repeatedly on the same meter over a period of 1–2 weeks may help to confirm the diagnosis of asthma by demonstrating excessive variability, but a normal PEF does not rule out either asthma or COPD. A high level of variability in lung function may also be found in ACOS.

After the results of spirometry and other investigations are available, the provisional diagnosis from the syndrome-based assessment must be reviewed and, if necessary, revised. As shown in Table 3, spirometry at a single visit is not always confirmatory of a diagnosis, and results must be considered in the context of the clinical presentation, and whether treatment has been commenced. Inhaled corticosteroids and long-acting bronchodilators influence results, particularly if a long withhold period is not used prior to performing spirometry. Further tests might therefore be necessary either to confirm the diagnosis or to assess the response to initial and subsequent treatment.

STEP 4: Commence initial therapy

Faced with a differential diagnosis equally balanced between asthma and COPD (i.e. ACOS) the default position should be to start treatment accordingly for asthma (Table 4). This recognizes the pivotal role of ICS in preventing morbidity and even death in patients with uncontrolled asthma symptoms, for whom even seemingly 'mild' symptoms (compared to those of moderate or severe COPD) might indicate significant risk of a life-threatening attack¹⁰.

- If the syndromic assessment suggests asthma or ACOS, or there is significant uncertainty about the diagnosis of COPD, it is prudent to start treatment as for asthma until further investigation has been performed to confirm or refute this initial position.
 - o Treatments will include an ICS (in a low or moderate dose, depending on level of symptoms).
 - o A long-acting beta2-agonist (LABA) should also be continued (if already prescribed), or added. However, it is important that patients should not be treated with a LABA without an ICS (often called LABA monotherapy) if there are features of asthma.
- If the syndromic assessment suggests COPD, appropriate symptomatic treatment with bronchodilators or combination therapy should be commenced, but not ICS alone (as monotherapy).²¹
- Treatment of ACOS should also include advice about other therapeutic strategies¹⁶ including:
 - o Smoking cessation
 - o Pulmonary rehabilitation
 - o Vaccinations
 - o Treatment of comorbidities, as advised in the respective GINA and GOLD reports.

In a majority of patients, the initial management of asthma and COPD can be satisfactorily carried out at primary care level. However, both the GINA and GOLD strategy reports make provision for referral for further diagnostic procedures at relevant points in patient management (see Step 5). This may be particularly important for patients with suspected ACOS, given that it is associated with worse outcomes and greater health care utilization.

Table 3. Spirometric measures in asthma, COPD and ACOS

Spirometric variable	Asthma	COPD	ACOS
Normal FEV ₁ /FVC pre- or post BD	Compatible with diagnosis	Not compatible with diagnosis	Not compatible unless other evidence of chronic airflow limitation
Post-BD FEV ₁ /FVC <0.7	Indicates airflow limitation but may improve spontaneously or on treatment	Required for diagnosis (GOLD)	Usually present
FEV ₁ ≥80% predicted	Compatible with diagnosis (good asthma control or interval between symptoms)	Compatible with GOLD classification of mild airflow limitation (categories A or B) if post- BD FEV ₁ /FVC <0.7	Compatible with diagnosis of mild ACOS
FEV ₁ <80% predicted	Compatible with diagnosis. Risk factor for asthma exacerbations	An indicator of severity of airflow limitation and risk of future events (e.g. mortality and COPD exacerbations)	An indicator of severity of airflow limitation and risk of future events (e.g. mortality and exacerbations)
Post-BD increase in FEV ₁ >12% and 200 ml from baseline (reversible airflow limitation)	Usual at some time in course of asthma, but may not be present when well-controlled or on controllers	Common and more likely when FEV ₁ is low, but ACOS should also be considered	Common and more likely when FEV ₁ is low, but ACOS should also be considered
Post-BD increase in FEV ₁ >12% and 400ml from baseline (marked reversibility)	High probability of asthma	Unusual in COPD. Consider ACOS	Compatible with diagnosis of ACOS

ACOS: asthma-COPD overlap syndrome; BD: bronchodilator; FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; GOLD: Global Initiative for Obstructive Lung Disease.

STEP 5: Referral for specialized investigations (if necessary)

Referral for expert advice and further diagnostic evaluation is necessary in the following contexts:

- Patients with persistent symptoms and/or exacerbations despite treatment.
- Diagnostic uncertainty, especially if an alternative diagnosis (e.g. bronchiectasis, post-tuberculous scarring, bronchiolitis, pulmonary fibrosis, pulmonary hypertension, cardiovascular diseases and other causes of respiratory symptoms) needs to be excluded.
- Patients with suspected asthma or COPD in whom atypical or additional symptoms or signs (e.g. haemoptysis, significant weight loss, night sweats, fever, signs of bronchiectasis or other structural lung disease) suggest an additional pulmonary diagnosis. This should prompt early referral, without necessarily waiting for a trial of treatment for asthma or COPD.
- When chronic airways disease is suspected but syndromic features of both asthma and COPD are few.
- Patients with comorbidities that may interfere with the assessment and management of their airways disease.
- Referral may also be appropriate for issues arising during on-going management of asthma, COPD or ACOS, as outlined in the GINA and GOLD strategy reports.

Table 5 summarizes specialized investigations that may be used to distinguish asthma and COPD.

Table 4. Summary of syndromic approach to diseases of chronic airflow limitation




Table 5. Specialized investigations sometimes used in distinguishing asthma and COPD

	Asthma	COPD
Lung function tests		
DLCO	Normal (or slightly elevated).	Often reduced.
Arterial blood gases	Normal between exacerbations	May be chronically abnormal between exacerbations in more severe forms of COPD
Airway hyperresponsiveness (AHR)	Not useful on its own in distinguishing asthma from COPD, but high levels of AHR favor asthma	
Imaging		
High resolution CT Scan	Usually normal but air trapping and increased bronchial wall thickness may be observed.	Low attenuation areas denoting either air trapping or emphysematous change can be quantitated; bronchial wall thickening and features of pulmonary hypertension may be seen.
Inflammatory biomarkers		
Test for atopy (specific IgE and/or skin prick tests)	Modestly increases probability of asthma; not essential for diagnosis	Conforms to background prevalence; does not rule out COPD
FENO	A high level (>50 ppb) in non-smokers supports a diagnosis of eosinophilic airway inflammation	Usually normal. Low in current smokers.
Blood eosinophilia	Supports asthma diagnosis	May be present during exacerbations
Sputum inflammatory cell analysis	Role in differential diagnosis is not established in large populations	

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

REFERENCES

1. Guerra S, Sherrill DL, Kurzius-Spencer M, et al. The course of persistent airflow limitation in subjects with and without asthma. *Respiratory Medicine* 2008;102:1473-82.
2. Silva GE, Sherrill DL, Guerra S, Barbee RA. Asthma as a risk factor for COPD in a longitudinal study. *Chest* 2004;126:59-65.
3. van Schayck CP, Levy ML, Chen JC, Isonaka S, Halbert RJ. Coordinated diagnostic approach for adult obstructive lung disease in primary care. *Prim Care Respir J* 2004;13:218-21.
4. Zeki AA, Schivo M, Chan A, Albertson TE, Louie S. The Asthma-COPD Overlap Syndrome: A Common Clinical Problem in the Elderly. *J Allergy* 2011;2011:861926.
5. Abramson MJ, Schattner RL, Sulaiman ND, Del Colle EA, Aroni R, Thien F. Accuracy of asthma and COPD diagnosis in Australian general practice: a mixed methods study. *Prim Care Respir J* 2012;21:167-73.
6. Gibson PG, Simpson JL. The overlap syndrome of asthma and COPD: what are its features and how important is it? *Thorax* 2009;64:728-35.
7. Mannino DM, Gagnon RC, Petty TL, Lydick E. Obstructive lung disease and low lung function in adults in the United States: data from the National Health and Nutrition Examination Survey, 1988-1994. *Arch Intern Med* 2000;160:1683-9.
8. Marsh SE, Travers J, Weatherall M, et al. Proportional classifications of COPD phenotypes. *Thorax* 2008;63:761-7.
9. Shirtcliffe P, Marsh S, Travers J, Weatherall M, Beasley R. Childhood asthma and GOLD-defined chronic obstructive pulmonary disease. *Intern Med J* 2012;42:83-8.
10. Louie S, Zeki AA, Schivo M, et al. The asthma-chronic obstructive pulmonary disease overlap syndrome: pharmacotherapeutic considerations. *Expert Rev Clin Pharmacol* 2013;6:197-219.
11. Miravittles M, Soler-Cataluna JJ, Calle M, Soriano JB. Treatment of COPD by clinical phenotypes: putting old evidence into clinical practice. *Eur Respir J* 2013;41:1252-6.
12. Soler-Cataluna JJ, Cosio B, Izquierdo JL, et al. Consensus document on the overlap phenotype COPD-asthma in COPD. *Arch Bronconeumol* 2012;48:331-7.
13. Kauppi P, Kupiainen H, Lindqvist A, et al. Overlap syndrome of asthma and COPD predicts low quality of life. *J Asthma* 2011;48:279-85.
14. Andersen H, Lampela P, Nevanlinna A, Saynajakangas O, Keistinen T. High hospital burden in overlap syndrome of asthma and COPD. *Clin Respir J* 2013;7:342-6.
15. Weatherall M, Travers J, Shirtcliffe PM, et al. Distinct clinical phenotypes of airways disease defined by cluster analysis. *Eur Respir J* 2009;34:812-8.
16. McDonald VM, Simpson JL, Higgins J, Gibson PG. Multidimensional assessment of older people with asthma and COPD: clinical management and health status. *Age Ageing* 2011;40:42-9.
17. Soriano JB, Davis KJ, Coleman B, Visick G, Mannino D, Pride NB. The proportional Venn diagram of obstructive lung disease: two approximations from the United States and the United Kingdom. *Chest* 2003;124:474-81.
18. Carolan BJ, Sutherland ER. Clinical phenotypes of chronic obstructive pulmonary disease and asthma: recent advances. *J Allergy Clin Immunol* 2013;131:627-34; quiz 35.
19. Hardin M, Silverman EK, Barr RG, et al. The clinical features of the overlap between COPD and asthma. *Respir Res* 2011;12:127.
20. Wardlaw AJ, Silverman M, Siva R, Pavord ID, Green R. Multi-dimensional phenotyping: towards a new taxonomy for airway disease. *Clin Exp Allergy* 2005;35:1254-62.
21. Global Initiative for Chronic Obstructive Pulmonary Disease (GOLD). Global Strategy for Diagnosis, Management and Prevention of COPD. 2014.
22. Halbert RJ, Isonaka S. International Primary Care Respiratory Group (IPCRG) Guidelines: integrating diagnostic guidelines for managing chronic respiratory diseases in primary care. *Prim Care Respir J* 2006;15:13-9.
23. Levy ML, Fletcher M, Price DB, Hausen T, Halbert RJ, Yawn BP. International Primary Care Respiratory Group (IPCRG) Guidelines: diagnosis of respiratory diseases in primary care. *Prim Care Respir J* 2006;15:20-34.
24. Price DB, Tinkelman DG, Halbert RJ, et al. Symptom-based questionnaire for identifying COPD in smokers. *Respiration* 2006;73:285-95.

25. Thiadens HA, de Bock GH, Dekker FW, et al. Identifying asthma and chronic obstructive pulmonary disease in patients with persistent cough presenting to general practitioners: descriptive study. *BMJ* 1998;316:1286-90.
26. Tinkelman DG, Price DB, Nordyke RJ, et al. Symptom-based questionnaire for differentiating COPD and asthma. *Respiration* 2006;73:296-305.
27. Van Schayck CP, Loozen JM, Wagena E, Akkermans RP, Wesseling GJ. Detecting patients at a high risk of developing chronic obstructive pulmonary disease in general practice: cross sectional case finding study. *BMJ* 2002;324:1370.

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

NOTES

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

NOTES

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

NOTES

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

NOTES

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE

COPYRIGHTED MATERIAL-DO NOT ALTER OR REPRODUCE